

Spinal hemangioblastoma arising from cervical nerve root


Jiuhong Li, MD,* Jiaojiang He, MD,* Susu Lu, MD, Xuhui Hui, PhD, and Haifeng Chen, MD

Neurology® 2020;95:791-792. doi:10.1212/WNL.0000000000010836

Correspondence

Dr. Chen
chenhaifengwc@163.com

Figure 1 MRI findings



Preoperative imaging revealed a hypointensity mass in sagittal T1 (A) and hyperintensity in T2-weighted imaging (B). The lesion was cystic-solid in sagittal (C) and coronal (D) imaging after administration of gadolinium and axial enhanced MRI (E) showed it was extramedullary. Postoperative contrast-enhanced MRI (F) depicted the tumor was totally resected.


A 39-year-old woman presented with neck and shoulder pain for 3 months. The MRI scan revealed a cystic-solid lesion located at C5 (figure 1, A–E). The patient received an operation and a small blood blister-like soft mass was detected. The tumor originated from proximal cervical nerve root and compressed the spinal medulla (figure 2, A–C). Pathologic diagnosis was hemangioblastoma (figure 2D). Postoperative gadolinium-enhanced MRI showed a gross total resection of the tumor (figure 1F). Spinal hemangioblastomas frequently originate from the medulla,¹ and nerve root originated hemangioblastoma is scarce. Our case provides a direct-viewing description and pathologic confirmation of a new subset of classification for origin of spinal hemangioblastomas.

*These authors contributed equally to this work.

From the Departments of Neurosurgery (J.L., J.H., X.H., H.C.) and Pathology (S.L.), West China Hospital of Sichuan University, Chengdu.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Figure 2 Operative and pathologic findings



During operation, a soft blood blister-like mass was detected (A). After carefully dissecting it, we found that the lesion arose from the proximal cervical nerve root (arrow), compressing the spinal medulla (asterisks) (B). The tumor was totally resected so that the nerve root (arrow) and spinal medulla (asterisks) were revealed (C). Hematoxylin & eosin (magnification, $\times 200$) (D) showed large intratumoral vascular channels, loose stromal elements, and prominent capillaries characteristic of a hemangioblastoma; the nerve root tissue (asterisks) is surrounded by tumor stroma.

Study funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

Appendix Authors

Name	Location	Contribution
Jiuhong Li	Chengdu, Sichuan, China	Study design, data collection and analysis, drafting the manuscript, analysis and interpretation of data
Jiaojiang He	Chengdu, Sichuan, China	Data collection, drafting the manuscript, revising the manuscript

Appendix (continued)

Name	Location	Contribution
Susu Lu	Chengdu, Sichuan, China	Acquisition of data, analysis and interpretation of data, drafting the manuscript
Xuhui Hui	Chengdu, Sichuan, China	Study concept and design, study supervision, revising the manuscript
Haifeng Chen	Chengdu, Sichuan, China	In charge of surgery, study concept and design, clinical care and investigative workup of the patient

Reference

- Westwick HJ, Giguere JF, Shamji MF. Incidence and prognosis of spinal hemangioblastoma: a surveillance epidemiology and end results study. *Neuroepidemiology* 2016;46:14–23.

Neurology®

Spinal hemangioblastoma arising from cervical nerve root

Jiuhong Li, Jiaojiang He, Susu Lu, et al.

Neurology 2020;95:791-792 Published Online before print September 15, 2020

DOI 10.1212/WNL.0000000000010836

This information is current as of September 15, 2020

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/95/17/791.full
References	This article cites 1 articles, 0 of which you can access for free at: http://n.neurology.org/content/95/17/791.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Education http://n.neurology.org/cgi/collection/all_education All Spinal Cord http://n.neurology.org/cgi/collection/all_spinal_cord MRI http://n.neurology.org/cgi/collection/mri Nerve tumor http://n.neurology.org/cgi/collection/nerve_tumor Spinal cord tumor http://n.neurology.org/cgi/collection/spinal_cord_tumor
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2020 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

