Identification of a pathogenic intronic KIF5A mutation in an ALS-FTD kindred

Sara Saez-Atienzar, PhD, Clifton L. Dalgard, PhD, Jinhui Ding, PhD, Adriano Chiò, MD, Camile Alba, PhD, Dan N. Hupalo, PhD, Matthew D. Wilkerson, PhD, Robert Bowser, PhD, Erik P. Pioro, MD, PhD, Richard Bedlack, MD, PhD, and Bryan J. Traynor, MD, PhD

Correspondence

Dr. Traynor
traynorb@mail.nih.gov

Neurology® 2020;95:1015-1018. doi:10.1212/WNL.00000000000011064

Not every gene nominated as a cause of human disease stands the test of time. As additional data become available, the evidence supporting the pathogenicity of a particular variant within a gene can be enhanced or diminished.¹ The amyotrophic lateral sclerosis (ALS) field, as much as any other, has been hesitant to address these controversies, leading to uncertainty among the research community.

In 2013, we published a study reporting that mutations in the *MATR3* gene were a cause of familial ALS.² That study was based, in part, on a pedigree in which we described p.Phe115Cys as the pathogenic variant based on exome sequence data obtained from 4 affected individuals. An additional member of this kindred (known as USALS#3, member III:10) was recently diagnosed as having ALS. Clinical genetic testing of this individual showed that they did not carry the *MATR3* mutation. Although this individual may be a phenocopy, a more parsimonious explanation was that a different mutation was responsible.

To address this issue, we performed whole-genome sequencing of this amyotrophic lateral sclerosis-frontotemporal dementia (ALS-FTD) family on an Illumina NovaSeq6000 sequencer to identify their true causative mutation (figure 1 and table). The participating institutions' institutional review boards approved the study (clinicaltrials.gov/ct2/show/NCT02014246), and informed consent was obtained from all subjects or their surrogate decision makers, according to the Declaration of Helsinki.

Analysis of the sequence data identified 218 variants that were rare and shared across the 5 affected individuals. One variant was located within intron 26 of the *KIF5A* gene, 14 base pairs from the start of exon 27 (chr12:57582588G>T, build hg38). Exon 27 within *KIF5A* is a known mutational hotspot underlying familial ALS.³ Exon trap experiments on cDNA obtained from our proband confirmed that this intronic mutation led to aberrant splicing of the *KIF5A* mRNA transcript. The altered transcript sequence was identical to that produced by other mutations in this intronic region because of skipping of exon 27 (figure 2).³ This family represents the most extensive kindred ascribed to a *KIF5A* mutation to date, and the affected individuals display both the short survival typically associated with ALS and the prolonged survival previously observed among some patients carrying mutations in this gene (table).³

Discussion

Our previous publication erroneously nominated the p.Phe115Cys variant in *MATR3* as the cause of disease within the USALS#3 kindred based on exome sequencing of affected

From the Neuromuscular Diseases Research Section (S.S.-A., B.J.T.), Laboratory of Neurogenetics, NIA, NIH; Department of Anatomy (C.L.D.), Physiology & Genetics, Uniformed Services University of the Health Sciences; The American Genome Center (C.L.D., C.A., D.N.H., M.D.W.), Collaborative Health Initiative Research Program, Uniformed Services University of the Health Sciences; Computational Biology Core (J.D.), Laboratory of Neurogenetics, NIA, NIH, Bethesda, MD; Rita Levi Montalcini Department of Neuroscience (A.C.), University of Turin, Italy and AOU Città della Salute e della Scienza, Turin, Italy; Division of Neurology and Neurobiology (R. Bowser), Barrow Neurological Institute, Phoenix, AZ; Department of Neurology (E.P.P.), Neuromuscular Center, Neurological Institute, Cleveland Clinic, OH; Department of Neurology (R. Bedlack), Duke University and the Durham VA Medical Center, NC; and Department of Neurology (B.J.T.), Johns Hopkins University, Baltimore, MD.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

The Article Processing Charge was funded by NIH.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Figure 1 Intronic mutation in *KIF5A* identified in an ALS-FTD kindred

The upper panel depicts a lollipop plot of *KIF5A* depicting the location of the intronic mutation identified in the USALS#3 kindred (red) and known mutations (gray). The chromatograms show mutant alleles from the indicated individuals (mutation is highlighted in pale pink). The pedigree shows the ALS-FTD kindred carrying the *KIF5A* intronic mutation (updated from ref 5). Individuals II:6, II:7, III:1, III:9, III:10, III:11 were diagnosed with ALS by neurologists. In addition, III:7 had executive dysfunction consistent with behavioral FTD based on formal neuropsychological testing, whereas III:1 and III:11 were observed to have mild cognitive impairment. The phenomenology of other individuals was reconstructed based on history from family members. Mt = mutant alleles; wt = wild-type alleles. Genotypes of presumed obligate carriers are in brackets. Arrow denotes proband. ALS = amyotrophic lateral sclerosis.

individuals.² Here, we correct the record to show that an intronic mutation within the known mutational hotspot of *KIF5A* is the actual cause of disease within this ALS-FTD family. The availability of DNA from an additional affected member within this pedigree was vital to identifying the causative mutation correctly. However, advancements within the genomics field and our understanding of ALS genetics were similarly crucial to resolving this family. In particular, our preexisting knowledge concerning *KIF5A* allowed us to single out that variant from the list of shared variants.³

Seven members of the kindred (figure 1 and table) developed executive dysfunction during their ALS illness, demonstrating a link between mutations in *KIF5A* and FTD. Mutations in *KIF5A* have now been linked to a wide variety of neurodegenerative conditions, including hereditary spastic paraparesis,⁴ Charcot-Marie-Tooth disease,⁴ ALS,³ and, more recently, the

KIF5A protein has been implicated as having a role in Alzheimer disease.⁵ These discoveries show the importance of the kinesin protein complex and axonal transport within neurons. Aside from being a striking example of pleiotropy within a single gene, it also suggests that, cumulatively, mutations within *KIF5A* may be a significant cause of neurologic disease.

Despite our recent findings, we maintain that mutations in *MATR3* are a cause of familial ALS. The p.Ser85Cys variant in *MATR3* remains the cause of neurologic disease within the other pedigree (USALS#4), segregating with disease among 11 affected members across multiple generations.² Although there is clear muscle involvement within this family, there is clinical evidence of upper and lower motor neuron involvement.² *MATR3* protein is present within neuronal cytoplasmic inclusions of more than half of sporadic ALS patients,⁶ and pathogenic *MATR3* mutants

Table Clinical features of affected individuals in the USALS#3 kindred

Individual	Clinical features			
	Diagnosis	Site of onset	Age at onset	Survival
I:1	ALS	Lower limb	Died at age 47	Prolonged course
II:1	Dementia	Cognition	NA	NA
II:2	ALS	NA	NA	NA
II:3	Dementia	Cognition	Died at age 84	NA
II:5	ALS-FTD	Cognition	NA	NA
II:6	ALS-FTD	Upper limb	70	5 y
II:7	ALS-FTD	Upper limb	57	26 y
III:1	ALS-FTD	Lower limb	63	5 y
III:9	ALS	Upper limb	52	8 y
III:10	ALS	Limb	63	1.5 y
III:11	ALS-FTD	Bulbar	50	6 y

Abbreviation: ALS = amyotrophic lateral sclerosis.

display neurotoxicity that is mitigated by cytoplasmic redistribution.⁷ Motor neuron loss and gliosis have been observed within the spinal cords of transgenic mice over-expressing mutant p.Ser85Cys MATR3.⁸ Finally, there are reports of other *MATR3* mutations in patients diagnosed with ALS.⁹

In conclusion, we identified an intronic mutation in *KIF5A* that segregated with disease in a large, multigenerational pedigree. Our efforts highlight the rapid advancements that are taking place in our understanding of the genetic

architecture of ALS and link mutations in *KIF5A* to cognitive impairment/frontotemporal dementia.

Study funding

This work was supported by the Intramural Research Programs of the NIH, National Institute on Aging (Z01-AG000949-02). The work was also funded by the Packard Center for ALS Research at Hopkins, the ALS Association, the Muscular Dystrophy Association, the Italian Ministry of Health (RF-2016-02362405), the Italian Ministry of Education, University and Research (Progetti di Ricerca di Rilevante

Figure 2 The intronic mutation alters the splicing of *KIF5A*

(A) RNA derived from blood (upper panel) from a healthy individual, an ALS patient not carrying the mutation, and individual III-11 carrying the *KIF5A* intronic mutation. RT-PCR was performed using RNA and previously described primers to amplify a wild-type (155 bp) splice form extending from exon 26 to exon 28.³ An extra band was observed at 127 base pairs indicating aberrant splicing in individual III-11 that was not present in the healthy and disease control subjects. RNA obtained from an IPS cell line (lower panel) derived from fibroblasts of individual III-11 and a control IPS cell line (A18945) showed the same pattern. (B) Sanger sequence analysis of the 127bp transcript/band observed in the patient confirmed the skipping of exon 27 of *KIF5A* yielding an out of frame and extended disrupted C-terminal peptide sequence.³ ALS = amyotrophic lateral sclerosis.

Interesse Nazionale, PRIN, grant n. 2017SNW5MB), the Joint Programme - Neurodegenerative Disease Research (JPND, Brain-Mend projects) granted by Italian Ministry of Education, University and Research, and by the European Community's Health Seventh Framework Programme (FP7/2007–2013, grant agreements no. 259867 and 278611), by the National Institute of Neurologic Disorders and Stroke (NIH grant number R35 NS097261), and by the Collaborative Health Initiative Research Program. This study was performed under the Department of Excellence grant of the Italian Ministry of Education, University and Research to the “Rita Levi Montalcini” Department of Neuroscience, University of Torino, Italy. This study utilized the high-performance computational capabilities of the Biowulf Linux cluster at the NIH, Bethesda, Maryland (hpc.nih.gov). The institutional review boards of participating institutions approved the study (NIH, 03-AG-N329), and informed consent was obtained from all subjects or their surrogate decision-makers, according to the Declaration of Helsinki.

Disclosure

S. Saez-Atienzar—Reports no disclosures. C.L. Dalgard—Reports no disclosures. J. Ding—Reports no disclosures. A. Chiò—Scientific Advisory Board Member for: Biogen Idec, Cytokinetics, Italfarmaco and Mitsubishi Tanabe, and editorial board member for the journal Amyotrophic Lateral Sclerosis and Frontotemporal Dementia. C. Alba—Reports no disclosures. D.N. Hupalo—Reports no disclosures. M.D. Wilkerson—Reports no disclosures. R. Bowser—founder of Iron Horse Diagnostics, Inc, a company commercialising assays for ALS and other neurologic diseases. E.P. Pioro—Scientific Advisory Board Member for: Biohaven Pharmaceuticals, Cytokinetics, Inc., ITF Pharma, Inc., MT Pharma America, Inc., and Consultant to: Avanir Pharmaceuticals, Inc., Cytokinetics, Inc., ITF Pharma, Inc., MT Pharma America, Inc., Neurotherapia, Inc., and Otsuka America, Inc. R. Bedlack—Scientific Advisory Board Member for: Brainstorm Cell DSMB, Mallinkrodt DSMB, Biogen DSMB, editorial board member for the journal Amyotrophic Lateral Sclerosis and Frontotemporal Dementia, and consultant for the ALS Association, Biohaven, ITF Pharma, MT Pharma America, New Biotic Inc, Woolsey Pharma, Leerink Swan and Company, Easton Associates, and the Gerson Lehman Group. B.J. Traynor—holds US, EU and Canadian patents on the clinical testing and therapeutic intervention for the hexanucleotide repeat expansion of C9orf72, and editorial board member for the journals JAMA Neurology, Journal of Neurology, Neurosurgery and Psychiatry, and Neurobiology of Aging. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Publication history

Received by *Neurology* May 9, 2020. Accepted in final form September 21, 2020.

Appendix Authors

Name	Location	Contribution
Sara Saez-Atienzar, PhD	National Institute on Aging, Bethesda, MD	Data collection and analysis and drafting and revision of the manuscript
Clifton L. Dalgard, PhD	Uniformed Services University of the Health Sciences, Bethesda, MD	Data collection and analysis and drafting and revision of the manuscript
Jinhui Ding, PhD	National Institute on Aging, Bethesda, MD	Data collection and analysis and drafting and revision of the manuscript
Adriano Chiò, MD	University of Turin, Italy	Data collection and analysis and drafting and revision of the manuscript
Camile Alba, PhD	Uniformed Services University of the Health Sciences, Bethesda, MD	Data collection and analysis and drafting and revision of the manuscript
Dan N. Hupalo, PhD	Uniformed Services University of the Health Sciences, Bethesda, MD	Data collection and analysis and drafting and revision of the manuscript
Matthew D. Wilkerson, PhD	Uniformed Services University of the Health Sciences, Bethesda, MD	Data collection and analysis and drafting and revision of the manuscript
Robert Bowser, PhD	Barrow Neurologic Institute, Phoenix, AZ	Data collection and analysis and drafting and revision of the manuscript
Erik P. Pioro, MD, PhD	Cleveland Clinic, OH	Data collection and analysis and drafting and revision of the manuscript
Richard Bedlack, MD, PhD	Duke University, Durham, NC	Data collection and analysis and drafting and revision of the manuscript
Bryan J. Traynor, MD, PhD	National Institute on Aging, Bethesda, MD	Design and conceptualized study, analyzed the data, and drafted the manuscript for intellectual content

References

- Wang J, Shen Y. When a “disease-causing mutation” is not a pathogenic variant. *Clin Chem* 2014;60:711–713.
- Johnson JO, Pioro EP, Boehringer A, et al. Mutations in the Matrin 3 gene cause familial amyotrophic lateral sclerosis. *Nat Neurosci* 2014;17:664–666.
- Nicolas A, Kenna KP, Renton AE, et al. Genome-wide analyses identify KIF5A as a novel ALS gene. *Neuron* 2018;97:1268–1283.
- Liu YT, Laura M, Hersheson J, et al. Extended phenotypic spectrum of KIF5A mutations: from spastic paraplegia to axonal neuropathy. *Neurology* 2014;83:612–619.
- Wang Q, Tian J, Chen H, Du H, Guo L. Amyloid beta-mediated KIF5A deficiency disrupts anterograde axonal mitochondrial movement. *Neurobiol Dis* 2019;127:410–418.
- Tada M, Doi H, Koyano S, et al. Matrin 3 is a component of neuronal cytoplasmic inclusions of motor neurons in sporadic amyotrophic lateral sclerosis. *Am J Pathol* 2018;188:507–514.
- Malik AM, Miguez RA, Li X, Ho YS, Feldman EL, Barmada SJ. Matrin 3-dependent neurotoxicity is modified by nucleic acid binding and nucleocytoplasmic localization. *Elife* 2018;7:e35977.
- Zhang X, Yamashita S, Hara K, et al. Mutant MATR3 mouse model to explain multisystem proteinopathy. *J Pathol* 2019;249:182–192.
- Leblond CS, Gan-Or Z, Spiegelman D, et al. Replication study of MATR3 in familial and sporadic amyotrophic lateral sclerosis. *Neurobiol Aging* 2016;37:209.e17–e209.e21.

Neurology®

Identification of a pathogenic intronic KIF5A mutation in an ALS-FTD kindred

Sara Saez-Atienzar, Clifton L. Dalgard, Jinhui Ding, et al.

Neurology 2020;95;1015-1018 Published Online before print October 19, 2020

DOI 10.1212/WNL.0000000000011064

This information is current as of October 19, 2020

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/95/22/1015.full
References	This article cites 9 articles, 2 of which you can access for free at: http://n.neurology.org/content/95/22/1015.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Genetics http://n.neurology.org/cgi/collection/all_genetics Amyotrophic lateral sclerosis http://n.neurology.org/cgi/collection/amyotrophic_lateral_sclerosis_Frontotemporal_dementia Frontotemporal dementia http://n.neurology.org/cgi/collection/frontotemporal_dementia
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.