

# *C9orf72*, age at onset, and ancestry help discriminate behavioral from language variants in FTL D cohorts

Beatrice Costa, BSc, Claudia Manzoni, PhD, Manuel Bernal-Quiros, PhD, Demis A. Kia, MD, Miquel Aguilar, MD, Ignacio Alvarez, MSc, Victoria Alvarez, PhD, Ole Andreassen, MD, PhD, Maria Anfossi, PhD, Silvia Bagnoli, PhD, Luisa Benussi, PhD, Livia Bernardi, PhD, Giuliano Binetti, MD, Daniel Blackburn, MD, PhD, Mercè Boada, MD, PhD, Barbara Borroni, MD, Lucy Bowns, Geir Bråthen, MD, PhD, Amalia C. Bruni, MD, Huei-Hsin Chiang, PhD, Jordi Clarimon, PhD, Shuna Colville, MSc, Maria E. Conidi, PhD, Tom E. Cope, MD, PhD, Carlos Cruchaga, PhD, Chiara Cupidi, MD, Maria Elena Di Battista, MD, PhD, Janine Diehl-Schmid, MD, Monica Diez-Fairen, MSc, Oriol Dols-Icardo, PhD, Elisabetta Durante, PhD, Dušan Flisar, MD, Francesca Frangipane, MD, Daniela Galimberti, PhD, Maura Gallo, PhD, Maurizio Gallucci, MD, Roberta Ghidoni, PhD, Caroline Graff, MD, PhD, Jordan H. Grafman, PhD, Murray Grossman, MD, John Hardy, PhD, Isabel Hernández, MD, PhD, Guy J.T. Holloway, MBBS, Edward D. Huey, MD, Ignacio Illán-Gala, MD, PhD, Anna Karydas, MSc, Behzad Khoshnood, PhD, Milica G. Kramberger, MD, PhD, Mark Kristiansen, PhD, Patrick A. Lewis, PhD, Alberto Lleó, MD, PhD, Gaganjit K. Madhan, MSc, Raffaele Maletta, MD, Aleš Maver, MD, PhD, Manuel Menendez-Gonzalez, MD, PhD, Graziella Milan, MD, Bruce Miller, MD, Merel O. Mol, MSc, Parastoo Momeni, PhD, Sonia Moreno-Grau, PhD, Chris M. Morris, PhD, Benedetta Nacmias, PhD, Christer Nilsson, MD, Valeria Novelli, PhD, Linn Öjsterstedt, MD, Alessandro Padovani, MD, Suvankar Pal, MBBS, MRCP, MD, Yasmin Panchbhaya, MSc, Pau Pastor, MD, PhD, Borut Peterlin, MD, PhD, Irene Piaceri, PhD, Stuart Pickering-Brown, PhD, Yolande A.L. Pijnenburg, MD, PhD, Annibale A. Puca, MD, Innocenzo Rainero, MD, PhD, Antonella Rendina, PhD, Anna M.T. Richardson, FRCP, Ekaterina Rogaeva, PhD, Boris Rogelj, PhD, Sara Rollinson, PhD, Giacomina Rossi, PhD, Carola Rossmeyer, MD, James B. Rowe, MD, PhD, Elisa Rubino, MD, PhD, Agustín Ruiz, MD, PhD, Raquel Sanchez-Valle, MD, PhD, Sigrid B. Sando, PhD, Alexander F. Santillo, MD, PhD, Jennifer Saxon, MSc, Elio Scarpini, MD, Maria Serpente, PhD, Nicoletta Smirne, BSc, Sandro Sorbi, MD, EunRan Suh, PhD, Fabrizio Tagliavini, MD, Jennifer C. Thompson, PhD, John Q. Trojanowski, MD, PhD, Vivianna M. Van Deerlin, MD, PhD, Julie Van der Zee, PhD, Christine Van Broeckhoven, DSc, PhD, Jeroen van Rooij, John C. Van Swieten, MD, Arianna Veronesi, MD, PhD, Emilia Vitale, PhD, Maria L. Waldö, MD, PhD, Cathy Woodward, MSc, Jennifer Yokoyama, PhD, Valentina Escott-Price, PhD, James M. Polke, PhD, and Raffaele Ferrari, PhD, for the International FTD-Genetics Consortium

## Correspondence

Dr. Ferrari  
r.ferrari@ucl.ac.uk  
or Dr. Manzoni  
c.manzoni@ucl.ac.uk

*Neurology*® 2020;95:e3288-e3302. doi:10.1212/WNL.000000000010914

## Abstract

### Objective

We sought to characterize *C9orf72* expansions in relation to genetic ancestry and age at onset (AAO) and to use these measures to discriminate the behavioral from the language variant syndrome in a large pan-European cohort of frontotemporal lobar degeneration (FTLD) cases.

### Methods

We evaluated expansions frequency in the entire cohort ( $n = 1,396$ ; behavioral variant frontotemporal dementia [bvFTD] [ $n = 800$ ], primary progressive aphasia [PPA] [ $n = 495$ ], and FTLD–motor neuron disease [MND] [ $n = 101$ ]). We then focused on the bvFTD and PPA cases and tested for association between expansion status, syndromes, genetic ancestry, and AAO applying statistical tests comprising Fisher exact tests, analysis of variance with Tukey post hoc tests, and logistic and nonlinear mixed-effects model regressions.

### Results

We found *C9orf72* pathogenic expansions in 4% of all cases (56/1,396). Expansion carriers differently distributed across syndromes: 12/101 FTLD-MND (11.9%), 40/800 bvFTD (5%), and 4/495 PPA (0.8%). While addressing population substructure through principal components analysis (PCA), we defined 2 patients groups with Central/Northern ( $n = 873$ ) and Southern European ( $n = 523$ ) ancestry. The proportion of expansion carriers was significantly higher in bvFTD compared to PPA (5% vs 0.8% [ $p = 2.17 \times 10^{-5}$ ]; odds

ratio (OR) 6.4; confidence interval (CI) 2.31–24.99]), as well as in individuals with Central/Northern European compared to Southern European ancestry (4.4% vs 1.8% [ $p = 1.1 \times 10^{-2}$ ; OR 2.5; CI 1.17–5.99]). Pathogenic expansions and Central/Northern European ancestry independently and inversely correlated with AAO. Our prediction model (based on expansions status, genetic ancestry, and AAO) predicted a diagnosis of bvFTD with 64% accuracy.

## Conclusions

Our results indicate correlation between pathogenic *C9orf72* expansions, AAO, PCA-based Central/Northern European ancestry, and a diagnosis of bvFTD, implying complex genetic risk architectures differently underpinning the behavioral and language variant syndromes.

From the Institute of Neurology (B.C., D.A.K., J.H., P.A.L., R.F.), School of Pharmacy (C.M.), and UCL Movement Disorders Centre (J.H.), University College London; School of Pharmacy (C.M., P.A.L.), University of Reading, Whiteknights; Neurogenetics Laboratory (M.B.-Q., C.W., J.M.P.), National Hospital for Neurology and Neurosurgery, London, UK; Aptima Clinic (Miquel Aguilar), Terrassa; Memory Disorders Unit, Department of Neurology (I.A., M.D.-F., P.P.), University Hospital Mutua de Terrassa, Barcelona; Hospital Universitario Central de Asturias (V.A., M.M.-G.), Oviedo, Spain; NORMENT (O.A.), Institute of Clinical Medicine, University of Oslo, Norway; Regional Neurogenetic Centre (María Anfosso, Livia Bernardi, A.C.B., M.E.C., Chiara Cupidi, F.F., Maura Gallo, R.M., N.S.), ASPCZ, Lamezia Terme; Department of Neuroscience, Psychology, Drug Research and Child Health (S.B., B.N., I.P., S.S.), University of Florence; Molecular Markers Laboratory (Luisa Benussi, Giuliano Binetti, R.G.), IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy; Sheffield Institute for Translational Neuroscience (SITraN), Department of Neuroscience (D.B.), University of Sheffield, UK; Research Center and Memory Clinic (M.B., I.H., S.M.-G., Agustín Ruiz), Fundació ACE, Institut Català de Neurociències Aplicades, Universitat Internacional de Catalunya (UIC), Barcelona, Spain; Centre for Neurodegenerative Disorders (B.B., A.P.), Department of Clinical and Experimental Sciences, University of Brescia, Italy; Department of Clinical Neurosciences (Lucy Bowns, T.E.C., J.B.R.), Cambridge University, UK; Department of Neurology (Geir Bråthen, S.B.S.), University Hospital of Trondheim, Norway; Dept NVS, Division of Neurogeriatrics (H.-H.C., C.G., B.K., L.Ö.), Karolinska Institutet, Bioclinicum Solna, Sweden; Department of Neurology (J.C., O.D.-I., I.I.-G., A.L.), IIB Sant Pau, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Spain; Anne Rowling Regenerative Neurology Clinic (S.C., G.J.T.H., S.P.) and Centre for Clinical Brain Sciences (S.P.), University of Edinburgh, UK; NeuroGenomics and Informatics, Department of Psychiatry (Carlos Cruchaga), Washington University, St. Louis, MO; Cognitive Impairment Center (M.E.D.B., Maurizio Gallucci) and Immunohematology and Transfusional Medicine Service (E.D., A.V.), Local Health Authority n.2 Marca Trevigiana, Treviso, Italy; Department of Psychiatry and Psychotherapy (J.D.-S., C.R.), School of Medicine, Technical University of Munich, Germany; Department of Neurology (D.F., M.G.K.) and Clinical Institute of Medical Genetics (A.M., B.P.), University Medical Center Ljubljana, Slovenia; Dino Ferrari Center (D.G., Elio Scarpini, M.S.), University of Milan, Italy; Cognitive Neuroscience Lab, Think and Speak Lab (J.H.G.), Shirley Ryan Ability Lab, Chicago, IL; Department of Pathology and Laboratory Medicine (Murray Grossman, EunRan Suh, J.Q.T., V.M.V.D.), Center for Neurodegenerative Diseases, Perelman School of Medicine at the University of Pennsylvania, Philadelphia; UCL Dementia Research Institute (J.H.), London; Reta Lila Weston Institute (J.H.), UCL Queen Square Institute of Neurology, UK; Institute for Advanced Study (J.H.), The Hong Kong University of Science and Technology, China; Royal Edinburgh Hospital (G.J.T.H.), UK; Taub Institute for Research on Alzheimer's Disease and the Aging Brain (E.D.H.), Columbia University, New York, NY; Department of Neurology, Memory and Aging Center (A.K., B.M., J.Y.), University of California, San Francisco; UCL Genomics (M.K., G.K.M., Y.P.), UCL Great Ormond Street Institute of Child Health, London, UK; Geriatric Center Frullone ASL Napoli 1 Centro (G.M.), Napoli, Italy; Department of Neurology (M.O.M., J.v.R., J.C.V.S.), Erasmus Medical Center, Rotterdam, the Netherlands; Rona Holdings (P.M.), Silicon Valley, CA; Newcastle Brain Tissue Resource, Institute of Neuroscience (C.M.M.), Newcastle University, Campus for Ageing and Vitality, Newcastle upon Tyne, UK; Department of Neurology (C.N.), Skåne University Hospital, Malmö, Sweden; Fondazione Policlinico Universitario A. Gemelli IRCCS (V.N.), Rome, Italy; Division of Neuroscience & Experimental Psychology (S.P.-B., A.M.T.R., S.R., J.C.T.), University of Manchester, UK; Amsterdam University Medical Center (Y.A.L.P.), VU University Medical Center, the Netherlands; Cardiovascular Research Unit (A.A.P.), IRCCS Multimedica, Milan; Neurology I, Department of Neuroscience (I.R., Elisa Rubino), University of Torino; NeuROMICS laboratory (G.M., Antonella Rendina, E.V.), Institute of Biochemistry and Cell Biology (IBBC), CNR Napoli, Italy; Manchester Centre for Clinical Neurosciences (A.M.T.R., J.S., J.C.T.), Salford Royal NHS Trust, Manchester, UK; Tanz Centre for Research in Neurodegenerative Diseases (Ekaterina Rogava), University of Toronto, Canada; Department of Biotechnology (B.R.), Jožef Stefan Institute, Ljubljana, Slovenia; Division of Neurology V and Neuropathology (G.R., F.T.), Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy; Alzheimer's Disease and Other Cognitive Disorders Unit (R.S.-V.), Hospital Clinic of Barcelona, Spain; Clinical Memory Research Unit, Department of Clinical Sciences Malmö (C.N., A.F.S.), and Division of Clinical Sciences Helsingborg, Department of Clinical Sciences Lund (M.L.W.), Lund University, Sweden; Neurodegenerative Brain Diseases Group (J.V.d.Z., C.V.B.), Center for Molecular Neurology, VIB, Antwerp, Belgium; Medical Research Council Centre for Neuropsychiatric Genetics and Genomics (V.E.-P.), Division of Psychological Medicine and Clinical Neurosciences and Dementia Research Institute, Cardiff University, UK; Instituto de Investigación Sanitaria del Principado de Asturias (V.A.), Oviedo, Asturias; Fundació per la Recerca Biomèdica i Social Mútua Terrassa (I.A., M.D.-F., P.P.), Barcelona; Centro de Investigación Biomédica en Red de Enfermedades Neurodegenerativas (CIBERNED) (M.B., J.C., O.D.-I., I.H., I.I.-G., A.L., S.M.-G., Agustín Ruiz), Instituto de Salud Carlos III, Madrid, Spain; MRC Cognition and Brain Sciences Unit (Lucy Bowns, T.E.C., J.B.R.), Cambridge University, UK; Department of Neuromedicine and Movement Science (Geir Bråthen, S.B.S.), Norwegian University of Science and Technology, Trondheim, Norway; Unit for Hereditary Dementias (H.-H.C., C.G., B.K., L.Ö.), Theme Aging, Karolinska University Hospital, Solna, Sweden; Medical Faculty (D.F., M.G.K.), University of Ljubljana, Slovenia; Fondazione IRCCS Ca'Granda (D.G., Elio Scarpini, M.S.), Ospedale Policlinico, Milan, Italy; Penn Center for Frontotemporal Degeneration (Murray Grossman), Philadelphia, PA; Universidad de Oviedo (M.M.-G.), Asturias, Spain; IRCCS Fondazione Don Carlo Gnocchi (B.N., S.S.), Florence; Istituto di Medicina Genomica (V.N.), Università Cattolica del sacro Cuore, Rome, Italy; Amsterdam Neuroscience (Y.A.L.P.), the Netherlands; Department of Medicine and Surgery (A.A.P.), University of Salerno, Baronissi (SA), Italy; Faculty of Chemistry and Chemical Technology (B.R.), University of Ljubljana, Slovenia; Institut d'Investigacions Biomèdiques August Pi i Sunyer (R.S.-V.), Barcelona, Spain; Department of Biomedical Sciences (J.V.d.Z., C.V.B.), University of Antwerp, Belgium; and Department of Comparative Biomedical Sciences (P.A.L.), The Royal Veterinary College, London, UK.

International FTD-Genetics Consortium (IFGC) coinvestigators are listed at [links.lww.com/WNL/B240](https://links.lww.com/WNL/B240).

Go to [Neurology.org/N](https://Neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

The Article Processing Charge was funded by MRC and Wellcome Trust.

This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Glossary

AAO = age at onset; **bvFTD** = behavioral variant frontotemporal dementia; **FTLD** = frontotemporal lobar degeneration; **IFGC** = International FTD Genetics Consortium; **LOOCV** = leave-one-out cross-validation; **MND** = motor neuron disease; **PCA** = principal components analysis; **PNFA** = progressive nonfluent aphasia; **PPA** = primary progressive aphasia; **rc** = repeat counts; **SNP** = single nucleotide polymorphism.

Frontotemporal lobar degeneration (FTLD) refers to the second most common form of young-onset dementia after Alzheimer disease.<sup>1</sup> The major clinical syndromes are behavioral variant frontotemporal dementia (bvFTD)<sup>2</sup> or language dysfunctions, broadly called primary progressive aphasia (PPA); the latter is subdivided into semantic dementia or semantic variant PPA and progressive nonfluent aphasia (PNFA) or nonfluent/agrammatic variant PPA.<sup>2,3</sup> FTLD can also occur together with motor neuron disease (MND) or amyotrophic lateral sclerosis in a continuous spectrum of phenotypes.<sup>4</sup>

In FTLD, repeat expansions in *C9orf72*<sup>5</sup> have been previously reported to occur in ~25%<sup>6–10</sup> of familial and ~6%<sup>11</sup> of sporadic cases (i.e., individuals with no clear familial history or genetic aetiology<sup>12</sup>). Several studies had shown high frequencies of pathogenic *C9orf72* expansions in Northern vs Southern European patients (North–South axis), especially in historically isolated populations (such as the Finnish<sup>13,14</sup>), leading to the hypothesis that a Scandinavian founder might be at the basis of the spread of the *C9orf72* expansion.<sup>15</sup> Other studies (based on the geographic location of the recruiting sites) challenged the North–South axis concept, reporting a high frequency (~25%) of pathogenic expansions in the Spanish population<sup>10</sup> or implying to the existence of more than 1 risk haplotype.<sup>16–19</sup>

Patients with FTLD with abnormal *C9orf72* repeat expansions exhibit marked phenotypic and pathologic heterogeneity, suggesting presence of additional (genetic and environmental) modifiers.<sup>20</sup> Despite conflicting studies reporting either direct or inverse correlation between repeat length and age at onset (AAO), *C9orf72* expansions have been suggested to act as a genetic modifier of AAO.<sup>16,21–24</sup>

We analyzed 1,396 FTLD cases gathered through the International FTD Genetics Consortium (IFGC) ([ifgcsite.wordpress.com/](http://ifgcsite.wordpress.com/)) phase III initiative, aiming at (1) characterizing *C9orf72* expansions in relation to genetic ancestry and AAO and (2) assessing the usefulness of these measures in discriminating the behavioral from the language variant syndrome.

## Methods

### Standard protocol approvals, registrations, and patient consents

Each contributing site obtained written informed consent from all patients to be part of extended genetic studies; the

current study is approved under institutional review board approval 9811/001.

### Cohort, clinical phenotyping

FTLD cases were collected between 2016 and 2018 (within the IFGC phase III project [[ifgcsite.wordpress.com/ongoing-projects/](http://ifgcsite.wordpress.com/ongoing-projects/)]). The samples were recruited by clinicians and research groups who are part of the IFGC network and based in Italy, Spain, Germany, the Netherlands, Belgium, the United Kingdom, Sweden, Norway, Slovenia, or the United States (supplementary table 1, [doi.org/10.5522/04/12418157](https://doi.org/10.5522/04/12418157)). Patients were diagnosed at each contributing site (supplementary table 2, [doi.org/10.5522/04/12418157](https://doi.org/10.5522/04/12418157)) in a harmonized fashion according to international consensus criteria such as those of Neary et al.<sup>2</sup> (for FTLD), Rascovsky et al.<sup>25</sup> (for bvFTD), Gorno-Tempini et al.<sup>3</sup> (for PPA [semantic dementia or PNFA]), and Strong et al.<sup>4</sup> (for FTLD-MND).


### Genotyping, *C9orf72* repeat expansions, and analysis cohorts

A total of 1,454 cases were successfully genotyped by means of the NeuroArray<sup>26</sup> on the Illumina Infinium platform. Genotypes were used to inform on population substructure via standard principal components analysis (PCA) (supplementary figure 1, [doi.org/10.5522/04/12418157](https://doi.org/10.5522/04/12418157)), which led to the exclusion of 44 population outliers, and allowed us to address population substructure within the cohort (we identified 2 distinct [Nordic and Mediterranean] clusters; supplementary figure 2, [doi.org/10.5522/04/12418157](https://doi.org/10.5522/04/12418157)). We also assessed cryptic relatedness and excluded 14 first- or second-degree related individuals, leaving a cohort of 1,396 cases (group 0)—for which *C9orf72* expansion status (i.e., presence/absence of pathogenic expansions) was known—for analyses. Frequencies of pathogenic expansions were assessed in group 0 and further analyses were performed in (1) 1,295 cases (group 1: n = 800 bvFTD and n = 495 PPA) with known *C9orf72* expansion status; (2) 1,179 cases (group 2; n = 756 bvFTD and n = 423 PPA) with known *C9orf72* expansion status and AAO data available; and (3) 734 cases (group 3; n = 462 bvFTD and n = 272 PPA) with AAO and repeat counts (rc; screened via repeat-primed PCR) (see references 27 and 28; supplementary Methods and supplementary figure 3, [doi.org/10.5522/04/12418157](https://doi.org/10.5522/04/12418157); and figure 1A).

### Statistical analyses

We first assessed the frequency of pathogenic expansions in the entire cohort (group 0). The information on presence/absence of expansions was used as a binary variable (0 = absence of

**Figure 1** Cohorts and analysis workflow



(A) Cohorts. (B) Analysis workflow. K-fold regression model. AAO = age at onset; ANOVA = analysis of variance; bvFTD = behavioral variant frontotemporal dementia; FTL-D = frontotemporal lobar degeneration; logistic/cluster = logistic regression using cluster as covariate; logistic/PC1 = logistic regression using PC1 as covariate; LOOCV = leave-one-out cross-validation regression model; MND = motor neuron disease; PPA = primary progressive aphasia.

expansion; 1 = presence of expansion). We then investigated differences in the frequencies of pathogenic expansions across bvFTD and PPA and the Nordic and Mediterranean clusters in group 1 (Fisher exact test) and in group 3 (logistic regression); in the latter, we used rc as a categorical variable (using no, short, intermediate, and long as factor levels) considering the following 4 categories: no expansions (rc = 2/3), short expansions (4 ≤ rc ≤ 8), intermediate expansions (9 ≤ rc ≤ 24), and long expansions (rc ≥ 25), the latter representing expansions in the pathogenic range (see references 10 and 22, supplementary Methods, and supplementary figure 3, doi.org/10.5522/04/12418157).

We then evaluated association between AAO and syndrome, genetic ancestry, and expansions (i.e., presence/absence used as a binary variable; see above) alone and with genetic ancestry as a covariate in group 2 (t test and logistic regression) and in group 3 (t test, analysis of variance with Tukey post hoc test, and logistic and linear mixed-effects model). In the latter case, we used rc as a categorical variable (see above).

Finally, we sought to build a model to predict syndrome (bvFTD vs PPA) using (1) presence/absence of pathogenic expansions (as binary variable [see above] for group 2) or (2) rc (as categorical variable [see above] for group 3), ancestry as binary variable, and AAO as continuous variable using logistic regression models (i.e., the leave-one-out cross-validation

[LOOCV] and the K-fold models). A summary of the analyses workflow can be found in figure 1B.

All analyses were performed using R studio (version 3.6.0, studio version 1.2.1335).

### C9orf72 locus risk haplotype

Twenty (rs1110264, rs1110155, rs2150336, rs1161680, rs2589054, rs1822723, rs4879515, rs895023, rs868856, rs1977661, rs903603, rs12349820, rs10122902, rs2282241, rs1948522, rs1982915, rs2453556, rs702231, rs696826, and rs247751) of the original 42 single nucleotide polymorphisms (SNPs) constituting the (Finnish) risk haplotype<sup>29</sup> were available on the NeuroArray.<sup>26</sup> We filtered out 7 markers in order to keep 13 informative SNPs (rs1822723, rs4879515, rs868856, rs1977661, rs903603, rs10122902, rs2282241, rs1948522, rs1982915, rs2453556, rs702231, rs696826, and rs2477518) matching 13 of the 20 used in Mok et al.<sup>15</sup> We evaluated the proportion of cases carrying at least 1 risk allele (as in Mok et al.<sup>15</sup>) for each marker assessing expansion vs nonexpansion carriers (with/without ancestry stratification).

### Data availability

All data generated or analyzed during this study are included in this published article and supplementary files 1 and 2 at doi.org/10.5522/04/12418157.

## Results

### C9orf72 expansions frequency and syndromes

We assessed the frequency of pathogenic expansions in the entire cohort and across the different syndromes in the group 0 cases (figure 1). Four percent of all cases (56/1,396 [4%]) carried pathogenic expansions. These were most frequent in FTLD-MND (12/101 [11.9%]) followed by bvFTD (40/800 [5%]) and PPA (4/495 [0.8%]). The higher prevalence of pathogenic expansions in bvFTD vs PPA was statistically significant (Fisher exact test:  $p = 2.17 \times 10^{-5}$ ; OR 6.4; 95% CI 2.31–24.99, table 1). We further explored this finding in the group 3 cases using logistic regression to assess association between expansion length (represented by 4 rc factor levels: short, intermediate, and long expansions, tested against no expansions) and syndromes (bvFTD vs PPA). Expansion length discriminated bvFTD from PPA with a trend that was significant in the intermediate ( $p = 4.7 \times 10^{-2}$ ; OR 1.6; CI 0.0061 [2.5%]–0.94 [97.5%]) and long ( $p = 1.9 \times 10^{-3}$ ; OR 7.2; CI 0.86 [2.5%]–3.45 [97.5%]) rc ranges (with a ~90% probability of a bvFTD diagnosis supported by the latter; supplementary table 3, doi.org/10.5522/04/12418157).

### C9orf72 expansions (and rc) and genetic ancestry

We performed PCA (PC1 vs PC2, supplementary figure 2A; PC1 vs PC3, supplementary figure 2B, doi.org/10.5522/04/12418157) to cluster the group 1 cases based on their genetic makeup. There were 2 major clusters: cluster 1 (Mediterranean) included most of the cases (439/500 [87.8%]) recruited from Southern European sites (Italy and Spain); cluster 2 (Nordic) included most of the cases (627/795 [78.8%]) recruited from Central and Northern European sites (Belgium, the Netherlands, Germany, the United Kingdom, Norway, and Sweden). Samples recruited from Eastern European (Slovenia) and North American sites distributed across both clusters, although with a higher prevalence within cluster 2 (167/795 [21%]) vs cluster 1 (42/500 [8.4%]).

**Table 1** Frequency of expansion carriers in the entire cohort and by syndrome

Cohort	Cases, n	Expansion carriers	Frequency, %
<b>bvFTD</b>	800	40	5 <sup>a</sup>
<b>PPA</b>	495	4	0.8 <sup>a</sup>
<b>FTLD-MND</b>	101	12	11.9
<b>Total</b>	1,396	56	4

Abbreviations: bvFTD = behavioral variant frontotemporal dementia; FTLD = frontotemporal lobar degeneration; MND = motor neuron disease; PPA = primary progressive aphasia.

Summary of expansion carrier frequency in the entire cohort (n = 1,396) and across syndromes. The higher prevalence of expansion carriers in bvFTD vs PPA is statistically significant: <sup>a</sup>Fisher exact test performed to statistically evaluate the difference between the occurrence of pathogenic expansions in bvFTD vs PPA:  $p = 2.17 \times 10^{-5}$ ; odds ratio 6.4; 95% confidence interval 2.31–24.99.

**Table 2** Frequency of expansion carriers in the Nordic and Mediterranean clusters

Genetic ancestry	Cases, n	Expansion carriers	Frequency, %
<b>Nordic</b>	795	35	4.4 <sup>a</sup>
<b>Mediterranean</b>	500	9	1.8 <sup>a</sup>

The higher prevalence of expansion carriers in the Nordic vs the Mediterranean cluster is statistically significant: <sup>a</sup>Fisher exact test:  $p = 1.1 \times 10^{-2}$ ; odds ratio 2.5; 95% confidence interval 1.17–5.99.

We observed a significantly higher prevalence of pathogenic expansions in the Nordic (35/795 [4.4%]) vs the Mediterranean (9/500 [1.8%]) cluster (Fisher exact test:  $p = 1.1 \times 10^{-2}$ ; OR 2.5; CI 1.17–5.99, table 2). We further evaluated this finding in the group 3 cases using logistic regression to assess association between expansion length (see above) and genetic ancestry. Expansion length discriminated the Nordic from Mediterranean cluster with a trend that was significant in the intermediate ( $p = 9.7 \times 10^{-4}$ , OR 2.2; CI 0.32 [2.5%]–1.25 [97.5%]) and long ( $p = 4.7 \times 10^{-4}$ , OR 9.3; CI 1.12 [2.5%]–3.7 [97.5%]) rc ranges (with a ~90% probability of Nordic ancestry supported by the latter; supplementary table 4, doi.org/10.5522/04/12418157).

Provided differences in syndromes prevalence and distribution across the Nordic and Mediterranean clusters—bvFTD (469/795 [59%] vs 331/500 [66.2%]) and PPA (326/795 [41%] vs 169/500 [33.8%]), respectively (supplementary table 5, doi.org/10.5522/04/12418157)—we analyzed the distribution of pathogenic expansions across syndromes and clusters. Stratified Fisher exact test showed significant differences in the distribution of the pathogenic expansions between bvFTD and PPA in the Nordic (but not the Mediterranean) cluster ( $p = 1 \times 10^{-4}$ ; OR 7.87; 95% CI 2.43–40.52), and between the Nordic and the Mediterranean clusters for the bvFTD (but not PPA) syndrome ( $p = 1.9 \times 10^{-2}$ ; OR 2.95; 95% CI 1.31–7.52), suggesting that ancestry (Nordic) and syndrome (bvFTD) are independently associated with pathogenic expansions (table 3).

### C9orf72 repeat expansions (and counts [rc]) and AAO

We assessed AAO in the group 2 cases (figure 1). Mean AAO was significantly different between the bvFTD (61.7) and PPA (64) syndromes ( $t$  test:  $p = 1.86 \times 10^{-5}$ ; CI –3.34 to –1.25) and the Nordic (61.3) and Mediterranean (64.3) clusters ( $t$  test:  $p = 1.16 \times 10^{-7}$ ; CI 1.86–4.03) (figure 2A and supplementary table 6, A and B, doi.org/10.5522/04/12418157). We then assessed the relationship between pathogenic expansions and AAO via logistic regression. First, we identified a significant correlation between a decrease in AAO and presence of pathogenic expansions ( $p = 7.7 \times 10^{-4}$ ;  $R^2 = 0.008$ ; CI –8.05 [2.5%] to –2.13 [97.5%]). When we included genetic ancestry in the model, we observed a significant correlation with a decrease in AAO, with no difference

**Table 3** Stratified Fisher exact tests comparing prevalence of pathogenic expansions across behavioral variant frontotemporal dementia (bvFTD) and primary progressive aphasia (PPA) and the Nordic and Mediterranean clusters

	Expansion range		Fisher exact test
	Pathogenic	Nonpathogenic	
<b>Subtype/ancestry</b>			
<b>bvFTD</b>			
Mediterranean	8	323	$p = 1.9 \times 10^{-2a}$
Nordic	32	437	
<b>Ancestry/subtype</b>			
<b>Mediterranean</b>			
bvFTD	8	323	$p = 1$
PPA	1	168	
<b>Nordic</b>			
bvFTD	32	437	$p = 1 \times 10^{-4b}$
PPA	3	323	


*p* Values presented in the table are corrected for multiple testing statistics. Prior correction *p* values were as follows: <sup>a</sup>(uncorrected) Fisher exact test:  $p = 4.7 \times 10^{-3}$ ; odds ratio 2.95; 95% confidence interval 1.31–7.52; significant difference in the prevalence of bvFTD expansion carriers in the Nordic vs the Mediterranean cluster; <sup>b</sup>(uncorrected) Fisher exact test  $p = 2.7 \times 10^{-5}$ ; odds ratio 7.87; 95% confidence interval 2.43–40.52; significant difference in the prevalence of expansion carriers in bvFTD vs PPA within the Nordic cluster.

in using either cluster ( $p = 2.3 \times 10^{-3}$ ; CI  $-7.5 [2.5\%]$  to  $-1.63 [97.5\%]$  for pathogenic expansions;  $p = 2.3 \times 10^{-7}$ ; CI  $-3.9 [2.5\%]$  to  $-1.77 [97.5\%]$  for cluster;  $R^2 = 0.03$ ) or PC1 ( $p = 2.1 \times 10^{-3}$ ; CI  $-7.5 [2.5\%]$  to  $-1.66 [97.5\%]$  for pathogenic expansions;  $p = 6.4 \times 10^{-7}$ ; CI  $30.1 [2.5\%]$ – $68.9 [97.5\%]$  for PC1;  $R^2 = 0.028$ ) as covariate and an almost 4-fold goodness of fit increase (supplementary table 7, A–C, doi.org/10.5522/04/12418157). Of note, when comparing the 2 regression models (with/without genetic ancestry as covariate) through the log-likelihood  $R^2$  ratio test, the difference (between the 2 models) appeared not to be due to chance ( $p < 10^{-12}$ ) (supplementary table 7, B and C, doi.org/10.5522/04/12418157).

We further evaluated the relationship between expansion length (represented by 4 rc factor levels—short, intermediate, and long expansions, tested against no expansions) and AAO in the group 3 cases (figure 1). First, we independently analyzed association between AAO and (1) genetic ancestry—mean AAO 60.9 and 64.6 in the Nordic and Mediterranean cluster, respectively (*t* test:  $p = 2.1 \times 10^{-7}$ ; CI 2.32–5.09; supplementary table 8A, doi.org/10.5522/04/12418157); (2) syndrome—mean AAO 61.7 and 63.5 in the bvFTD and PPA syndromes, respectively (*t* test:  $p = 9.1 \times 10^{-3}$ ; CI  $-3.11$  to  $-0.44$ ; supplementary table 8B, doi.org/10.5522/04/12418157), and (3) expansion length—mean AAO 63.2 for both no and short expansions, 61 for intermediate expansions, and 58 for long expansions (analysis of variance  $p = 3.6 \times 10^{-2}$ ; CI  $-10.2$  to

$-0.23$  for long vs no expansions) (supplementary table 8D, figure 2B and C: doi.org/10.5522/04/12418157). We then assessed the relationship between expansion length (see above) and AAO via logistic regression. First, we identified a significant correlation between a decrease in AAO and both intermediate and long expansions ( $p = 4 \times 10^{-2}$ ; CI  $-4.36 [2.5\%]$  to  $-0.96 [97.5\%]$  for intermediate and  $p = 7 \times 10^{-3}$ ; CI  $-9.05 [2.5\%]$  to  $-1.43 [97.5\%]$  for long expansions;  $R^2 = 0.017$ ) (supplementary table 9A, doi.org/10.5522/04/12418157). When we included genetic ancestry in the model, we observed a significant correlation with a decrease in AAO, no difference in using either cluster ( $p = 4.7 \times 10^{-2}$ ; CI  $-7.65 [2.5\%]$  to  $-0.05 [97.5\%]$  for long vs no expansion;  $p = 2.38 \times 10^{-6}$ ; CI  $-4.73 [2.5\%]$  to  $-1.97 [97.5\%]$  for cluster;  $R^2 = 0.045$ ) or PC1 ( $p = 5.98 \times 10^{-2}$ ; CI  $-7.5 [2.5\%]$  to  $0.14 [97.5\%]$  for long vs no expansion;  $p = 1.2 \times 10^{-6}$ ; CI 39.8 [2.5%]–92.9 [97.5%] for PC1;  $R^2 = 0.047$ ) as covariate and an almost 3-fold goodness of fit increase (supplementary table 9, A–C, doi.org/10.5522/04/12418157). Of note, when comparing the 2 regression models (with/without genetic ancestry as covariate) through the log-likelihood  $R^2$  ratio test, the difference (between the 2 models) appeared not to be due to chance ( $p < 10^{-12}$ ) (supplementary table 9, B and C, doi.org/10.5522/04/12418157). These findings were further supported by nonlinear mixed-effects model regression using genetic ancestry as random effect covariate (for long vs no expansion; see supplementary table 10, doi.org/10.5522/04/12418157).

**Figure 2** Association between age at onset (AAO) and ancestry, syndrome, and expansion length



(A) AAO in the group 2 cases. Mean AAO behavioral variant frontotemporal dementia (bvFTD) (61.7) and primary progressive aphasia (PPA) (64) (t test:  $p = 1.86 \times 10^{-5}$ ; confidence interval [CI] -3.34 to 1.25); mean AAO Nordic (61.3) and Mediterranean (64.3) clusters (t test:  $p = 1.16 \times 10^{-7}$ ; CI 1.86-4.03). (B) AAO in the group 3 cases. Mean AAO bvFTD (61.7) and PPA (63.5) (t test:  $p = 9.1 \times 10^{-3}$ ; CI -3.11 to 0.44); mean AAO Nordic (60.9) and Mediterranean (64.6) (t test:  $p = 2.1 \times 10^{-7}$ ; CI 2.32-5.09). (C) AAO in the group 3 cases. Mean AAO for both no and short expansions (63.2), for intermediate expansions (61), and for long expansions (58) evaluated via analysis of variance test.

### C9orf72 locus risk haplotype

All of the risk alleles for the 13 markers—shortest informative stretch of the original risk haplotype<sup>15,29</sup> available to us—were seen in (1) 40/56 (71.4%) expansion carriers vs 380/1,340 (28.4%) nonexpansion carriers in the entire cohort; (2) 33/47 (70.2%) expansion carriers vs 228/826 (27.6%) nonexpansion carriers in the Nordic cluster; and (3) 7/9 (77.8%) expansion carriers vs 152/514 (29.6%) nonexpansion carriers in the Mediterranean cluster. Comparing the proportion of risk allele carriers (expansion vs nonexpansion carriers) for each single marker, 5/13 markers (rs4879515, rs868856, rs903603, rs2282241, rs2453556) were significant in the Nordic cluster, and none in the Mediterranean cluster (supplementary figure 4, doi.org/10.5522/04/12418157). Rs2477518 showed variable frequencies for the risk allele (T) across expansion vs nonexpansion carriers (and the 2 clusters), thus making this most probably a negligible marker within this stretch, as hinted previously.<sup>15,17</sup> Rs3849942, previously suggested as a surrogate marker for the risk haplotype,<sup>15</sup> was not among the SNPs available to us. We used rs868856, displaying strongest linkage disequilibrium with rs3849942 ( $D' = 0.96$ ;  $R^2 = 0.7$ ; [ldlink.nci.nih.gov/](http://ldlink.nci.nih.gov/)), as informative

proxy: the risk allele segregated differently across expansion vs nonexpansion carriers in the Nordic and Mediterranean cluster (as for rs2453556), possibly suggesting these 2 as the most conserved markers of the original risk haplotype across populations in expansion carriers (highlighted in blue in supplementary figure 4, doi.org/10.5522/04/12418157).

### Syndrome prediction

We then sought to build a model to predict syndrome (bvFTD vs PPA) and assess its accuracy. We analyzed both groups 2 and 3 cases using expansion status (presence/absence of expansion for group 2 and the 4 rc factor levels for group 3 [see Methods]), genetic ancestry (using either cluster or PC1) as binary variables, and AAO as a continuous variable in logistic regression models. We observed an accuracy of  $\sim 0.64$  (group 2; supplementary table 11, doi.org/10.5522/04/12418157) and  $\sim 0.62$  (group 3; supplementary table 12, doi.org/10.5522/04/12418157) in predicting bvFTD; there were no differences in the outcome when using either cluster or PC1 as covariates in both (LOOCV and K-fold) models.

## Discussion


This study aimed to characterize *C9orf72* expansions in relation to genetic ancestry and AAO and to assess the usefulness of these measures in discriminating the behavioral from the language variant syndrome in a large pan-European cohort of 1,396 FTL D cases.

To our knowledge, the current work is unique in that, prior to characterizing the expansions, we excluded population-substructure bias using genome-wide genotyping data to cluster the cases on the basis of their genetic makeup. After PCA, we identified 2 distinct clusters including samples with geographic ancestry corresponding to Southern Europe (Mediterranean cluster) and Central/Northern Europe (Nordic cluster). Our analyses not only showed that patients from the Nordic cluster presented significantly higher frequency of pathogenic *C9orf72* expansions compared to the Mediterranean cluster, but also that a core stretch of markers ( $n = 8$ ) of the Finnish risk haplotype<sup>29</sup> appeared to be conserved across the Nordic expansion carriers, whereas there was a similar tendency for (just) 2 of such markers in the Mediterranean expansion carriers. Several studies had shown high frequencies of long *C9orf72* expansions in Northern vs Southern European patients (North–South axis).<sup>13–15</sup> Other studies (based on the geographic location of the recruiting sites) challenged the North–South axis concept,<sup>10</sup> or the founder effect implying the existence of more than 1 risk haplotype.<sup>16–19</sup> All this taken together, our current data appear to support the North–South axis hypothesis and suggest that rearrangements (and instability)<sup>16,19</sup> at the *C9orf72* locus might have occurred, reducing the level of conservation of the original risk haplotype across the European population.

We found pathogenic expansions in ~4% of all cases and that the proportion of expansion carriers was significantly higher in bvFTD compared to PPA. The fact that we overall identified significant association between pathogenic expansions and a diagnosis of bvFTD and Central/Northern European ancestry—findings in line with previous reports<sup>8,10,13,20,30–34</sup>—suggests that *C9orf72* expansions might serve as useful genetic fingerprint to define subpopulations of FTL D (figure 3). Of note, we observed a trend of association with syndrome (bvFTD) and genetic ancestry (Central/Northern European) already supported by the intermediate repeat counts ( $9 \leq rc \leq 24$ ) category. This appears in line with previous reports suggesting that individuals with 7–24 alleles might have an increased risk to convert to carriers of pathologic repeat expansions<sup>10,22</sup> and may, altogether, be useful information in the context of diagnostics.

Despite some previous conflicting reports of direct (or inverse) correlation between *C9orf72* expansions and AAO,<sup>16,21,23</sup> we (as others<sup>22,24</sup>) found a significant inverse correlation between *C9orf72* expansion length and AAO. In addition, and interestingly, our data also indicate that Central/Northern European genetic ancestry contributes to a decreased AAO (independently from the expansions),

**Figure 3** Patient subpopulations (behavioral variant frontotemporal dementia [bvFTD] and primary progressive aphasia [PPA] syndromes) based on *C9orf72* expansion genetic signatures and ancestry



possibly implying a more complex genetic signature (or architecture), and subsequently molecular mechanisms, underpinning this feature. Clearly, disease mechanisms that involve *C9orf72* expansion length and AAO are complex, thus it is likely that additional factors might further modulate their relationship and effect on the phenotype (see also Babić Leko et al.<sup>5</sup>).

While using expansion length, genetic ancestry, and AAO in a regression model to discriminate behavioral from language variant subtypes, we found that such measures supported a prediction of bvFTD with 64% accuracy.

Our results have a number of implications. First, provided that significant variation exists in the genetic architecture of the Caucasian population,<sup>35</sup> genetic variability characterizing and differentiating Nordic vs Mediterranean subjects (such as in the case of our cohort) might influence predisposition to harboring longer repeat expansions. In other repeat expansion diseases—e.g., Huntington disease or other microsatellite diseases, including myotonic dystrophy and spinocerebellar ataxias<sup>35</sup>—the presence of specific haplogroups in Western European populations occurs with a manifold increase in prevalence of repeats compared to other ethnic groups and populations.<sup>36</sup> Second, different genetic risk architectures underpinning different (and possibly genetically more homogeneous) subpopulations of patients may exist within the FTL D population.

In a nutshell, our results imply that a significantly higher proportion of FTL D cases, with Nordic rather than Mediterranean genetic ancestry, is likely to develop bvFTD in presence of intermediate and long (pathogenic) expansions, whereas long (pathogenic) expansions are (almost) negligible in PPA, regardless of ancestry. Clearly, multiple factors including genetic heterogeneity, epigenetic changes, ethnicity, as well as environmental factors



and habits that may subsist within and across multicultural cohorts, all together, contribute to disease predisposition, onset, and progression.<sup>22,37,38</sup> These concepts, reinforced by our study, warrant further characterization of genetic, environmental, and additional clinical measures to fine-tune models able to predict disease outcome to complement diagnostic criteria, and possibly assist in the identification of informative cohorts for tailored clinical trials and the development of effective personalized therapies.

## Acknowledgment

The authors thank the patients and their families and the IFGC ([ifgcsite.wordpress.com/](http://ifgcsite.wordpress.com/)) phase III network.

## Study funding

R. Ferrari and B. Costa are supported by funding from the Alzheimer's Society (grants 284 and 447). V. Alvarez and M. Menendez-Gonzalez are supported by the Fondos Feder (grant PI 15/00878). O. Andreassen is supported by the Research Council of Norway (grant 223273), Norwegian Health Association, and the KG Jebsen Stiftelsen. L. Benussi, R. Ghidoni, G. Rossi, and F. Tagliavini are supported by the Italian Ministry of Health–Ricerca Corrente. G. Bråthen and R. Ghidoni are supported by the Italian Ministry of Health (grant RF-2016-02361492). D. Blackburn is supported by the Sheffield Biomedical Research Centre. L. Bernardi is supported by the National Institute for Health Research Cambridge Biomedical Research Centre and Biomedical Research Unit in Dementia (NIHR, grant RG64473). T.E. Cope is supported by the Association of British Neurologists; the recruitment and clinical characterization of research participants at Washington University were supported by NIH (grants R01AG044546, RF1AG053303, R01AG058501, U01AG058922, P50 AG05681, P01 AG03991, and P01 AG026276). F. Frangipane is supported by ONLUS Lamezia Terme. C. Graff is supported by grants from JPND Prefrontals Swedish Research Council (VR) 529-2014-7504, Swedish Research Council (VR) 2015-02926, Swedish Research Council (VR) 2018-02754, Swedish FTD Initiative-Schörling Foundation, Swedish Brain Foundation, Swedish Alzheimer Foundation, Stockholm County Council ALF, Karolinska Institutet Doctoral Funding, and StratNeuro, Swedish Demensfonden. M. Gallo is supported by the NIH (grant AG054519). M. Gallucci, E. Scarpini, J.C. Thompson and V.M. Van Deerlin are supported by the NIH (grant AG017586, and P30 AG10124 and U01 AG062418). J. Hardy and P.A. Lewis are supported by an MRC Programme grant (MR/N026004/1). J. Hardy is supported by the UK Dementia Research Institute, which receives its funding from DRI Ltd, funded by the UK Medical Research Council, Alzheimer's Society and Alzheimer's Research UK, Wellcome Trust (award 202903/Z/16/Z), Dolby Family Fund, National Institute for Health Research University College London Hospitals Biomedical Research Centre, BRCNIHR Biomedical Research Centre at University College London Hospitals NHS Foundation Trust, and University College London. R. Maletta is supported by the Associazione per la

Ricerca Neurogenetica. B. Miller is supported by the NIH (grants P50-AG023501 and P01-AG1972403) (BLM). C.M. Morris is supported by the Newcastle Brain Tissue Resource, the UK Medical Research Council (grant G0400074), the NIHR Newcastle Biomedical Research Centre awarded to the Newcastle upon Tyne NHS Foundation Trust and Newcastle University, and a grant from the Alzheimer's Society and Alzheimer's Research UK as part of the Brains for Dementia Research Project. B. Nacmias and S.B. Sando are supported by the Ricerca di Ateneo 2019. E. Rogaeva is supported by the Canadian Consortium on Neurodegeneration in Aging. B. Rogelj is supported by grants from the Slovenian Research Agency (grants P4-0127, J3-8201, J3-9263). J. van Rooij is supported by the Wellcome Trust (grant 103838), by the Medical Research Council (grant number SUAG004/91365), and by the National Institute for Health Research Cambridge Biomedical Research Centre and Biomedical Research Unit in Dementia (NIHR, grant RG64473). A. Rendina is supported by the Innovative Medicines Initiative 2 Joint Undertaking, which receives support from the European Union's Horizon 2020 research and innovation programme and EFPIA (grant 115975), by Acción Estratégica en Salud integrated in the Spanish National R + D + I Plan (grants PI13/02434 and PI16/01861), by ISCIII (Instituto de Salud Carlos III) Subdirección General de Evaluación, and by the Fondo Europeo de Desarrollo Regional (FEDER-Una manera de Hacer Europa). M. Boada and A. Ruiz are funded by Fundación bancaria La Caixa and Grifols SA (GR@ACE project). R. Sanchez-Valle is funded by the Spanish National Institute of Health Carlos III (ISCIII) under the aegis of the EU Joint Programme–Neurodegenerative Disease Research (JPND) (grant AC14/00013) and Fundacio Marato de TV3 (20143810) (RSV). J.Q. Trojanowski is supported by the NIH (grant AG09215). C. Van Broeckhoven and J. Van der Zee are supported by the Flemish Government initiated Impulse Program on Networks for Dementia Research (VIND) and the Methusalem Excellence Program, by the Research Foundation Flanders (FWO), and by the University of Antwerp Research Fund (Belgium). M.L. Waldö is supported by the Elly Berggren Foundation. J. Yokoyama is funded by NIA K01 AG049152.

## Disclosure

C. Cruchaga receives research support from Biogen, Eisai, Alector, and Parabon. The funders of the study had no role in the collection, analysis, or interpretation of data, in the writing of the report, or in the decision to submit the paper for publication. C. Cupidi is a member of the advisory board of ADx Healthcare and Vivid Genomics. B. Costa, C. Manzoni, M. Bernal-Quiros, D.A. Kia, M. Aguilar, I. Alvarez, V. Alvarez, O. Andreassen, M. Anfossi, S. Bagnoli, L. Benussi, L. Bernardi, G. Binetti, D. Blackburn, M. Boada, B. Borroni, L. Bowns, G. Bråthen, A.C. Bruni, H.-H. Chiang, J. Clarimon, S. Colville, M.E. Conidi, T.E. Cope, C. Cruchaga, M.E. Di Battista, J. Diehl-Schmid, M. Diez-Fairen, O. Dols-Icardo, E. Durante, D. Flisar, F. Frangipane, D. Galimberti, M. Gallo, M. Gallucci, R.

Ghidoni, C. Graff, J.H. Grafman, M. Grossman, J. Hardy, I. Hernández, G.J.T. Holloway, E.D. Huey, I. Illán-Gala, A. Karydas, B. Khoshnood, M.G. Kramberger, M. Kristiansen, P.A. Lewis, A. Lleó, G.K. Madhan, R. Maletta, A. Maver, M. Menendez-Gonzalez, G. Milan, B. Miller, M.O. Mol, P. Momeni, S. Moreno-Grau, C.M. Morris, B. Nacmias, C. Nilsson, V. Novelli, L. Öijerstedt, A. Padovani, S. Pal, Y. Panchbhaya, P. Pastor, B. Peterlin, I. Piaceri, S. Pickering-Brown, Y.A.L. Pijnenburg, A.A. Puca, I. Rainero, A. Rendina, A.M.T. Richardson, E. Rogaeva, B. Rogelj, S. Rollinson, G. Rossi, C. Rossmeier, J.B. Rowe, E. Rubino, A. Ruiz, R. Sanchez-Valle, S.B. Sando, A.F. Santillo, J. Saxon, E. Scarpini, M. Serpente, N. Smirne, S. Sorbi, E. Suh, F. Tagliavini, J.C. Thompson, J.Q. Trojanowski, V.M. Van Deerlin, J. Van der Zee, C. Van Broeckhoven, J. van Rooij, J.C. Van Swieten, A. Veronesi, E. Vitale, M.L. Waldö, C. Woodward, J. Yokoyama, V. Escott-Price, J.M. Polke, and R. Ferrari report no disclosures relevant to the manuscript. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

## Publication history

Received by *Neurology* April 15, 2020. Accepted in final form August 12, 2020.

## Appendix 1 Authors

Name	Location	Contribution
<b>Beatrice Costa, BSc</b>	University College London, Institute of Neurology, UK	C9orf72 expansions screening, data interpretation and drafting of manuscript
<b>Claudia Manzoni, PhD</b>	School of Pharmacy, University College London, UK	Project design, data analyses and interpretation, manuscript drafting
<b>Manuel Bernal-Quiros, PhD</b>	Neurogenetics Laboratory, National Hospital for Neurology and Neurosurgery, London, UK	C9orf72 expansions screening, data interpretation and drafting of manuscript
<b>Demis A. Kia, MD</b>	University College London, Institute of Neurology, UK	Data interpretation and drafting of manuscript
<b>Miquel Aguilar, MD</b>	Aptima Clinic, Terrassa, Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Ignacio Alvarez, MSc</b>	Memory Disorders Unit, Department of Neurology, University Hospital Mutua de Terrassa, Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Victoria Alvarez, PhD</b>	Hospital Universitario Central de Asturias, Oviedo, Asturias, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Ole A. Andreassen, MD, PhD</b>	NORMENT, Institute of Clinical Medicine, University of Oslo, Norway	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata

## Appendix 1 (continued)

Name	Location	Contribution
<b>Maria Anfossi, PhD</b>	Regional Neurogenetic Centre, ASPCZ, Lamezia Terme, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Silvia Bagnoli, PhD</b>	Department of Neuroscience, Psychology, Drug Research and Child Health, University of Florence, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Luisa Benussi, PhD</b>	Molecular Markers Laboratory, IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Livia Bernardi, PhD</b>	Regional Neurogenetic Centre, ASPCZ, Lamezia Terme, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Giuliano Binetti, MD</b>	Molecular Markers Laboratory, IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Daniel J. Blackburn, MD, PhD</b>	Sheffield Institute for Translational Neuroscience (SITraN), Department of Neuroscience, University of Sheffield, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Mercè Boada, MD, PhD</b>	Research Center and Memory Clinic, Fundació ACE, Institut Català de Neurociències Aplicades, Universitat Internacional de Catalunya (UIC), Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Barbara Borroni, MD</b>	Centre for Neurodegenerative Disorders, Department of Clinical and Experimental Sciences, University of Brescia, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Lucy Bownes</b>	Department of Clinical Neurosciences, Cambridge University, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Geir Bråthen, MD, PhD</b>	Department of Neurology, University Hospital of Trondheim, Norway	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Amalia C. Bruni, MD</b>	Regional Neurogenetic Centre, ASPCZ, Lamezia Terme, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata

Continued

## Appendix 1 (continued)

Name	Location	Contribution
<b>Huei-Hsin Chiang, PhD</b>	Karolinska Institutet, Dept NVS, Division of Neurogeriatrics, Bioclinicum, Solna, Sweden	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Jordi Clarimon, PhD</b>	Department of Neurology, IIB Sant Pau, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Shuna Colville, MSc</b>	Anne Rowling Regenerative Neurology Clinic, University of Edinburgh, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Maria E. Conidi, PhD</b>	Regional Neurogenetic Centre, ASPCZ, Lamezia Terme, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Tom E. Cope, MD, PhD</b>	Department of Clinical Neurosciences, Cambridge University, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Carlos Cruchaga, PhD</b>	NeuroGenomics and Informatics, Washington University, Department of Psychiatry, St. Louis, MO	Critical review of manuscript for intellectual content
<b>Chiara Cupidi, MD</b>	Regional Neurogenetic Centre, ASPCZ, Lamezia Terme, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Maria Elena Di Battista, MD, PhD</b>	Cognitive Impairment Center, Local Health Authority n.2 Marca Trevigiana, Treviso, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Janine Diehl-Schmid, MD</b>	Technical University of Munich, School of Medicine, Department of Psychiatry and Psychotherapy, Germany	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Monica Diez-Fairen, MSc</b>	Memory Disorders Unit, Department of Neurology, University Hospital Mutua de Terrassa, Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Oriol Dols-Icardo, PhD</b>	Department of Neurology, IIB Sant Pau, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata

## Appendix 1 (continued)

Name	Location	Contribution
<b>Elisabetta Durante, PhD</b>	Immunohematology and Transfusional Medicine Service, Local Health Authority n.2 Marca Trevigiana, Treviso, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Dušan Flisar, MD</b>	Department of Neurology, University Medical Center Ljubljana, Slovenia	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Francesca Frangipane, MD</b>	Regional Neurogenetic Centre, ASPCZ, Lamezia Terme, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Daniela Galimberti, PhD</b>	University of Milan, Dino Ferrari Center, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Maura Gallo, PhD</b>	Regional Neurogenetic Centre, ASPCZ, Lamezia Terme, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Maurizio Gallucci, MD</b>	Cognitive Impairment Center, Local Health Authority n.2 Marca Trevigiana, Treviso, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Roberta Ghidoni, PhD</b>	Molecular Markers Laboratory, IRCCS Istituto Centro San Giovanni di Dio Fatebenefratelli, Brescia, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Caroline Graff, MD, PhD</b>	Karolinska Institutet, Dept NVS, Division of Neurogeriatrics, Bioclinicum, Solna, Sweden	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Jordan H. Grafman, PhD</b>	Cognitive Neuroscience Lab, Think and Speak Lab, Shirley Ryan AbilityLab, Chicago, IL	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Murray Grossman, MD</b>	Department of Neurology, Perelman School of Medicine at the University of Pennsylvania, Philadelphia	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>John Hardy, PhD</b>	UCL Dementia Research Institute Wing 1.2 Cruciform Building, London, UK	Critical review of manuscript for intellectual content
<b>Isabel Hernández, MD, PhD</b>	Research Center and Memory Clinic. Fundació ACE, Institut Català de Neurociències Aplicades, Universitat Internacional de Catalunya (UIC), Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata

## Appendix 1 (continued)

Name	Location	Contribution
<b>Guy J.T. Holloway, MBBS, MRCPsych</b>	Royal Edinburgh Hospital, Edinburgh, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Edward D. Huey, MD</b>	Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia University, New York, NY	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Ignacio Illán-Gala, MD, PhD</b>	Department of Neurology, IIB Sant Pau, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Anna Karydas, MSc</b>	Department of Neurology, Memory and Aging Center, University of California, San Francisco	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Behzad Khoshnood, PhD</b>	Karolinska Institutet, Dept NVS, Division of Neurogeriatrics, Bioclinicum, Solna, Sweden	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Milica G. Kramberger, MD, PhD</b>	Department of Neurology, University Medical Center Ljubljana, Slovenia	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Mark Kristiansen, PhD</b>	UCL Genomics, UCL Great Ormond Street Institute of Child Health, London, UK	Critical review of manuscript for intellectual content
<b>Patrick A. Lewis, PhD</b>	The Royal Veterinary College, Department of Comparative Biomedical Sciences, London, UK	Critical review of manuscript for intellectual content
<b>Alberto Lleó, MD, PhD</b>	Department of Neurology, IIB Sant Pau, Hospital de la Santa Creu i Sant Pau, Universitat Autònoma de Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Gaganjit K. Madhan, MSc</b>	UCL Genomics, UCL Great Ormond Street Institute of Child Health, London, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Raffaele Maletta, MD</b>	Regional Neurogenetic Centre, ASPCZ, Lamezia Terme, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Aleš Maver, MD, PhD</b>	Clinical Institute of Medical Genetics, University Medical Center Ljubljana, Slovenia	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata

## Appendix 1 (continued)

Name	Location	Contribution
<b>Manuel Menendez-Gonzalez, MD, PhD</b>	Hospital Universitario Central de Asturias, Oviedo, Asturias, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Graziella Milan, MD</b>	Geriatric Center Frullone ASL Napoli 1 Centro, Napoli, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Bruce L. Miller, MD</b>	Department of Neurology, Memory and Aging Center, University of California, San Francisco	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Merel O. Mol, MSc</b>	Department of Neurology, Erasmus Medical Center, Rotterdam, the Netherlands	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Parastoo Momeni, PhD</b>	Rona Holdings, Silicon Valley, CA	Critical review of manuscript for intellectual content, contribution of metadata
<b>Sonia Moreno-Grau, PhD</b>	Research Center and Memory Clinic, Fundació ACE, Institut Català de Neurociències Aplicades, Universitat Internacional de Catalunya (UIC), Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Christopher M. Morris, PhD</b>	Newcastle Brain Tissue Resource, Institute of Neuroscience, Newcastle University, Edwardson Building, Campus for Ageing and Vitality, Newcastle upon Tyne, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Benedetta Nacmias, PhD</b>	Department of Neuroscience, Psychology, Drug Research and Child Health, University of Florence, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Christer Nilsson, MD</b>	Department of Neurology, Skåne University Hospital, Malmö, Sweden	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Valeria Novelli, PhD</b>	Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Linn Oijerstedt, MD</b>	Karolinska Institutet, Dept NVS, Division of Neurogeriatrics, Bioclinicum, Solna, Sweden	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata

Continued

## Appendix 1 (continued)

Name	Location	Contribution
<b>Alessandro Padovani, MD</b>	Centre for Neurodegenerative Disorders, Department of Clinical and Experimental Sciences, University of Brescia, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Suvankar Pal, MBBS, MRCP, MD</b>	Anne Rowling Regenerative Neurology Clinic, University of Edinburgh, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Yasmin Panchbhaya, MSc</b>	UCL Genomics, UCL Great Ormond Street Institute of Child Health, London, UK	Critical review of manuscript for intellectual content
<b>Pau Pastor, MD, PhD</b>	Memory Disorders Unit, Department of Neurology, University Hospital Mutua de Terrassa, Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Borut Peterlin, MD, PhD</b>	Clinical Institute of Medical Genetics, University Medical Center Ljubljana, Slovenia	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Irene Piaceri, PhD</b>	Department of Neuroscience, Psychology, Drug Research and Child Health, University of Florence, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Stuart Pickering-Brown, PhD</b>	Division of Neuroscience & Experimental Psychology, The University of Manchester, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Yolande A.L. Pijnenburg, MD, PhD</b>	Amsterdam University Medical Center, VU University Medical Center, the Netherlands	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Annibale A. Puca, MD</b>	Cardiovascular Research Unit, IRCCS Multimedica, Milan, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Innocenzo Rainero, MD, PhD</b>	Neurology I, Department of Neuroscience, University of Torino, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Antonella Rendina, PhD</b>	NeurOMICS laboratory, Institute of Biochemistry and Cell Biology (IBBC), CNR Napoli, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata

## Appendix 1 (continued)

Name	Location	Contribution
<b>Anna M.T. Richardson, FRCP</b>	Manchester Centre for Clinical Neurosciences, Salford Royal NHS Trust, Manchester, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Ekaterina Rogaeva, PhD</b>	Tanz Centre for Research in Neurodegenerative Diseases, University of Toronto, Canada	Critical review of manuscript for intellectual content
<b>Boris Rogelj, PhD</b>	Department of Biotechnology, Jožef Stefan Institute, Ljubljana, Slovenia	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Sara Rollinson, PhD</b>	Division of Neuroscience & Experimental Psychology, The University of Manchester, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Giacomina Rossi, PhD</b>	Division of Neurology V and Neuropathology; Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Carola Roßmeier, MD</b>	Technical University of Munich, School of Medicine, Department of Psychiatry and Psychotherapy, Germany	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>James B. Rowe, MD, PhD</b>	Department of Clinical Neurosciences, Cambridge University, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Elisa Rubino, MD, PhD</b>	Neurology I, Department of Neuroscience, University of Torino, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Agustín Ruiz, MD, PhD</b>	Research Center and Memory Clinic. Fundació ACE, Institut Català de Neurociències Aplicades, Universitat Internacional de Catalunya (UIC), Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Raquel Sanchez-Valle, MD, PhD</b>	Alzheimer's Disease and Other Cognitive Disorders Unit, Hospital Clínic of Barcelona, Spain	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Sigrid B. Sando, PhD</b>	Department of Neurology, University Hospital of Trondheim, Norway	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata

## Appendix 1 (continued)

Name	Location	Contribution
<b>Alexander F. Santillo, MD, PhD</b>	Clinical Memory Research Unit, Lund University, Sweden	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Jennifer Saxon, MSc</b>	Manchester Centre for Clinical Neurosciences, Salford Royal NHS Trust, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Elio Scarpini, MD</b>	University of Milan, Dino Ferrari Center, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Maria Serpente, PhD</b>	University of Milan, Dino Ferrari Center, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Nicoletta Smirne, BSc</b>	Regional Neurogenetic Centre, ASPCZ, Lamezia Terme, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Sandro Sorbi, MD</b>	Department of Neuroscience, Psychology, Drug Research and Child Health, University of Florence, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>EunRan Suh, PhD</b>	Department of Pathology and Laboratory Medicine, Perelman School of Medicine at the University of Pennsylvania, Philadelphia	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Fabrizio Tagliavini, MD</b>	Fondazione IRCCS Istituto Neurologico Carlo Besta, Milan, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Jennifer C. Thompson, PhD</b>	Manchester Centre for Clinical Neurosciences, Salford Royal NHS Trust, Manchester, UK	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>John Q. Trojanowski, MD, PhD</b>	Department of Pathology and Laboratory Medicine, Perelman School of Medicine at the University of Pennsylvania, Philadelphia	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Vivianna M. Van Deerlin, MD, PhD</b>	Department of Pathology and Laboratory Medicine, Perelman School of Medicine at the University of Pennsylvania, Philadelphia	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata

## Appendix 1 (continued)

Name	Location	Contribution
<b>Julie van der Zee, PhD</b>	Neurodegenerative Brain Diseases group, Center for Molecular Neurology, VIB, Antwerp, Belgium	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Christine Van Broeckhoven, DSc, PhD</b>	Neurodegenerative Brain Diseases group, Center for Molecular Neurology, VIB, Antwerp, Belgium	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Jeroen G.J. van Rooij</b>	Department of Neurology, Erasmus Medical Center, Rotterdam, the Netherlands	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>John C. van Swieten, MD, PhD</b>	Department of Neurology, Erasmus Medical Center, Rotterdam, the Netherlands	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Arianna Veronesi, MD, PhD</b>	Cognitive Impairment Center, Local Health Authority n.2 Marca Trevigiana, Treviso, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Emilia Vitale, PhD</b>	NeuroMICS laboratory, Institute of Biochemistry and Cell Biology (IBBC), CNR Napoli, Italy	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Maria L. Waldö, MD, PhD</b>	Division of Clinical Sciences Helsingborg, Department of Clinical Sciences Lund, Lund University, Sweden	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Cathy Woodward, MSc</b>	Neurogenetics Laboratory, National Hospital for Neurology and Neurosurgery, London, UK	C9orf72 expansions screening, data interpretation and drafting of manuscript
<b>Jennifer S. Yokoyama, PhD</b>	Department of Neurology, Memory and Aging Center, University of California, San Francisco	Critical review of manuscript for intellectual content, contribution of samples and demographics metadata
<b>Valentina Escott-Price, PhD</b>	Medical Research Council Centre for Neuropsychiatric Genetics and Genomics, Division of Psychological Medicine and Clinical Neurosciences, Cardiff University, UK and Dementia Research Institute, Cardiff University, UK	Project design, data analyses and interpretation, manuscript drafting
<b>James M. Polke, PhD</b>	Neurogenetics Laboratory, National Hospital for Neurology and Neurosurgery, London, UK	C9orf72 expansions screening, data interpretation and drafting of manuscript
<b>Raffaele Ferrari, PhD</b>	Department of Neurodegenerative Disease, University College London, Institute of Neurology, UK	Project design, data analyses and interpretation, manuscript drafting

## Appendix 2 Coinvestigators

Coinvestigators are listed at [links.lww.com/WNL/B240](https://links.lww.com/WNL/B240)

### References

1. Ratnavalli E, Brayne C, Dawson K, Hodges JR. The prevalence of frontotemporal dementia. *Neurology* 2002;58:1615–1621.
2. Neary D, Snowden JS, Gustafson L, et al. Frontotemporal lobar degeneration: a consensus on clinical diagnostic criteria. *Neurology* 1998;51:1546–1554.
3. Gorno-Tempini ML, Hillis AE, Weintraub S, et al. Classification of primary progressive aphasia and its variants. *Neurology* 2011;76:1006–1014.
4. Strong MJ, Abrahams S, Goldstein LH, et al. Amyotrophic lateral sclerosis–frontotemporal spectrum disorder (ALS-FTSD): revised diagnostic criteria. *Amyotroph Lateral Scler Frontotemporal Degener* 2017;18:153–174.
5. Babić Leko M, Zupunski V, Kirinčić J, et al. Molecular mechanisms of neurodegeneration related to C9orf72 hexanucleotide repeat expansion. *Behav Neurol* 2019;2019:2909168.
6. DeJesus-Hernandez M, Mackenzie IR, Boeve BF, et al. Expanded GGGGCC hexanucleotide repeat in noncoding region of C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* 2011;72:245–256.
7. Ferrari R, Manzoni C, Hardy J. Genetics and molecular mechanisms of frontotemporal lobar degeneration: an update and future avenues. *Neurobiol Aging* 2019;78:98–110.
8. Pottier C, Ravenscroft TA, Sanchez-Contreras M, Rademakers R. Genetics of FTL: overview and what else we can expect from genetic studies. *J Neurochem* 2016;138(suppl 1):32–53.
9. Seelaar H, Rohrer JD, Pijnenburg YA, Fox NC, van Swieten JC. Clinical, genetic and pathological heterogeneity of frontotemporal dementia: a review. *J Neurol Neurosurg Psychiatry* 2011;82:476–486.
10. van der Zee J, Gijssels I, Dillen L, et al. A Pan-European study of the C9orf72 repeat associated with FTL: geographic prevalence, genomic instability, and intermediate repeats. *Hum Mutat* 2013;34:363–373.
11. LT T. The genetics of monogenic frontotemporal dementia. *Dement Neuropsychol* 2015;219–229.
12. Turner MR, Al-Chalabi A, Chio A, et al. Genetic screening in sporadic ALS and FTD. *J Neurol Neurosurg Psychiatry* 2017;88:1042–1044.
13. Majounie E, Renton AE, Mok K, et al. Frequency of the C9orf72 hexanucleotide repeat expansion in patients with amyotrophic lateral sclerosis and frontotemporal dementia: a cross-sectional study. *Lancet Neurol* 2012;11:323–330.
14. Ramos EM, Koros C, Dokuru DR, et al. Frontotemporal dementia spectrum: first genetic screen in a Greek cohort. *Neurobiol Aging* 2019;75:224.e221–224.e228.
15. Mok K, Traynor BJ, Schymick J, et al. Chromosome 9 ALS and FTD locus is probably derived from a single founder. *Neurobiol Aging* 2012;33:209.e203–209.e208.
16. Beck J, Poulter M, Hensman D, et al. Large C9orf72 hexanucleotide repeat expansions are seen in multiple neurodegenerative syndromes and are more frequent than expected in the UK population. *Am J Hum Genet* 2013;92:345–353.
17. Chiang HH, Forsell C, Lindstrom AK, et al. No common founder for C9orf72 expansion mutation in Sweden. *J Hum Genet* 2017;62:321–324.
18. Fratta P, Polke JM, Newcombe J, et al. Screening a UK amyotrophic lateral sclerosis cohort provides evidence of multiple origins of the C9orf72 expansion. *Neurobiol Aging* 2015;36:546 e541–547.
19. Xi Z, van Blitterswijk M, Zhang M, et al. Jump from pre-mutation to pathologic expansion in C9orf72. *Am J Hum Genet* 2015;96:962–970.
20. Van Mossevelde S, Engelborghs S, van der Zee J, Van Broeckhoven C. Genotype-phenotype links in frontotemporal lobar degeneration. *Nat Rev Neurol* 2018;14:363–378.
21. Fournier C, Barbier M, Camuzat A, et al. Relations between C9orf72 expansion size in blood, age at onset, age at collection and transmission across generations in patients and presymptomatic carriers. *Neurobiol Aging* 2019;74:234.e231–234.e238.
22. Gijssels I, Van Mossevelde S, van der Zee J, et al. The C9orf72 repeat size correlates with onset age of disease, DNA methylation and transcriptional downregulation of the promoter. *Mol Psychiatry* 2016;21:1112–1124.
23. van Blitterswijk M, DeJesus-Hernandez M, Niemantsverdriet E, et al. Association between repeat sizes and clinical and pathological characteristics in carriers of C9ORF72 repeat expansions (Xpansize-72): a cross-sectional cohort study. *Lancet Neurol* 2013;12:978–988.
24. Van Mossevelde S, van der Zee J, Gijssels I, et al. Clinical evidence of disease anticipation in families segregating a C9orf72 repeat expansion. *JAMA Neurol* 2017;74:445–452.
25. Rascovsky K, Hodges JR, Knopman D, et al. Sensitivity of revised diagnostic criteria for the behavioural variant of frontotemporal dementia. *Brain* 2011;134:2456–2477.
26. Blauwendraat C, Faghri F, Pihlstrom L, et al. NeuroChip, an updated version of the NeuroX genotyping platform to rapidly screen for variants associated with neurological diseases. *Neurobiol Aging* 2017;57:247 e249–247 e213.
27. Ferrari R, Mok K, Moreno JH, et al. Screening for C9ORF72 repeat expansion in FTL. *Neurobiol Aging* 2012;33:1850.e1–e11.
28. Renton AE, Majounie E, Waite A, et al. A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-linked ALS-FTD. *Neuron* 2011;72:257–268.
29. Laaksovirta H, Peuralinna T, Schymick JC, et al. Chromosome 9p21 in amyotrophic lateral sclerosis in Finland: a genome-wide association study. *Lancet Neurol* 2010;9:978–985.
30. Benussi L, Rossi G, Glionna M, et al. C9ORF72 hexanucleotide repeat number in frontotemporal lobar degeneration: a genotype-phenotype correlation study. *J Alzheimers Dis* 2015;45:319.
31. Devenney E, Bartley L, Hoon C, et al. Progression in behavioral variant frontotemporal dementia: a longitudinal study. *JAMA Neurol* 2015;72:1501–1509.
32. Galimberti D, Fenoglio C, Serpente M, et al. Autosomal dominant frontotemporal lobar degeneration due to the C9ORF72 hexanucleotide repeat expansion: late-onset psychotic clinical presentation. *Biol Psychiatry* 2013;74:384–391.
33. Ramos EM, Dokuru DR, Van Berlo V, et al. Genetic screen in a large series of patients with primary progressive aphasia. *Alzheimers Dement* 2019;15:553–560.
34. Simon-Sanchez J, Dopper EG, Cohn-Hokke PE, et al. The clinical and pathological phenotype of C9ORF72 hexanucleotide repeat expansions. *Brain* 2012;135:723–735.
35. Ralph P, Coop G. The geography of recent genetic ancestry across Europe. *PLoS Biol* 2013;11:e1001555.
36. Warby SC, Montpetit A, Hayden AR, et al. CAG expansion in the Huntington disease gene is associated with a specific and targetable predisposing haplogroup. *Am J Hum Genet* 2009;84:351–366.
37. Huang T, Shu Y, Cai YD. Genetic differences among ethnic groups. *BMC Genomics* 2015;16:1093.
38. Zhang M, Ferrari R, Tartaglia MC, et al. A C6orf10/LOC101929163 locus is associated with age of onset in C9orf72 carriers. *Brain* 2018;141:2895–2907.

# Neurology®

## ***C9orf72*, age at onset, and ancestry help discriminate behavioral from language variants in FTL D cohorts**

Beatrice Costa, Claudia Manzoni, Manuel Bernal-Quiros, et al.  
*Neurology* 2020;95:e3288-e3302 Published Online before print September 17, 2020  
DOI 10.1212/WNL.0000000000010914

**This information is current as of September 17, 2020**

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://n.neurology.org/content/95/24/e3288.full">http://n.neurology.org/content/95/24/e3288.full</a>
<b>References</b>	This article cites 37 articles, 5 of which you can access for free at: <a href="http://n.neurology.org/content/95/24/e3288.full#ref-list-1">http://n.neurology.org/content/95/24/e3288.full#ref-list-1</a>
<b>Subspecialty Collections</b>	This article, along with others on similar topics, appears in the following collection(s): <b>All Genetics</b> <a href="http://n.neurology.org/cgi/collection/all_genetics">http://n.neurology.org/cgi/collection/all_genetics</a> <b>Frontotemporal dementia</b> <a href="http://n.neurology.org/cgi/collection/frontotemporal_dementia">http://n.neurology.org/cgi/collection/frontotemporal_dementia</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://www.neurology.org/about/about_the_journal#permissions">http://www.neurology.org/about/about_the_journal#permissions</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://n.neurology.org/subscribers/advertise">http://n.neurology.org/subscribers/advertise</a>

*Neurology*® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

