Serum albumin and beta-amyloid deposition in the human brain

Jee Wook Kim, MD, PhD, Min Soo Byun, MD, PhD, Jun Ho Lee, MD, Dahyun Yi, PhD, So Yeon Jeon, MD, Bo Kyung Sohn, MD, Jun-Young Lee, MD, PhD, Seong A Shin, MS, Yu Kyeong Kim, MD, PhD, Koung Mi Kang, MD, Chul-Ho Sohn, MD, PhD, and Dong Young Lee, MD, PhD, for the KBASE Research Group

Correspondence

Dr. Lee
selfpsy@snu.ac.kr

Neurology® 2020;95:e815-e826. doi:10.1212/WNL.00000000000010005

Abstract

Objectives

To investigate the relationships of serum albumin with in vivo Alzheimer disease (AD) pathologies, including cerebral β -amyloid (A β) protein deposition, neurodegeneration of AD-signature regions, and cerebral white matter hyperintensities (WMH), in the human brain.

Methods

A total of 396 older adults without dementia underwent comprehensive clinical assessments, measurement of serum albumin level, and multimodal brain imaging, including [^{11}C] Pittsburgh compound B-PET, ^{18}F -fluorodeoxyglucose-PET, and MRI. Serum albumin was categorized as follows: <4.4 g/dL (low albumin), 4.4 to 4.5 g/dL (middle albumin), and >4.5 g/dL (high albumin; used as a reference category). A β positivity, AD-signature region cerebral glucose metabolism (AD-CM), AD-signature region cortical thickness (AD-CT), and WMH volume were used as outcome measures.

Results

Serum albumin level (as a continuous variable) was inversely associated with A β deposition and A β positivity. The low albumin group showed a significantly higher A β positivity rate compared to the high albumin group (odds ratio 3.40, 95% confidence interval 1.67–6.92, $p = 0.001$), while the middle albumin group showed no difference (odds ratio 1.74, 95% confidence interval 0.80–3.77, $p = 0.162$). Neither serum albumin level (as a continuous variable) nor albumin categories were related to AD-CM, AD-CT, or WMH volume.

Conclusions

Low serum albumin may increase the risk of AD dementia by elevating amyloid accumulation. In terms of AD prevention, more attention needs to be paid to avoid a low serum albumin level, even within the clinical normal range, by clinicians.

From the Department of Neuropsychiatry (J.W.K.), Hallym University Dongtan Sacred Heart Hospital, Hwaseong-si, Gyeonggi-do; Department of Psychiatry (J.W.K.), Hallym University College of Medicine, Chuncheon, Gangwon-do; Institute of Human Behavioral Medicine (M.S.B., D.Y., D.Y.L.), Medical Research Center Seoul National University; Departments of Neuropsychiatry (J.H.L., D.Y.L.) and Radiology (K.M.K., C.-H.S.), Seoul National University Hospital; Department of Psychiatry (S.Y.J.), Chungnam National University Hospital, Daejeon; Sanggye Paik Hospital (B.K.S.), Department of Psychiatry, Inje University College of Medicine; Departments of Neuropsychiatry (J.-Y.L.) and Nuclear Medicine (S.A.S., Y.K.K.), SMG-SNU Boramae Medical Center; and Department of Psychiatry (J.-Y.L., D.Y.L.), Seoul National University College of Medicine, Republic of Korea.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Coinvestigators are listed at links.lww.com/WNL/B134

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Glossary

A β = β -amyloid; **AD** = Alzheimer disease; **AD-CM** = AD-signature cerebral glucose metabolism; **AD-CT** = AD-signature cortical thickness; **AMBAR** = Alzheimer's Management by Albumin Replacement; **CN** = cognitively normal; **FDG** = ¹⁸F-fluorodeoxyglucose; **FLAIR** = fluid-attenuated inversion recovery; **KBASE** = Korean Brain Aging Study for Early Diagnosis and Prediction of Alzheimer's Disease; **MCI** = mild cognitive impairment; **MCL** = minimum cost of living; **MR** = magnetic resonance; **PiB** = Pittsburgh compound B; **ROI** = region of interest; **SUVr** = standardized uptake value ratio; **WMH** = white matter hyperintensities.

Alzheimer disease (AD) is characterized by extracellular deposition of β -amyloid protein (A β) and intraneuronal neurofibrillary tangles in the brain.¹ Because A β deposition is the earliest pathology of AD and begins $\approx$ 10 to 15 years before the onset of the cognitive symptoms,^{2–4} identification of factors affecting cerebral A β deposition may facilitate the development of strategies for preventing AD dementia.

Serum albumin, which is the most abundant protein in human plasma,⁵ is regarded as one of the most potent A β sequestering systems in that it binds 90% to 95% of the A β in blood plasma.^{6,7} The dynamic equilibrium of A β between brain and blood plasma may be shifted toward the bloodstream by peripheral serum albumin that binds the A β .⁸ Therefore, a reduction in A β binding to serum albumin in the blood may lead to a decrease in the capacity for A β excretion from the brain to the blood, resulting in A β deposition in the brain.⁹

Preclinical studies suggested that serum albumin may inhibit A β fibril formation by binding A β monomers⁷ or oligomers.¹⁰ Several human studies have indicated that a low serum albumin level is associated with cognitive impairment^{11–13} and AD dementia.^{14,15} However, little information is available on whether serum albumin is related to A β deposition in the living human brain. Therefore, the present study was performed to examine the relationships of serum albumin and in vivo cerebral A β deposition in older adults without dementia. We additionally investigated the associations of serum albumin with AD-signature neurodegeneration and white matter hyperintensities (WMH) as a measure of cerebrovascular injury. While some human studies have reported an inverse association of serum albumin and cerebrovascular disease,^{16,17} few studies have investigated the relationships of serum albumin with neurodegenerative changes in the brain.

Methods

Participants

The present study was performed as part of the Korean Brain Aging Study for Early Diagnosis and Prediction of Alzheimer's Disease (KBASE), which is an ongoing prospective cohort study.¹⁸ As of November 2016, a total of 396 older adults without dementia (284 cognitively normal [CN] individuals and 112 individuals with mild cognitive

impairment [MCI]) between 55 and 90 years of age were enrolled in the study. The CN group consisted of participants with a Clinical Dementia Rating¹⁹ score of 0 and no diagnosis of MCI or dementia. All individuals with MCI met the current consensus criteria for amnesic MCI, which are as follows: (1) memory complaints confirmed by an informant, (2) objective memory impairments, (3) preserved global cognitive function, (4) independence in functional activities, and (5) no dementia. With regard to criterion 2, the age-, education-, and sex-adjusted z scores for at least 1 of 4 episodic memory tests were <-1.0 . The 4 memory tests were the Word List Memory, Word List Recall, Word List Recognition, and Constructional Recall tests, which are included in the Korean version of the Consortium to Establish a Registry for Alzheimer's Disease neuropsychological battery.²⁰ All individuals with MCI had a Clinical Dementia Rating score of 0.5. The exclusion criteria were as follows: (1) presence of a major psychiatric illness; (2) significant neurologic (e.g., cerebrovascular disease) or medical conditions that could affect mental function; (3) contraindications for MRI (e.g., pacemaker or claustrophobia); (4) illiteracy; (5) presence of significant visual/hearing difficulties or severe communication or behavioral problems that would make clinical examinations or brain scans difficult; (6) taking an investigational drug; and (7) pregnant or breastfeeding. The presence of any item included in the exclusion criteria was determined by research clinicians referring to the results of laboratory examinations and MRI and the clinical data collected by trained nurses during systematic interviews of participants and their reliable informants during the screening period. More detailed information on the recruitment of the KBASE cohort is presented in a previous report from our research group.¹⁸

Standard protocol approvals, registrations, and patient consents

This study protocol was approved by the Institutional Review boards of the Seoul National University Hospital (C-1401-027-547) and SMG-SNU Boramae Medical Center (26-2015-60), Seoul, South Korea, and the study was conducted in accordance with the recommendations of the current version of the Declaration of Helsinki. The subjects or their legal representatives gave written informed consent.

Clinical assessments

All participants underwent comprehensive clinical and neuropsychological assessments administered by trained

Table 1 Participant characteristics^a

Characteristic	Overall (n = 396)	Aβ/AD-CM (n = 388)	AD-CT (n = 380)	WMH (n = 348)
Age, y	70.48 (8.04)	70.41 (8.03)	70.39 (8.07)	70.28 (8.04)
Female, n (%)	222 (56.06)	218 (56.19)	215 (56.58)	191 (54.89)
Education, y	11.01 (4.80)	11.14 (4.83)	11.16 (4.81)	11.23 (4.76)
APOE ε4 positivity, n (%)	78 (19.70)	76 (19.59)	73 (19.21)	66 (18.97)
CN, n (%)	284 (71.72)	281 (72.42)	275 (72.37)	256 (73.56)
Serum albumin level				
Overall serum albumin, g/dL	4.47 (0.24)	4.46 (0.24)	4.46 (0.24)	4.47 (0.24)
Categorized serum albumin, n (%)				
Low: 3.6–4.3 g/dL	117 (29.55)	117 (30.15)	115 (30.26)	99 (28.45)
Middle: 4.4–4.5 g/dL	132 (33.33)	127 (32.73)	125 (29.89)	116 (33.33)
High: 4.6–5.3 g/dL	147 (37.12)	144 (37.11)	140 (36.84)	133 (38.22)
Alcohol use, n (%)				
Never	211 (53.28)	207 (53.49)	204 (53.83)	185 (53.31)
Former	52 (13.13)	50 (12.92)	49 (12.93)	45 (12.97)
Drinker	132 (33.33)	130 (33.59)	126 (33.25)	117 (33.72)
Medication use within 4 wk, n (%)				
No	84 (21.21)	84 (21.71)	84 (22.16)	73 (21.04)
Yes	311 (78.54)	303 (78.29)	295 (77.84)	274 (78.96)
Decrease in food intake over the past 3 mo, n (%)				
No	327 (83.00)	369 (95.60)	362 (95.51)	330 (95.38)
Yes	67 (17.00)	17 (4.40)	17 (4.49)	16 (4.62)
Medical condition, n (%)				
Liver disease	69 (17.42)	69 (17.78)	68 (17.89)	62 (17.82)
Kidney disease	16 (4.04)	16 (4.12)	15 (3.95)	16 (4.60)
Diabetes mellitus	66 (16.67)	66 (17.01)	62 (16.32)	59 (16.95)
Hyperlipidemia	136 (34.34)	135 (34.79)	132 (34.74)	126 (36.21)
Cerebral Aβ deposition				
Global Aβ retention, SUVR		1.21 (0.26)		
Aβ positivity, n (%)		68 (17.17)		
Neurodegeneration				
AD-CM, SUVR		1.40 (0.12)		
AD-CT, mm			2.82 (0.20)	
WMH volume, cm ³				5.95 (5.43)

Abbreviations: Aβ = β-amyloid protein; AD-CM = Alzheimer disease–signature cerebral glucose metabolism; AD-CT = Alzheimer disease–signature cortical thickness; CN = cognitively normal; SUVR = standardized uptake value ratio; WMH = white matter hyperintensities.

^a Unless otherwise indicated, data are expressed as mean (SD).

psychiatrists and neuropsychologists according to the KBASE assessment protocol,¹⁸ which incorporates the Korean version of the Consortium to Establish a Registry

for Alzheimer's Disease.^{21,22} Medication use within 4 weeks and alcohol intake status (never/former/drinker) were evaluated by nurse interviews and by a review of medical

Table 2 Relationships between serum albumin and neuroimaging biomarkers

Serum albumin	OR	95% CI	p Value
Dependent variable: Aβ positivity (n = 388)			
Model 1 ^a	0.13	0.04–0.39	<0.001
Model 2 ^b	0.16	0.05–0.52	0.002
Model 3 ^c	0.17	0.05–0.58	0.004
	B	95% CI	p Value
Dependent variable: global Aβ retention (n = 388)			
Model 1 ^a	–0.14	–0.21 to –0.06	<0.001
Model 2 ^b	–0.11	–0.18 to –0.04	0.002
Model 3 ^c	–0.09	–0.16 to –0.03	0.008
Dependent variable: AD-CM (n = 388)			
Model 1 ^a	0.01	–0.05 to 0.06	0.816
Model 2 ^b	–<0.01	–0.06 to 0.05	0.885
Model 3 ^c	<0.01	–0.05 to 0.06	0.883
Dependent variable: AD-CT (n = 380)			
Model 1 ^a	0.07	–0.01 to 0.16	0.092
Model 2 ^b	0.03	–0.05 to 0.10	0.457
Model 3 ^c	0.02	–0.06 to 0.09	0.641
Dependent variable: WMH volume (n = 348)			
Model 1 ^a	0.38	–2.03 to 2.79	0.758
Model 2 ^b	0.79	–1.58 to 3.16	0.513
Model 3 ^c	0.68	–1.74 to 3.10	0.581

Abbreviations: A β = β -amyloid protein; AD-CM = Alzheimer disease–signature cerebral glucose metabolism; AD-CT = Alzheimer disease–signature cortical thickness; CI = confidence interval; OR = odds ratio; WMH = white matter hyperintensities.

Global A β retention was used after natural log-transformation to achieve a normal distribution.

^a Not adjusted.

^b Adjusted for age, sex, and APOE ϵ 4.

^c Adjusted for age, sex, APOE ϵ 4, education, clinical diagnosis, medication use, alcohol use, annual income, diabetes mellitus, hyperlipidemia, liver disease, and kidney disease.

records. Participants were divided into 3 groups according to annual income: below the minimum cost of living (MCL), above the MCL but below twice the MCL, and twice the MCL or more (law.go.kr). The MCL was determined according to the administrative rules published by the Ministry of Health and Welfare, Republic of Korea, in November 2012. The MCL was 572,168 Korea Won for single-person households and added 286,840 Korea Won for each additional housemate. The presence of comorbid diabetes mellitus, dyslipidemia, liver disease, and kidney disease was assessed from data collected by trained nurses during systematic interviews of participants and their reliable informants. Information about nutritional state, including the change in food intake over the past 3 months due to loss of appetite, digestive problems, or chewing or swallowing difficulties, was systematically obtained by using the Mini Nutritional Assessment tool.²³ To acquire accurate

information, reliable informants were interviewed and medical records were reviewed.

Laboratory tests of blood samples

Blood samples were obtained by venipuncture after an overnight fast. The concentration of albumin was determined by bromocresol green dye binding assay (ADVIA 1800; Siemens, Washington, DC). The normal range for albumin level is 3.5 to 5.5 g/dL. Participants were categorized on the basis of serum albumin level using tertiles referring to previous studies^{24,25} as follows: low albumin, <4.4 g/dL; middle albumin, 4.4 to 4.5 g/dL; and high albumin, >4.5 g/dL (used as a reference category). In addition, genomic DNA was extracted from whole blood, and APOE genotyping was performed as described previously.²⁶ APOE ϵ 4 positivity was defined as the presence of at least 1 ϵ 4 allele.

Figure 1 Partial correlation plots for the relationships of serum albumin with neuroimaging biomarkers

Multiple linear regression analysis was performed for (A) global A β retention, (B) AD-CM, (C) AD-CT, and (D) WMH after adjustment for age, sex, and *APOE* $\epsilon 4$. Global A β retention was used after natural log transformation to achieve a normal distribution. A β = beta-amyloid protein; AD-CM = Alzheimer disease–signature cerebral glucose metabolism; AD-CT = Alzheimer disease–signature cortical thickness; WMH = white matter hyperintensities.

Measurement of cerebral A β deposition

All participants underwent simultaneous 3D [^{11}C] Pittsburgh compound B (PiB)-PET and 3D T1-weighted MRI scan with a 3.0T Biograph mMR (PET-MR) scanner (Siemens) according to the manufacturer's guidelines. Details of the PiB-PET imaging acquisition and preprocessing were described previously.²⁷ An automatic anatomic labeling algorithm and a region-combining method²⁸ were applied to determine regions of interest (ROIs) to characterize the PiB retention levels in the frontal, lateral parietal, posterior cingulate-precuneus, and lateral temporal regions. The standardized uptake value ratio (SUVR) value for each ROI was calculated by dividing the mean value for all voxels within each ROI by the mean cerebellar uptake value in the same image. A global cortical ROI consisting of the 4 ROIs was also defined, and a global A β retention value was generated by dividing the mean value for all voxels of the global cortical ROI by the mean cerebellar uptake value in the same image.^{28,29} Given the characteristic skewed distribution of cerebral A β deposition values, global A β retention data were used after natural log-transformation and removal of extreme outliers to achieve a normal distribution with reference to previous reports.^{30,31} Amyloid positivity was also used as an outcome variable. Each participant was classified as A β -positive (A β +) if the SUVR value was >1.4 in at least 1 of the 4 ROIs

or as A β -negative (A β -) if the SUVR values were ≤ 1.4 for all 4 ROIs.^{28,32}

Measurement of AD-signature neurodegeneration

All participants underwent ^{18}F -fluorodeoxyglucose (FDG)-PET imaging with the abovementioned PET-MR machine. Details of the FDG-PET image acquisition and preprocessing were described previously.²⁷ AD-signature FDG ROIs such as the angular gyri, posterior cingulate cortex, and inferior temporal gyri, which are sensitive to the changes associated with AD,³² were determined. AD-signature cerebral glucose metabolism (AD-CM) was defined as the voxel-weighted mean SUVR extracted from the AD-signature FDG ROIs. Details of MRI acquisition and preprocessing were described previously.²⁷ AD-signature cortical thickness (AD-CT) was defined as the mean cortical thickness values obtained from AD-signature regions, including the entorhinal, inferior temporal, middle temporal, and fusiform gyrus, as described previously.³²

Measurement of WMH

All participants underwent MRI scans with fluid-attenuated inversion recovery (FLAIR) using the abovementioned 3.0T PET-MR scanner. We followed the validated automatic procedure reported previously.³³ Briefly, the procedure

Table 3 Relationships between stratified serum albumin and neuroimaging biomarkers

	Stratified serum albumin				
	Low (<4.4 g/dL)		Middle (4.4–4.5 g/dL)		High (>4.5 g/dL)
	OR (95% CI)	p Value	OR (95% CI)	p Value	
Aβ positivity (n = 388)					
Model 1 ^a	3.96 (2.01–7.81)	<0.001	1.55 (0.74–3.26)	0.250	Reference
Model 2 ^b	3.40 (1.67– 6.92)	0.001	1.74 (0.80–3.77)	0.162	Reference
Model 3 ^c	3.08 (1.47–6.46)	0.003	1.77 (0.79–3.99)	0.168	Reference
	B (95% CI)	p Value	B (95% CI)	p Value	
Global Aβ retention (n = 388)					
Model 1 ^a	0.09 (0.05 to 0.13)	<0.001	<0.01 (–0.03 to 0.05)	0.719	Reference
Model 2 ^b	0.07 (0.03 to 0.11)	<0.001	0.01 (–0.03 to 0.05)	0.623	Reference
Model 3 ^c	0.06 (0.02 to 0.10)	0.006	<0.01 (–0.03 to 0.05)	0.751	Reference
AD-CM (n = 388)					
Model 1 ^a	–0.01 (–0.05 to 0.02)	0.372	0.01 (–0.03 to 0.04)	0.735	Reference
Model 2 ^b	–0.01 (–0.04 to 0.02)	0.659	0.01 (–0.02 to 0.04)	0.669	Reference
Model 3 ^c	–0.01 (–0.04 to 0.02)	0.592	0.01 (–0.02 to 0.04)	0.712	Reference
AD-CT (n = 380)					
Model 1 ^a	–0.07 (–0.12 to –0.02)	0.009	0.01 (–0.04 to 0.06)	0.680	Reference
Model 2 ^b	–0.04 (–0.08 to 0.01)	0.124	0.01 (–0.03 to 0.05)	0.625	Reference
Model 3 ^c	–0.03 (–0.07 to 0.02)	0.246	0.01 (–0.03 to 0.05)	0.616	Reference
WMH (n = 340)					
Model 1 ^a	0.29 (–1.13 to 1.71)	0.687	–0.16 (–1.52 to 1.21)	0.821	Reference
Model 2 ^b	–0.04 (–1.45 to 1.37)	0.959	–0.35 (–1.69 to 0.99)	0.609	Reference
Model 3 ^c	0.04 (–1.40 to 1.48)	0.956	–0.26 (–1.62 to 1.10)	0.707	Reference

Abbreviations: Aβ = β-amyloid protein; AD-CM = Alzheimer disease–signature cerebral glucose metabolism; AD-CT = Alzheimer disease–signature cortical thickness; CI = confidence interval; OR = odds ratio; WMH = white matter hyperintensities.

Global Aβ retention was used after natural log-transformation to achieve a normal distribution.

^a Not adjusted.

^b Adjusted for age, sex, and APOE ε4.

^c Adjusted for age, sex, APOE ε4, education, clinical diagnosis, medication use, alcohol use, annual income, diabetes mellitus, hyperlipidemia, liver disease, and kidney disease.

consisted of 11 steps: spatial coregistration of T1 and FLAIR images; fusion of T1 and FLAIR images; segmentation of T1; attainment of transformation parameters; deformation and obtainment of the white matter mask; obtainment of FLAIR within the white matter mask; intensity normalization of the masked FLAIR; nomination of candidate WMH with a designated threshold; creation of a junction map; and elimination of the junction. There were 2 modifications in the current processing procedure compared to the original study: (1) an optimal threshold of 70 was applied because it was more suitable for our data compared to the threshold of 65 used in the original study; and, (2) given that individuals with acute cerebral infarcts were not enrolled in our sample, we did not use diffusion-weighted imaging in the current

automated procedure. With the use of the final WMH candidate image, the WMH volume was extracted in the native space in each subject.

Statistical analysis

Multiple linear regression analysis with serum albumin (continuous variable) as the independent variable and AD-neuroimaging parameters (Aβ deposition, AD-CM, AD-CT, and WMH) as the dependent variables was performed. Multiple logistic regression analysis with serum albumin (continuous variables or categorical variables, i.e., low albumin, middle albumin, and high albumin) as the independent variable and Aβ positivity as the dependent variable was also conducted. Three models were tested for stepwise control of

Figure 2 Relationship of stratified serum albumin with A β deposition

Multiple regression analyses were performed for (A) β -amyloid (A β) positivity and (B) global A β retention after adjustment for age, sex, and *APOE* ϵ 4. Error bars represent standard deviation.

potential confounders that could affect the relationships between albumin and AD biomarkers. The first model did not include any covariates. The second model included age (<70 vs $\geq$ 70 years), sex (female vs male), and *APOE* ϵ 4 positivity (+/–) as covariates. The third model included all potential covariates: age, sex, *APOE* ϵ 4, education, clinical diagnosis (CN vs MCI), medication use within 4 weeks (+/–), alcohol use (no/former/drinker), annual income, diabetes mellitus (+/–), hyperlipidemia (+/–), liver disease (+/–), and kidney disease (+/–), which have been considered possible confounders in previous studies.^{25,34–38} In the analysis, high albumin was used as a reference (i.e., high albumin vs middle albumin, or high albumin vs low albumin). As sensitivity analyses, the same analyses were performed for the subjects with no decrease in food intake over the past 3 months due to loss of appetite, digestive problems, or chewing or swallowing difficulties to eliminate any influence of physical or mental conditions that can potentially relate to both serum albumin level and brain status. Statistical analyses were performed with IBM SPSS Statistics 24 (IBM Corp, Armonk, NY). In all analyses, values of $p < 0.05$ were taken to indicate statistical significance.

Data availability

The data for this analysis are owned by the KBASE research group. Requests for data access can be submitted to the administrative coordinator of the group by e-mail (kbasecohort@gmail.com).

Results

Participant characteristics

The demographic and clinical characteristics of the participants are presented in table 1. All participants had serum albumin levels within normal range. Among them, 117 were

categorized as having low albumin, 132 as having middle albumin, and 147 as having high albumin group.

Association of serum albumin with cerebral amyloid deposition

Serum albumin level (as a continuous variable) was significantly associated with global A β retention (table 2 and figure 1A) and A β positivity after controlling for the covariates (table 2). Furthermore, the low albumin category was significantly associated with higher A β positivity compared to the high albumin category after controlling for the covariates, while the middle albumin category showed no relation to A β positivity (table 3 and figure 2).

Association of serum albumin with AD-signature neurodegeneration and WMH

Serum albumin level (as a continuous variable) was not related to AD-CM, AD-CT, or WMH (table 2 and figure 1, B–D). Moreover, no differences were observed in AD-CM, AD-CT, or WMH between the serum albumin categories (table 3).

Sensitivity analyses

The same analyses for the individuals with no decrease in food intake over the past 3 months produced similar results for A β deposition (or A β positivity), AD-CM, AD-CT, and WMH (tables 4 and 5 and figure 3).

Discussion

The results of the present study showed that low serum albumin was associated with an increased cerebral A β positivity rate compared to high serum albumin in older adults without dementia. The findings presented here were consistent with results of previous human studies on the relationships between serum albumin level and cognitive decline or AD dementia.

Table 4 Relationships between serum albumin and neuroimaging biomarkers in individuals with no recent decrease in food intake

Serum albumin	OR	95% CI	p Value
Dependent variable: Aβ positivity (n = 320)			
Model 1 ^a	0.10	0.03 to 0.35	<0.001
Model 2 ^b	0.15	0.04 to 0.57	0.005
Model 3 ^c	0.18	0.04 to 0.70	0.014
	B	95% CI	p Value
Dependent variable: global Aβ retention (n = 320)			
Model 1 ^a	−0.15	−0.24 to −0.07	<0.001
Model 2 ^b	−0.11	−0.19 to −0.03	0.006
Model 3 ^c	−0.10	−0.18 to −0.02	0.014
Dependent variable: AD-CM (n = 320)			
Model 1 ^a	0.01	−0.05 to 0.07	0.792
Model 2 ^b	−0.01	−0.07 to 0.05	0.768
Model 3 ^c	−<0.01	−0.06 to 0.06	0.903
Dependent variable: AD-CT (n = 314)			
Model 1 ^a	0.09	−<0.01 to 0.18	0.060
Model 2 ^b	0.03	−0.06 to 0.11	0.555
Model 3 ^c	0.01	−0.07 to 0.09	0.836
Dependent variable: WMH volume (n = 286)			
Model 1 ^a	0.46	−2.31 to 3.23	0.743
Model 2 ^b	1.01	−1.76 to 3.78	0.474
Model 3 ^c	1.08	−1.78 to 3.94	0.460

Abbreviations: Aβ = β-amyloid protein; AD-CM = Alzheimer disease–signature cerebral glucose metabolism; AD-CT = Alzheimer disease–signature cortical thickness; CI = confidence interval; OR = odds ratio; WMH = white matter hyperintensities. Global Aβ retention was used after natural log-transformation to achieve a normal distribution.

^a Not adjusted.

^b Adjusted for age, sex, and APOE ε4.

^c Adjusted for age, sex, APOE ε4, education, clinical diagnosis, medication use, alcohol use, annual income, diabetes mellitus, hyperlipidemia, liver disease, and kidney disease.

One population-based longitudinal study showed that low albumin within the normal range was an independent risk marker for cognitive decline in community-living older adults.³⁴ Previous clinical studies in hospital inpatients also showed that low serum albumin was associated with cognitive impairment^{11,39} and late-onset AD dementia compared with controls.^{14,40,41}

The inverse association of serum albumin with Aβ deposition observed in the present study supports the so-called sink hypothesis,^{42,43} which suggests that Aβ shifts from the brain into the blood plasma, forming a decrease in the Aβ concentration gradient between the brain and blood plasma. In this process, serum albumin binds most of the Aβ in the blood plasma and, in turn, lowers the blood plasma Aβ concentration. Therefore, serum albumin plays an important role in the

Aβ shift from the brain to the blood plasma for balancing the dynamic equilibrium of Aβ between the brain and blood plasma. Low serum albumin results in less binding to Aβ in blood, which increases blood plasma Aβ concentration, resulting in increased Aβ deposition in the brain by blocking the Aβ shift due to its lower concentration difference. An alternative hypothesis suggests that serum albumin is an endogenous inhibitor of Aβ fibril formation.^{7,10} Serum albumin may act on Aβ in 3 ways—monomer stabilizer, dissociation catalyst, or monomer competitor—and thus may prevent fibril formation. However, the possibility of reverse causality should also be considered. Poor nutrition in patients with dementia can reduce serum albumin level. This may explain the relationships between low serum albumin and AD dementia reported previously.^{14,15} Because all of our subjects were without dementia, however, reverse causality could not

Table 5 Relationships between stratified serum albumin and neuroimaging biomarkers in individuals with no recent decrease in food intake

	Stratified serum albumin				
	Low (<4.4 g/dL)		Middle (4.4–4.5 g/dL)		High (>4.5 g/dL)
	OR (95% CI)	<i>p</i> Value	OR (95% CI)	<i>p</i> Value	
Aβ positivity (n = 320)					
Model 1 ^a	4.76 (2.22–10.21)	<0.001	1.86 (0.82–4.21)	0.135	Reference
Model 2 ^b	3.80 (1.72–8.40)	0.001	1.97 (0.85–4.57)	0.116	Reference
Model 3 ^c	3.30 (1.43–7.61)	0.005	2.19 (0.89–5.39)	0.087	Reference
	B (95% CI)	<i>p</i> Value	B (95% CI)	<i>p</i> Value	
Global Aβ retention (n = 320)					
Model 1 ^a	0.10 (0.06 to 0.15)	<0.001	0.01 (–0.03 to 0.06)	0.593	Reference
Model 2 ^b	0.08 (0.03 to 0.13)	0.001	0.01 (–0.03 to 0.05)	0.647	Reference
Model 3 ^c	0.06 (0.02 to 0.11)	0.007	0.01 (–0.03 to 0.05)	0.639	Reference
AD-CM (n = 320)					
Model 1 ^a	–0.02 (–0.05 to 0.02)	0.287	–0.01 (–0.04 to 0.03)	0.713	Reference
Model 2 ^b	–0.01 (–0.04 to 0.03)	0.628	–<0.01 (–0.04 to 0.03)	0.813	Reference
Model 3 ^c	–0.01 (–0.05 to 0.03)	0.590	–<0.01 (–0.04 to 0.03)	0.816	Reference
AD-CT (n = 314)					
Model 1 ^a	–0.08 (–0.13 to –0.02)	0.005	0.01 (–0.04 to 0.06)	0.774	Reference
Model 2 ^b	–0.04 (–0.09 to 0.01)	0.111	0.01 (–0.03 to 0.06)	0.641	Reference
Model 3 ^c	–0.03 (–0.07 to 0.02)	0.297	0.02 (–0.03 to 0.06)	0.499	Reference
WMH (n = 286)					
Model 1 ^a	0.39 (–1.23 to 2.00)	0.639	–0.17 (–1.68 to 1.35)	0.828	Reference
Model 2 ^b	0.02 (–1.60 to 1.64)	0.982	–0.38 (–1.88 to 1.11)	0.613	Reference
Model 3 ^c	–0.03 (–1.71 to 1.64)	0.969	–0.41 (–1.95 to 1.13)	0.598	Reference

Abbreviations: Aβ = β-amyloid protein; AD-CM = Alzheimer disease–signature cerebral glucose metabolism; AD-CT = Alzheimer disease–signature cortical thickness; CI = confidence interval; OR = odds ratio; WMH = white matter hyperintensities.

Global Aβ retention was used after natural log-transformation to achieve a normal distribution.

^a Not adjusted.

^b Adjusted for age, sex, and APOE ε4.

^c Adjusted for age, sex, APOE ε4, education, clinical diagnosis, medication use, alcohol use, annual income, diabetes mellitus, hyperlipidemia, liver disease, and kidney disease.

explain our observation of the relationship between albumin and Aβ deposition.

In terms of Aβ positivity risk, the odds ratio for low albumin category was 3.40 (95% confidence interval 1.67–6.92, $p = 0.001$) compared with the high albumin category. This means that, even within the normal range of serum albumin, individuals with serum albumin <4.4 g/dL have an ≈3 times higher risk of pathologic Aβ deposition. This finding also indicates that nutritional interventions such as high dietary protein or albumin replacement may be helpful for those with low serum albumin in terms of AD prevention or treatment. Recently, the Alzheimer's Management by Albumin Replacement (AMBAR) trial was conducted to

demonstrate the clinical efficacy of therapeutic albumin replacement with plasma exchange as a new treatment of AD.^{44,45} Our result may provide an additional rationale for this kind of trial.

Unlike the relationship with Aβ, serum albumin was not related to either AD-signature neurodegeneration or WMH. While the relationships between low serum albumin and AD dementia have been reported as mentioned above, very little information is available on the relationships of serum albumin with degenerative changes in the brain.^{14,15} The lack of association between serum albumin and AD-CM or AD-CT in the present study indicated that serum albumin does not directly affect metabolism or structural changes in

Figure 3 Partial correlation plot for the relationship of serum albumin with global A β retention in individuals with no recent decrease in food intake

Multiple linear regression analysis was performed after adjustment for age, sex, and APOE ϵ 4. Global β -amyloid (A β) retention was used after natural log-transformation to achieve a normal distribution.

AD-related brain regions in older adults. We did not find any relationship between serum albumin and WMH. In addition, some human studies have reported an inverse association between serum albumin and cerebrovascular disease.^{16,17} The discrepancy may be related to the difference in range of serum albumin concentrations. While our subjects had serum albumin levels within the normal range, the previous studies mentioned above included subjects with serum albumin levels below the normal range (total levels 3.0–5.7 g/dL, low levels 3.0–3.9 g/dL in 1 study¹⁶; total levels 2.7–5.5 g/dL, low levels 2.7–4.2 g/dL in the other¹⁷; and total levels 3.6–5.3 g/dL, low levels 3.6–4.3 g/dL in our study). Therefore, serum albumin level below the normal range may increase cerebrovascular injury, as shown in the previous studies, while low serum albumin within the normal range may have no such effect.

The present study had several limitations that should be considered. First, because this was a cross-sectional study, we could not confirm a causal relationship between chronic albumin status in blood and brain A β deposition. Some findings support the long-term stability of serum albumin level in healthy older adults.^{46–49} We also excluded individuals with severe medical conditions that could affect mental function. In addition, the results were not changed even when the individuals with decreased food intake were excluded. Together, our results may support a long-term causal effect of serum albumin on brain A β deposition. Nevertheless, further long-term follow-up studies are required to clarify the causal relationships. In addition, the lack of repeated assessments of serum albumin level may have resulted in measurement errors because of diurnal variation.⁵⁰ However, to minimize such errors, all blood samples for serum albumin measurement

were obtained at the same time of the day (8–9 AM) in all participants.

The findings of present study suggest that low serum albumin may increase the risk of AD dementia by elevating amyloid accumulation. In terms of AD prevention, more attention needs to be paid to avoid low serum albumin level, even within the clinical normal range, by clinicians.

Study funding

This study was supported by a grant from the Ministry of Science, ICT, and Future Planning, Republic of Korea (grant NRF-2014M3C7A1046042) and a grant from the Korea Health Technology R&D Project through the Korea Health Industry Development Institute, funded by the Ministry of Health & Welfare, Republic of Korea (grant HI18C0630 & HI19C0149).

Disclosure

The authors report no disclosures relevant to the manuscript. Go to Neurology.org/N for full disclosures.

Publication history

Received by *Neurology* May 14, 2019. Accepted in final form January 27, 2020.

Appendix 1 Authors

Name	Location	Contribution
Jee Wook Kim, MD, PhD	Hallym University Dongtan Sacred Heart Hospital, Hwaseong, Republic of Korea	Study concept and design; analysis and interpretation of data; statistical analysis; and drafting and critically revising the manuscript for intellectual content
Min Soo Byun, MD, PhD	Medical Research Center Seoul National University, Republic of Korea	Acquisition, analysis, and interpretation of data; statistical analysis; and critically revising the manuscript for intellectual content
Jun Ho Lee, MD	Seoul National University Hospital, Republic of Korea	Acquisition, analysis, and interpretation of data; and critically revising the manuscript for intellectual content
Dahyun Yi, PhD	Medical Research Center Seoul National University, Republic of Korea	Acquisition, analysis, and interpretation of data; statistical analysis; and critically revising the manuscript for intellectual content
So Yeon Jeon	Chungnam National University Hospital, Daejeon, Republic of Korea	Acquisition, analysis, and interpretation of data; and critically revising the manuscript for intellectual content
Bo Kyung Sohn, MD	Sanggye Paik Hospital, Inje University College of Medicine, Seoul, Republic of Korea	Acquisition, analysis, and interpretation of data; and critically revising the manuscript for intellectual content

Appendix 1 (continued)

Name	Location	Contribution
Jun-Young Lee, MD, PhD	SMG-SNU Boramae Medical Center, Seoul National University College of Medicine, Republic of Korea	Acquisition, analysis, and interpretation of data; and critically revising the manuscript for intellectual content
Seong A Shin, BA	SMG-SNU Boramae Medical Center, Seoul, Republic of Korea	Acquisition, analysis, and interpretation of data; and critically revising the manuscript for intellectual content
Yu Kyeong Kim, MD, PhD	SMG-SNU Boramae Medical Center, Seoul, Republic of Korea	Acquisition, analysis, and interpretation of data; and critically revising the manuscript for intellectual content
Koung Mi Kang, MD	Seoul National University Hospital, Republic of Korea	Acquisition, analysis, and interpretation of data; and critically revising the manuscript for intellectual content
Chul-Ho Sohn, MD, PhD	Seoul National University Hospital, Republic of Korea	Acquisition, analysis, and interpretation of data; and critically revising the manuscript for intellectual content
Dong Young Lee, MD, PhD	Seoul National University Hospital, Seoul National University College of Medicine, Republic of Korea	Study concept and design; acquisition, analysis, and interpretation of data; statistical analysis; and drafting and critically revising the manuscript for intellectual content

Appendix 2 Coinvestigators

Coinvestigators are listed at links.lww.com/WNL/B134.

References

- Cummings JL. Alzheimer's disease. *N Engl J Med* 2004;351:56–67.
- Tobiansky R, Blizard R, Livingston G, Mann A. The Gospel Oak Study stage IV: the clinical relevance of subjective memory impairment in older people. *Psychol Med* 1995;25:779–786.
- Braak H, Braak E. Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol* 1991;82:239–259.
- Jack CR, Jr, Knopman DS, Jagust WJ, et al. Hypothetical model of dynamic biomarkers of the Alzheimer's pathological cascade. *Lancet Neurol* 2010;9:119–128.
- Carter DC, Ho JX. Structure of serum albumin. *Adv Protein Chem* 1994;45:153–203.
- Biere AL, Ostaszewski B, Stimson ER, Hyman BT, Maggio JE, Selkoe DJ. Amyloid beta-peptide is transported on lipoproteins and albumin in human plasma. *J Biol Chem* 1996;271:32916–32922.
- Kuo YM, Kokjohn TA, Kalback W, et al. Amyloid-beta peptides interact with plasma proteins and erythrocytes: implications for their quantitation in plasma. *Biochem Biophys Res Commun* 2000;268:750–756.
- Zlokovic BV. Clearing amyloid through the blood-brain barrier. *J Neurochem* 2004;89:807–811.
- Stanyon HF, Viles JH. Human serum albumin can regulate amyloid-beta peptide fiber growth in the brain interstitium: implications for Alzheimer disease. *J Biol Chem* 2012;287:28163–28168.
- Milosevic J, Raditsis A, Melacini G. Human serum albumin inhibits Aβ fibrillization through a "monomer-competitor" mechanism. *Biophys J* 2009;97:2585–2594.
- Zuccala G, Marzetti E, Cesari M, et al. Correlates of cognitive impairment among patients with heart failure: results of a multicenter survey. *Am J Med* 2005;118:496–502.
- Mizrahi EH, Blumstein T, Arad M, Adunsky A. Serum albumin levels predict cognitive impairment in elderly hip fracture patients. *Am J Alzheimers Dis Other Dement* 2008;23:85–90.
- Llewellyn DJ, Langa KM, Friedland RP, Lang IA. Serum albumin concentration and cognitive impairment. *Curr Alzheimer Res* 2010;7:91–96.
- Maes M, De Vos N, Wauters A, et al. Inflammatory markers in younger vs elderly normal volunteers and in patients with Alzheimer's disease. *J Psychiatr Res* 1999;33:397–405.
- Kim TS, Pae CU, Yoon SJ, et al. Decreased plasma antioxidants in patients with Alzheimer's disease. *Int J Geriatr Psychiatry* 2006;21:344–348.
- Xu WH, Dong C, Rundek T, Elkind MS, Sacco RL. Serum albumin levels are associated with cardioembolic and cryptogenic ischemic strokes: Northern Manhattan Study. *Stroke* 2014;45:973–978.
- Phillips A, Shaper AG, Whincup PH. Association between serum albumin and mortality from cardiovascular disease, cancer, and other causes. *Lancet* 1989;2:1434–1436.
- Byun MS, Yi D, Lee JH, et al. Korean brain aging study for the early diagnosis and prediction of Alzheimer's disease: methodology and baseline sample characteristics. *Psychiatry Invest* 2017;14:851–863.
- Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules. *Neurology* 1993;43:2412–2414.
- Lee DY, Lee KU, Lee JH, et al. A normative study of the CERAD neuropsychological assessment battery in the Korean elderly. *J Int Neuropsychol Soc* 2004;10:72–81.
- Morris JC, Heyman A, Mohs RC, et al. The Consortium to Establish a Registry for Alzheimer's Disease (CERAD), part I: clinical and neuropsychological assessment of Alzheimer's disease. *Neurology* 1989;39:1159–1165.
- Lee JH, Lee KU, Lee DY, et al. Development of the Korean version of the Consortium to Establish a Registry for Alzheimer's Disease assessment packet (CERAD-K): clinical and neuropsychological assessment batteries. *J Gerontol B Psychol Sci Soc Sci* 2002;57:P47–P53.
- Vellas B, Guigoz Y, Garry PJ, et al. The Mini Nutritional Assessment (MNA) and its use in grading the nutritional state of elderly patients. *Nutrition* 1999;15:116–122.
- Kuller LH, Eichner JE, Orchard TJ, Grandits GA, McCallum L, Tracy RP. The relation between serum albumin levels and risk of coronary heart disease in the Multiple Risk Factor Intervention Trial. *Am J Epidemiol* 1991;134:1266–1277.
- Dik MG, Jonker C, Hack CE, Smit JH, Comijs HC, Eikelenboom P. Serum inflammatory proteins and cognitive decline in older persons. *Neurology* 2005;64:1371–1377.
- Wenham PR, Price WH, Blandell G. Apolipoprotein E genotyping by one-stage PCR. *Lancet* 1991;337:1158–1159.
- Park JC, Han SH, Yi D, et al. Plasma tau/amyloid-beta1-42 ratio predicts brain tau deposition and neurodegeneration in Alzheimer's disease. *Brain* 2019;142:771–786.
- Reiman EM, Chen K, Liu X, et al. Fibrillar amyloid-beta burden in cognitively normal people at 3 levels of genetic risk for Alzheimer's disease. *Proc Natl Acad Sci USA* 2009;106:6820–6825.
- Choe YM, Sohn BK, Choi HJ, et al. Association of homocysteine with hippocampal volume independent of cerebral amyloid and vascular burden. *Neurobiol Aging* 2014;35:1519–1525.
- Westwood S, Baird AL, Hye A, et al. Plasma protein biomarkers for the prediction of CSF amyloid and tau and [(18)F]-Flutemetamol PET scan result. *Front Aging Neurosci* 2018;10:409.
- Janssen L, Sobott F, De Deyn PP, Van Dam D. Signal loss due to oligomerization in ELISA analysis of amyloid-beta can be recovered by a novel sample pre-treatment method. *MethodsX* 2015;2:112–123.
- Jack CR, Jr, Wiste HJ, Weigand SD, et al. Age-specific population frequencies of cerebral beta-amyloidosis and neurodegeneration among people with normal cognitive function aged 50–89 years: a cross-sectional study. *Lancet Neurol* 2014;13:997–1005.
- Tsai JZ, Peng SJ, Chen YW, et al. Automated segmentation and quantification of white matter hyperintensities in acute ischemic stroke patients with cerebral infarction. *PLoS One* 2014;9:e104011.
- Ng TP, Niti M, Feng L, Kua EH, Yap KB. Albumin, apolipoprotein E-ε4 and cognitive decline in community-dwelling Chinese older adults. *J Am Geriatr Soc* 2009;57:101–106.
- Weiner DE, Bartolomei K, Scott T, et al. Albuminuria, cognitive functioning, and white matter hyperintensities in homebound elders. *Am J Kidney Dis* 2009;53:438–447.
- Ota A, Kondo N, Murayama N, et al. Serum albumin levels and economic status in Japanese older adults. *PLoS One* 2016;11:e0155022.
- Folsom AR, Ma J, Eckfeldt JH, Nieto FJ, Metcalf PA, Barnes RW. Low serum albumin: association with diabetes mellitus and other cardiovascular risk factors but not with prevalent cardiovascular disease or carotid artery intima-media thickness: the Atherosclerosis Risk in Communities (ARIC) Study Investigators. *Ann Epidemiol* 1995;5:186–191.
- Sui M, Jia X, Yu C, et al. Relationship between hypoalbuminemia, hyperlipidemia and renal severity in patients with lupus nephritis: a prospective study. *Cent Eur J Immunol* 2014;39:243–252.
- Cattin L, Bordin P, Fonda M, et al. Factors associated with cognitive impairment among older Italian inpatients: Gruppo Italiano di Farmacovigilanza nell'Anziano (G.I.F.A.). *J Am Geriatr Soc* 1997;45:1324–1330.
- Hampel H, Kotter HU, Moller HJ. Blood-cerebrospinal fluid barrier dysfunction for high molecular weight proteins in Alzheimer disease and major depression: indication for disease subsets. *Alzheimer Dis Assoc Disord* 1997;11:78–87.
- Jeandel C, Nicolas MB, Dubois F, Nabet-Belleville F, Penin F, Cuny G. Lipid peroxidation and free radical scavengers in Alzheimer's disease. *Gerontology* 1989;35:275–282.
- DeMattos RB, Bales KR, Cummins DJ, Paul SM, Holtzman DM. Brain to plasma amyloid-beta efflux: a measure of brain amyloid burden in a mouse model of Alzheimer's disease. *Science* 2002;295:2264–2267.
- Boada M, Ortiz P, Anaya F, et al. Amyloid-targeted therapeutics in Alzheimer's disease: use of human albumin in plasma exchange as a novel approach for Aβ mobilization. *Drug News Perspect* 2009;22:325–339.

44. Boada M, Lopez O, Nunez L, et al. Plasma exchange for Alzheimer's Disease Management by Albumin Replacement (AMBAR) trial: study design and progress. *Alzheimers Dement* 2019;5:61–69.
45. Boada M, Ramos-Fernandez E, Guivernau B, et al. Treatment of Alzheimer disease using combination therapy with plasma exchange and haemapheresis with albumin and intravenous immunoglobulin: rationale and treatment approach of the AMBAR (Alzheimer Management by Albumin Replacement) study. *Neurologia* 2016;31:473–481.
46. Campion EW, deLabry LO, Glynn RJ. The effect of age on serum albumin in healthy males: report from the Normative Aging Study. *J Gerontol* 1988;43:M18–M20.
47. Leask RG, Andrews GR, Caird FL. Normal values for sixteen blood constituents in the elderly. *Age Ageing* 1973;2:14–23.
48. Reed AH, Cannon DC, Winkelman JW, Bhasin YP, Henry RJ, Pileggi VJ. Estimation of normal ranges from a controlled sample survey, I: sex- and age-related influence on the SMA 12-60 screening group of tests. *Clin Chem* 1972;18:57–66.
49. Misra DP, Loudon JM, Staddon GE. Albumin metabolism in elderly patients. *J Gerontol* 1975;30:304–306.
50. Andersen IB, Brasen CL, Christensen H, et al. Standardised resting time prior to blood sampling and diurnal variation associated with risk of patient misclassification: results from selected biochemical components. *PLoS One* 2015;10:e0140475.

Neurology®

Serum albumin and beta-amyloid deposition in the human brain

Jee Wook Kim, Min Soo Byun, Jun Ho Lee, et al.

Neurology 2020;95:e815-e826 Published Online before print July 20, 2020

DOI 10.1212/WNL.0000000000010005

This information is current as of July 20, 2020

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/95/7/e815.full
References	This article cites 50 articles, 9 of which you can access for free at: http://n.neurology.org/content/95/7/e815.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Alzheimer's disease http://n.neurology.org/cgi/collection/alzheimers_disease PET http://n.neurology.org/cgi/collection/pet
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.