



Abstracts

Articles appearing in the December 2020 issue

New Recessive Mutations in *SYT2* Causing Severe Presynaptic Congenital Myasthenic Syndromes

Objective To report the identification of 2 new homozygous recessive mutations in the synaptotagmin 2 (*SYT2*) gene as the genetic cause of severe and early presynaptic forms of congenital myasthenic syndromes (CMSs).

Methods Next-generation sequencing identified new homozygous intronic and frameshift mutations in the *SYT2* gene as a likely cause of presynaptic CMS. We describe the clinical and electromyographic patient phenotypes, perform ex vivo splicing analyses to characterize the effect of the intronic mutation on exon splicing, and analyze the functional impact of this variation at the neuromuscular junction (NMJ).

Results The 2 infants presented a similar clinical phenotype evoking first a congenital myopathy characterized by muscle weakness and hypotonia. Next-generation sequencing allowed to the identification of 1 homozygous intronic mutation c.465+1 G>A in patient 1 and another homozygous frameshift mutation c.328_331dup in patient 2, located respectively in the 5' splice donor site of *SYT2* intron 4 and in exon 3. Functional studies of the intronic mutation validated the abolition of the splice donor site of exon 4 leading to its skipping. In-frame skipping of exon 4 that encodes part of the C2A calcium-binding domain of *SYT2* is associated with a loss-of-function effect resulting in a decrease of neurotransmitter release and severe pre- and postsynaptic NMJ defects.

Conclusions This study identifies new homozygous recessive *SYT2* mutations as the underlying cause of severe and early presynaptic form of CMS expanding the genetic spectrum of recessive *SYT2*-related CMS associated with defects in neurotransmitter release.

[NPub.org/NG/9615a](https://pub.org/NG/9615a)

Genotype-Phenotype Correlations in Patients With de Novo *KCNQ2* Pathogenic Variants

Objective Early identification of de novo *KCNQ2* variants in patients with epilepsy raises prognostic issues toward optimal management. We analyzed the clinical and genetic information from a cohort of patients with de novo *KCNQ2* pathogenic variants to dissect genotype-phenotype correlations.

Methods Patients with de novo *KCNQ2* pathogenic variants were identified from Italy, Denmark, and Belgium. Atomic resolution Kv7.2 structures were also generated using homology modeling to map the variants.

Results We included 34 patients with a mean age of 4.7 years. Median seizure onset was 2 days, mainly with focal seizures with autonomic signs. Twenty-two patients (65%) were seizure free at the mean age of 1.2 years. More than half of the patients (17/32) displayed severe/profound intellectual disability; however, 4 (13%) of them had a normal cognitive outcome.

A total of 28 de novo pathogenic variants were identified, most missense (25/28), and clustered in conserved regions of the protein; 6 variants recurred, and 7 were novel. We did not identify a relationship between variant position and seizure offset or cognitive outcome in patients harboring missense variants. Besides, recurrent variants were associated with overlapping epilepsy features but also variable evolution regarding the intellectual outcome.

Conclusions We highlight the complexity of variant interpretation to assess the impact of a class of de novo *KCNQ2* mutations. Genetic modifiers could be implicated, but the study paradigms to successfully address the impact of each single mutation need to be developed.

[NPub.org/NG/9615b](https://pub.org/NG/9615b)



Most-Read Articles

As of February 19, 2021

Homozygous deletion in *MICU1* presenting with fatigue and lethargy in childhood

D. Lewis-Smith, K. J. Kamer, H. Griffin, et al. 2016;e59. doi.org/10.1212/NXG.0000000000000059

KCNQ2 encephalopathy: Features, mutational hot spots, and ezogabine treatment of 11 patients

J.J. Millichap, K.L. Park, T. Tsuchida, et al. 2016;2:e96. doi.org/10.1212/NXG.0000000000000096

Mendelian randomization shows a causal effect of low vitamin D on multiple sclerosis risk

B. Rhead, M. Bäärnhielm, M. Gianfrancesco, et al. 2016;2:e97. doi.org/10.1212/NXG.0000000000000097

GBA p.T369M substitution in Parkinson disease: Polymorphism or association? A meta-analysis

V. Mallett, J. P. Ross, R. N. Alcalay, et al. 2016;2:e104. doi.org/10.1212/NXG.0000000000000104

The Alzheimer's Disease Sequencing Project: Study design and sample selection

G.W. Beecham, J.C. Bis, E.R. Martin, et al. 2017;3:e194. doi.org/10.1212/NXG.0000000000000194

Neurology®

What's Happening in *Neurology*® *Genetics*
Neurology 2021;96;703
DOI 10.1212/WNL.00000000000011778

This information is current as of April 12, 2021

Updated Information & Services

including high resolution figures, can be found at:
<http://n.neurology.org/content/96/15/703.full>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
http://www.neurology.org/about/about_the_journal#permissions

Reprints

Information about ordering reprints can be found online:
<http://n.neurology.org/subscribers/advertise>

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2021 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.

