Teaching NeuroImages: Ohtahara Syndrome due to Unilateral Perisylvian Polymicrogyria

Sabarish Sekar, MD, DNB, Pavankumar Rudrabhatla, MD, Vinayagamani Selvadasan, MD, DNB, DM, Bejoy Thomas, MD, DNB, PDCC, and Chandrasekharan Kesavadas, MD

Correspondence

Dr. Sekar
sabarish.ss.dr@gmail.com

Neurology® 2021;96:e2456-e2457. doi:10.1212/WNL.00000000000011497

Figure MRI and EEG

Coronal T2 (A) and coronal T1 (B, C) images show left perisylvian polymicrogyria (A), abnormally thick and bumpy coarse appearance of cortex (B), and widening of sylvian fissure (C). (D) Arterial spin labeling (postlabeling delay 2,500 ms) image shows left perisylvian hypoperfusion corresponding to polymicrogyria (arrowhead). (E) EEG shows typical burst-suppression pattern (time base: 20 mm/s, gain: 10 μ V/mm, high-frequency filter: 70 Hz, low-frequency filter: 1 Hz).

A 10-month-old boy, born of nonconsanguineous parents, without any perinatal insult, presented with tonic spasms from the 13th day of life. He started having tonic seizures from 4 months of age, with global developmental delay. On examination, there were no neurocutaneous markers, facial dysmorphism, or focal neurologic deficits. EEG (figure) showed generalized and multifocal epileptiform discharges along with burst-suppression pattern. MRI of brain revealed left perisylvian polymicrogyria (figure).

Ohtahara syndrome is an electroclinical syndrome, characterized by infantile-onset epileptic encephalopathy and typical burst-suppression pattern on EEG, which remains unchanged during sleep and wakefulness.¹ It is most commonly associated with structural malformations including neuronal migration disorder or dysgenesis, mutations in genes including *ARX*, *CDKL5*, *SLC25A22*, *STXBP1*, and *KCNQ2*, and various metabolic disorders.^{1,2} Presence of structural malformations, like left-sided perisylvian polymicrogyria in our case, precludes

MORE ONLINE

→Teaching slides

<http://links.lww.com/WNL/B303>

From the Department of Imaging Sciences and Interventional Radiology (S.S., B.T., C.K.), Department of Neurology (P.R.), and Department of Imaging Sciences and Interventional Radiology (V.S.), Sree Chitra Tirunal Institute for Medical Sciences and Technology, Trivandrum, India.

Go to Neurology.org/N for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

extensive genetic and metabolic analysis and differentiates it from early myoclonic encephalopathy.

Study Funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Appendix Authors

Name	Location	Contribution
Sabarish Sekar, MD, DNB	Department of Imaging Sciences and Intervention Radiology, Sree Chitra Tirunal Institute for Medical Sciences and Technology, Thiruvananthapuram, Kerala, India	Major role in acquisition of data, study concept, design and drafting of the manuscript
Pavankumar Rudrabhatla, MD	Department of Neurology, Sree Chitra Tirunal Institute for Medical Sciences and Technology, Thiruvananthapuram, Kerala, India	Major role in acquisition of data and EEG discussion

Appendix (continued)

Name	Location	Contribution
Vinayagamani Selvadasan, MD, DNB, DM	Department of Imaging Sciences and Intervention Radiology, Sree Chitra Tirunal Institute for Medical Sciences and Technology, Thiruvananthapuram, Kerala, India	Study concept and design, revision of manuscript
Bejoy Thomas, MD, DNB, PDCC	Department of Imaging Sciences and Intervention Radiology, Sree Chitra Tirunal Institute for Medical Sciences and Technology, Thiruvananthapuram, Kerala, India	Critical revision of manuscript
Chandrasekharan Kesavadas, MD	Department of Imaging Sciences and Intervention Radiology, Sree Chitra Tirunal Institute for Medical Sciences and Technology, Thiruvananthapuram, Kerala, India	Critical revision of manuscript

References

1. Beal JC, Cherian K, Moshe SL. Early-onset epileptic encephalopathies: Ohtahara syndrome and early myoclonic encephalopathy. *Pediatr Neurol* 2012;47:317–323.
2. Pavone P, Spalice A, Polizzi A, Parisi P, Ruggieri M. Ohtahara syndrome with emphasis on recent genetic discovery. *Brain Dev* 2012;34:459–468.

Neurology®

Teaching NeuroImages: Ohtahara Syndrome due to Unilateral Perisylvian Polymicrogyria

Sabarish Sekar, Pavankumar Rudrabhatla, Vinayagamani Selvasadan, et al.
Neurology 2021;96:e2456-e2457 Published Online before print January 5, 2021
DOI 10.1212/WNL.00000000000011497

This information is current as of January 5, 2021

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/96/19/e2456.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/96/19/e2456.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Cortical dysplasia http://n.neurology.org/cgi/collection/cortical_dysplasia EEG http://n.neurology.org/cgi/collection/eeg_ Infantile spasms http://n.neurology.org/cgi/collection/infantile_spasms
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2021 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.