Bilateral Thalamic Lesions Associated With Atezolizumab-Induced Autoimmune Encephalitis

Haruo Nishijima, MD, PhD, Chieko Suzuki, MD, PhD, Tomoya Kon, MD, PhD, Takashi Nakamura, MD, Hisashi Tanaka, MD, PhD, Yui Sakamoto, MD, and Masahiko Tomiyama, MD, PhD

Neurology® 2021;96:126-127. doi:10.1212/WNL.00000000000011297

Correspondence

Dr. Nishijima

hnishijima-ky@umin.ac.jp

Figure Reversible Thalamic Lesions With Atezolizumab-Induced Encephalitis Before and After Immunotherapy

MRI initially showed high signals bilaterally in thalamus on T2-weighted (A), fluid-attenuated inversion recovery (B), and diffusion-weighted images (C), and apparent diffusion coefficient map (D). Lesion size reduced on fluid-attenuated inversion recovery images 3 (E) and 7 weeks (F) after immunotherapy.

A 72-year-old woman presented with a 6-week history of gait disturbance and mild disturbance of consciousness. For 9 months, she had been treated for advanced non-small cell lung cancer with atezolizumab. MRI revealed symmetrical high signal in the thalamus bilaterally (figure). She did not have evidence of cancer recurrence or metastases. Serum autoimmune antibodies were absent (anti-AQP4, anti-MOG, anti-amphiphysin, CV2, PNMA2 [Ma2/Ta], Ri, Yo, Hu, recoverin, SOX1, titin, zic4, GAD65, and Tr [DNER]). In the CSF, she had high immunoglobulin G index and positive oligoclonal bands but normal myelin basic protein, absent anti-NMDA receptor and anti-MOG antibodies, and no evidence of infection. Cytology was normal. She was diagnosed with autoimmune encephalitis associated with atezolizumab treatment and

From the Department of Neurology, Institute of Brain Science (H.N., C.S., T.K., T.N., M.T.), Department of Respiratory Medicine (H.T.), and Department of Neuropsychiatry (Y.S.), Hirosaki University Graduate School of Medicine; and Department of Neurology (H.N.), Hirosaki University Hospital, Hirosaki City, Aomori, Japan.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

treated with steroids and IV immunoglobulin 9 weeks after the onset of symptoms. The size of the lesions decreased after immunotherapy started, but she remained bedridden.

Study Funding

No targeted funding reported.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Appendix Authors

Name	Location	Contribution
Haruo Nishijima, MD, PhD	Hirosaki University Graduate School of Medicine, Japan	Designed and conceptualized study, analyzed the data, drafted the manuscript for intellectual content
Chieko Suzuki, MD, PhD	Hirosaki University Graduate School of Medicine, Japan	Patient care and acquisition of data, interpreted the data, revised the manuscript for intellectual content

Appendix (continued)

Name	Location	Contribution
Tomoya Kon, MD, PhD	Hirosaki University Graduate School of Medicine, Japan	Patient care and acquisition of data, revised the manuscript for intellectual content
Takashi Nakamura, MD	Hirosaki University Graduate School of Medicine, Japan	Patient care and acquisition of data
Hisashi Tanaka, MD, PhD	Hirosaki University Graduate School of Medicine, Japan	Patient care and acquisition of data, revised the manuscript for intellectual content
Yui Sakamoto, MD	Hirosaki University Graduate School of Medicine, Japan	Patient care and acquisition of data, revised the manuscript for intellectual content
Masahiko Tomiyama, MD, PhD	Hirosaki University Graduate School of Medicine, Japan	Interpreted the data, revised the manuscript for intellectual content

NEW EPISODE

Increasing Out-of-Pocket Costs for Neurologic Care for Privately Insured Patients (see p. 94)

In the first segment, Dr. Stacey Clardy receives an update from Dr. Pushpa Narayanaswami on the management of myasthenia gravis. In the second part of the podcast, Dr. Jeffrey Ratliff talks with Dr. Chloe Hill about patients' out-of-pocket costs in neurology.

Disclosures can be found at [Neurology.org](https://www.neurology.org).

CME Opportunity: Listen to this week's Neurology® Podcast and earn 0.5 AMA PRA Category 1 CME Credits™ by completing the online podcast quiz.

Neurology®

Bilateral Thalamic Lesions Associated With Atezolizumab-Induced Autoimmune Encephalitis

Haruo Nishijima, Chieko Suzuki, Tomoya Kon, et al.

Neurology 2021;96:126-127 Published Online before print December 2, 2020

DOI 10.1212/WNL.0000000000011297

This information is current as of December 2, 2020

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/96/3/126.full
Citations	This article has been cited by 1 HighWire-hosted articles: http://n.neurology.org/content/96/3/126.full##otherarticles
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Clinical Neurology http://n.neurology.org/cgi/collection/all_clinical_neurology Autoimmune diseases http://n.neurology.org/cgi/collection/autoimmune_diseases Encephalitis http://n.neurology.org/cgi/collection/encephalitis MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2020 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.