ANGPTL6 Genetic Variants Are an Underlying Cause of Familial Intracranial Aneurysms

Isabel C. Hostettler, MD, Benjamin O'Callaghan, MSc, Enrico Bugiardini, MD, Emer O'Connor, MD, Jana Vandrovicova, PhD, Indran Davagnanam, MD, Varinder Alg, MD, Stephen Bonner, PhD, Daniel Walsh, PhD, Diederik Bulters, FRCS, Neil Kitchen, PhD, Martin M. Brown, FRCP, Joan Grieve, MD, David J. Werring, PhD, FRCP, and Henry Houlden, PhD FRCP, on behalf of the Genetics and Observational Subarachnoid Haemorrhage (GOSH) study investigators

Correspondence

Dr. Houlden
h.houlden@ucl.ac.uk

Neurology® 2021;96:e947-e955. doi:10.1212/WNL.0000000000001125

Abstract

Purpose

To understand the role of the angiopoietin-like 6 gene (*ANGPTL6*) in intracranial aneurysms (IAs), we investigated its role in a large cohort of familial IAs.

Methods

Individuals with family history of IA were recruited to the Genetic and Observational Subarachnoid Haemorrhage (GOSH) study. The *ANGPTL6* gene was sequenced using Sanger sequencing. Identified genetic variants were compared to a control population.

Results

We found 6 rare *ANGPTL6* genetic variants in 9/275 individuals with a family history of IA (3.3%) (5 missense mutations and 1 nonsense mutation leading to a premature stop codon), none present in controls. One of these had been previously reported: c.392A>T (p.Glu131Val) on exon 2; another was very close: c.332G>A (p.Arg111His). Two further genetic variants lie within the fibrinogen-like domain of the *ANGPTL6* gene, which may influence function or level of the *ANGPTL6* protein. The last 2 missense mutations lie within the coiled-coil domain of the *ANGPTL6* protein. All genetic variants were well conserved across species.

Conclusion

ANGPTL6 genetic variants are an important cause of IA. Defective or lack of *ANGPTL6* protein is therefore an important factor in blood vessel proliferation leading to IA; dysfunction of this protein is likely to cause abnormal proliferation or weakness of vessel walls. With these data, not only do we emphasize the importance of screening familial IA cases for *ANGPTL6* and other genes involved in IA, but also highlight the *ANGPTL6* pathway as a potential therapeutic target.

Classification of Evidence

This is a Class III study showing some specificity of presence of the *ANGPTL6* gene variant as a marker of familial intracranial aneurysms in a small subset of individuals with familial aneurysms.

MORE ONLINE

→ Class of Evidence

Criteria for rating therapeutic and diagnostic studies

[NPub.org/coe](https://npub.org/coe)

From the Stroke Research Centre (I.C.H., V.A., M.M.B., D.J.W.), MRC Centre for Neuromuscular Diseases (B.O., E.B.), and Department of Neuromuscular Disorders (E.B., J.V.), UCL Queen Square Institute of Neurology; Neurogenetics Laboratory (I.C.H., B.O., E.O., H.H.) and Departments of Neuroradiology (I.D.) and Neurosurgery (N.K., J.G.), the National Hospital for Neurology and Neurosurgery, University College London Hospitals NHS Foundation Trust, Queen Square, London; Department of Anaesthesia (S.B.), the James Cook University Hospital, Middlesbrough; Department of Neurosurgery (D.W.), King's College Hospital NHS Foundation Trust, London; and Department of Neurosurgery (D.B.), University Hospital Southampton NHS Foundation Trust.

Go to Neurology.org/N for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

Genetics and Observational Subarachnoid Haemorrhage (GOSH) study investigators coinvestigators are listed in the appendix 2 at the end of the article.

The Article Processing Charge was funded by Wellcome Trust and MRC.

This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Glossary

aSAH = aneurysmal subarachnoid hemorrhage; **ExAC** = Exome Aggregation Consortium; **FLD** = fibrinogen-like domain; **gnomAD** = Genome Aggregation Database; **GOSH** = Genetic and Observational Subarachnoid Haemorrhage; **IA** = intracranial aneurysm; **MAF** = minor allele frequency; **MRC** = Medical Research Council; **NMD** = nonsense-mediated decay; **SAH** = subarachnoid haemorrhage; **SIFT** = Sorting Intolerant From Tolerant; **UIA** = unruptured intracranial aneurysm.

Three percent of the general population harbor an unruptured intracranial aneurysm (UIA).¹ Ruptured UIAs cause aneurysmal subarachnoid hemorrhage (aSAH), a devastating condition with high mortality and morbidity and great socioeconomic burden.^{2,3} Despite efforts to discover factors associated with aneurysm formation and rupture, little is known about mechanisms driving aneurysm formation, growth, and rupture as well as the genetic background of this disease. Some inherited disorders, such as autosomal dominant polycystic kidney disease, have been associated with increased aneurysm formation as well as rupture.^{4,5} Increasing evidence exists that genetic factors also play a role in intracranial aneurysm (IA) formation in patients without underlying genetic diseases.^{6,7}

Most IAs are acquired lesions arising under hemodynamic stress and associated defective wall response.⁸ Disturbances in angiogenic factors can lead to changes in vessel structure and cerebral artery wall stability. Genes involved in angiogenesis have recurrently been investigated. The angiopoietin-like 6 (*ANGPTL6*) gene is a protein-coding gene belonging to the *ANGPTL* family, which has been associated with regulatory capacities in angiogenesis.^{9–11} *ANGPTL6* has been identified as a circulating proangiogenic factor increasing endothelial permeability.^{9,12} A recent publication found significant enrichment in rare coding variants within *ANGPTL6* in patients with familial IA.¹³ They demonstrated a reduction of 50% in the serum concentration of *ANGPTL6* in individuals heterozygous for c.1378A>T.¹³

We aimed to externally validate the previous findings, to search for additional rare variants, and to evaluate the rate of rare variants in a large cohort of individuals with family history of IA.

Methods

Patient Selection

For the main analysis, we included patients with available DNA and familial IA where at least 1 affected first- or second-degree relative had IA and was recruited into the Genetic and Observational Subarachnoid Haemorrhage (GOSH) study to evaluate whether previously reported variants would be detected in those with only 1 affected first- or second-degree family member as well (figure 1). Only 1 patient had only 1 affected first- or second-degree relative. Patients with unruptured or ruptured IA could be

included in this study. We compare patients with familial IA and the rest of the cohort in table 1 in order to highlight differences. A description of the whole cohort is published elsewhere.¹⁴ Written informed consent was obtained from all participants, or in case of lack of capacity, from a representative. We did not include patients with perimesencephalic subarachnoid hemorrhage (SAH) (defined by blood distribution mainly or only in the cisterns around the midbrain), absence of intracranial aneurysm, or SAH due to trauma as well as mycotic aneurysms. None of the patients included had been diagnosed with Ehlers-Danlos syndrome, Marfan syndrome, or polycystic kidney disease. IA was confirmed on CT/magnetic resonance angiography or digital subtraction angiography. Because the primary aim of this study was to replicate the association between IA and identifier variants in *ANGPTL6*, and genetic data from affected/unaffected relatives were not collected with the exception of a mother and daughter pair, segregation analysis was not performed.

The study was approved by the corresponding ethics committee (reference 09/H0716/S4).

We used brain biopsy controls from our own laboratory as well as Medical Research Council (MRC) brain exome controls, excluding individuals with known previous stroke (n = 59), to compare the frequency of rare genetic variants found in cases with a control cohort.

The primary research goal is to externally validate and evaluate additional rare variants in the *ANGPTL6* gene. This is a Class III evidence study.

Sanger Sequencing

We extracted genomic DNA and performed PCR to amplify exons 2–6 of the *ANGPTL6* gene followed by Sanger sequencing. Primer sequences can be provided upon request. We cleaned up the PCR product using Exo-Fast, before performing dye-terminator sequencing PCR with BigDye Terminator v3.1 (Thermo Fisher). The sequencing PCR product was cleaned using Sephadex (Sigma Aldrich) before being loaded onto the AB1 3730xl genetic analyzer (Applied Biosystems). We used SeqScape v.3.0 software (Applied Biosystems) for sequence analysis.

In accordance with the previous publication, we filtered out variants with a minor allele frequency (MAF) higher than 1% in the Genome Aggregation Database (gnomAD), v2.1.1.¹³

Figure 1 Patients and Controls Selection Flow Diagram

GOSH = Genetic and Observational Subarachnoid Haemorrhage; IA = intracranial aneurysm; MRC = Medical Research Council.

Brain Bank Neuropathologically Normal and MRC Controls

Libraries for brain biopsy controls were generated using TruSeq Exome Enrichment Kit (Illumina), according to the manufacturer's protocol, and sequenced on Illumina HiSeq, generating 150bp paired end reads. The average coverage obtained was 36x. Reads were aligned to

GRCh37 using Novoalign and variants were called using GATK HaplotypeCaller. Whole exome sequencing data from the MRC brain bank samples were obtained from EGA (EGAS00001001599), mainly sequenced using the Illumina HiSeq 2000 array (ega-archive.org/).

We compared the MAF of rare variants in our cases with our in-house brain control exomes, the MRC controls, and data available on population databases, specifically gnomAD (gnomad.broadinstitute.org) and Exome Aggregation Consortium (ExAC) (exac.broadinstitute.org) (figure 2).¹⁵ We then further checked the estimated effect of the according rare mutation using different computation predictive programs. As functional prediction scores, we used PolyPhen-2 and Sorting Intolerant From Tolerant (SIFT). These are used to help interpret the sequence variant.

Standard Protocol Approvals, Registrations, and Patient Consents

This study received approval from the ethical standards committee on human experimentation for any experiments using human subjects (UKCRN ID-7512, 09/H0716/54).

Data Availability

Anonymized data will be shared by request from any qualified investigator. EGAS00001001599 can be requested from the European Genome-Phenome Archive.

Table 1 Baseline Characteristics and Risk Factors

	Familial aneurysms, n = 275	Nonfamilial aneurysms, n = 2059	p Value	UIA, n = 90	aSAH, n = 185	p Value
Age, y	51.8 (12.6)	54.5 (12.7)	0.001	53.6 (10.9)	51 (13.3)	0.1
Female sex	217 (78.9)	1,423 (69.1)	0.001	75 (83.3)	142 (76.8)	0.21
Current smoker	108 (39.3)	885 (43)	0.24	26 (28.9)	82 (44.3)	0.02
Current drinker	176 (64)	1,384 (67.2)	0.29	52 (58.8)	124 (67)	0.14
PMH						
HTN	98 (35.6)	725 (35.2)	0.89	38 (42.2)	60 (32.4)	0.11
Hypercholesterolemia	60 (21.8)	508 (24.7)	0.3	33 (36.7)	27 (14.6)	<0.001
DM	6 (2.2)	102 (5)	0.05	2 (2.2)	4 (2.2)	0.97
OAC	22 (8)	96 (4.7)	0.02	4 (4.44)	18 (9.7)	0.14
Antiplatelets	33 (12)	271 (13.2)	0.59	19 (21.19)	14 (7.6)	0.002
Statins	48 (17.5)	417 (20.3)	0.28	26 (28.9)	22 (11.9)	0.001
Anti-HTN	81 (29.5)	599 (29.1)	0.9	32 (35.6)	49 (26.5)	0.12
Previous SAH	4 (1.5)	36 (1.8)	0.73	0	4 (2.2)	omit

Abbreviations: anti-HTN = antihypertensive medication; aSAH = aneurysmal subarachnoid hemorrhage; DM = diabetes mellitus; HTN = hypertension; OAC = oral anticoagulation; PMH = past medical history; SAH = subarachnoid hemorrhage; UIA = unruptured intracranial aneurysm. Values are mean (SD) or n (%).

Figure 2 Family Trees of Individuals With Rare Genetic Variants

Results

We included 275 patients with family history of IA. Of these 275 patients, 185 (67.3%) had had an aSAH. Baseline characteristics and risk factors are summarized in table 1. Comparing familial with sporadic IA, we found that individuals with familial IA were, as expected, younger than their non-familial equivalent. Interestingly, individuals with familial IA were more frequently female, and were more frequently on oral anticoagulation, but not on antiplatelets. We did not find a difference in hypertension. In our cohort of patients with family history of IA, 9/275 individuals (3.3%) harbored a rare (<0.01% gnomAD MAF) exonic mutation (figure 2 for the family trees of 7/9 individuals).

We found a total of 13 different nucleotide changes in *ANGPTL6*. Four of these changes resulted in the synonymous amino acid and 3 had a gnomAD MAF above 1% (2 of which were also present in the control group) and were consequently not considered. In the remaining 6 rare genetic variants, we noted 5 different missense mutations and 1 nonsense mutation (table 2). None of these rare genetic variants was present in our control population of 1,800 individuals. Figure 3 demonstrates imaging studies of the aneurysms of 3 of our patients.

Previously Reported Variants and Associated Variants

We replicated the previous finding of c.392A>T (p.Glu131Val), which has previously been reported as deleterious as per SIFT.¹³ In our cohort, this mutation was present in 2 individuals (MAF 0.004; figure 2, individuals 1 and 2). It

was not present in our control population or the ExAC database, but it was present in gnomAD population controls, with a MAF of 0.001. p.Glu131Val is moderately conserved across species, and lies within the linker region between the coiled-coil and fibrinogen-like domains (FLD), but no functional role has been described (figures 4 and 5). We found a previously unreported mutation very close to c.392A>T: c.332G>A (p.Arg111His) in 1 individual (MAF 0.002; figure 2, individual 8). This variant was not present in our control population or ExAC or gnomAD databases and is estimated to be deleterious as per SIFT. The mutation lies within the coiled-coil domain, which is important for the function of angiotensin-like proteins.¹⁶ p. Arg111His is moderately conserved across species but no functional role for this specific residue has been described.

Novel Rare Variants

In addition to c.332G>A (p.Arg111His), which lies in close proximity to the previously reported c.392A>T (p.Glu131Val), we found 4 additional rare genetic variants that have not been reported. Most notably, we found a c.851C>G (p.Ser284*) nonsense mutation leading to a premature stop codon in 1 individual (MAF 0.002; figure 2, individual 3). This variant was not present in our own control cohort; however, it was present in ExAC and gnomAD, with MAF of 0.0004 and 0.0005, respectively. This mutation occurs in Ex4 and is predicted to lead to nonsense-mediated decay (NMD) or production of a truncated protein lacking the majority of the FLD. This stop-gain mutation has been described to result in complete loss of secretion of *ANGPTL6* protein in cells.¹⁷ It is moderately conserved across species.

Table 2 Rare Nonsynonymous Variants in the *ANGPTL6* Gene in Patients With Familial Intracranial Aneurysm

Nucleotide change	Exon	Protein consequence	dbSNP	Het	Hom	MAF (n = 500)	ExAC MAF	gnomAD MAF	SIFT	PolyPhen
c.243G>C	Ex2	p.81R>S	rs1166038416	1	0	0.0018	NP	0.00011	Tolerated (0.74)	Benign (0.025)
c.245T>A	Ex2	p.82L>Q	rs979714863	3	0	0.0055	NP	0.000068	Tolerated (0.25)	Benign (0.003)
c.332G>A	Ex2	p.111R>H	rs1310828976	1	0	0.0018	NP	NP	Deleterious (0.04)	Benign (0.254)
c.392A>T ¹³	Ex2	p.131E>V	rs576667683	2	0	0.0036	NP	0.0015	Deleterious (0)	possibly damaging (0.724)
c.851C>G	Ex4	p.284S>	rs201622589	1	0	0.0018	0.0004	0.00045		
c.1073G>A	Ex5	p.358R>H	rs779399477	1	0	0.0018	0.000025	0.000031	Tolerated (0.33)	Benign (0.013)

Abbreviations: Ex = Exon; ExAC = Exome Aggregation Consortium; gnomAD = Genome Aggregation Database; Het = Heterozygous; Hom = Homozygous; MAF = minor allele frequency; NP = not present; SIFT = Sorting Intolerant From Tolerant.

Another mutation present in our cases lying within the FLD of the *ANGPTL6* gene is c.1073G>A (p.Arg358His), which was again present in 1 individual (MAF 0.002; figure 2, individual 4). This variant was not present in our control group but present in ExAC and gnomAD databases with low MAF of 0.00002 and 0.00003, respectively. p.Arg358His lies within the FLD, which is also important for the activity of angiopoietin-like proteins and is moderately conserved across species, but no functional role of this specific residue has been described.¹⁶

Finally, we found 2 further missense mutations: c.243G>C (p.Arg81Ser) in 1 individual (MAF 0.002; figure 2, individual 5) and c.245T>A (p.Gln82Leu) in 3 individuals (MAF 0.005; figure 2, individuals 6 and 7; family tree for the third individual not available). Interestingly, 2 of these individuals were related (mother and daughter). These 2 genetic variants affect adjacent residues, and both lie within the coiled-coil domain of *ANGPTL6*. Both amino acids are moderately conserved across species but for neither of them has a functional role been described.

Discussion

We present a study of the *ANGPTL6* gene in a large cohort with family history of IA replicating 1 of the previously presented variants: c.392A>T (p.Glu131Val) in the linker region. In addition, we found c.332G>A (p.Arg111His) located very close to this previously described mutation within the coiled-coiled domain. We also found 2 genetic variants in the FLD of the *ANGPTL6* gene, 1 being a nonsense mutation that leads to a stop-gain, which has been described to result in complete loss of *ANGPTL6* secretion. None of the genetic variants presented here was found in our 1,800 controls.

Most likely due to the continuous expansion and sometimes merging of databases such as gnomAD and ExAC, some of the previously described variants have now also been described in these control populations. We did not replicate any of the other described variants but found 4 additional ones. All of the rare variants reported here were much rarer in the analyzed control populations, if present at all.

Our study provides additional insight building on the previous study on *ANGPTL6* in individuals with familial IA.¹³ The c.851C>G (p.Ser284*) stop-gain mutation we identified lies in *ANGPTL6* Ex4 and likely results in NMD of the mutant transcript, which could contribute to haploinsufficiency. It is also possible that this mutation leads to the expression of a truncated protein lacking the majority of the FLD, reducing *ANGPTL6* protein function or stability. In fact, a previous study reported this variant leads to a complete loss of secretion of the *ANGPTL6* protein in cells.¹⁷ The actual effect of reduced *ANGPTL6* secretion remains unknown and warrants further investigation. We also identify novel genetic variants in the FLD and coiled-coil domain of *ANGPTL6*, which might be impairing the proangiogenic function or overall stability of the protein.¹⁶ Although no function specific to *ANGPTL6* has been established, the FLD is suggested to regulate angiogenic activities.^{10,16} The same is true for the coiled-coil domain. Several functions of this domain have been reported, including activity in polymer formation, molecular recognition, cytoskeletal regulation, and pH sensing.¹⁸ Based on the information available, a dominant negative mechanism cannot be conclusively ruled out, especially in missense changes. A potential mechanism could be that the mutation alters the protein structural binding, preventing *ANGPTL6* signaling. The *ANGPTL* family also regulates lipid and glucose metabolism and by doing so indirectly

Figure 3 Familial Intracranial Aneurysm (IA) With *ANGPTL6* Genetic Variants Had Typical History and Imaging of IA

Patient 1: a 48-year-old woman presenting with subarachnoid hemorrhage (SAH) within the right sylvian fissure and suprasellar cistern on the axial noncontrast CT examination (A), from a large right terminal M1 segment right middle cerebral artery aneurysm demonstrated on a maximal intensity projection axial CT angiogram (B), and an oblique view of a selective injection into the right internal carotid artery on digital subtraction cerebral angiography (C). Patient 2: a 41-year-old man presenting with SAH within the right sylvian fissure, anterior interhemispheric fissure, and right quadrigeminal cistern on the axial noncontrast CT examination (D), from an A1-A2 junction aneurysm of the left anterior cerebral artery demonstrated on axial CT angiography (E), and a Towne projection from the selective injection of the left internal carotid artery on digital subtraction cerebral angiography (F). Patient 3: a 57-year-old woman presenting with extensive SAH within the sylvian fissures, anterior interhemispheric fissure, as well as the perimesencephalic and ambient cisterns on the axial noncontrast CT examination (G), and from a posterior communicating segment aneurysm of the left internal carotid artery demonstrated on the lateral projection from the selective injection of the left internal carotid artery on digital subtraction cerebral angiography pre-(H) and post-(I) coil embolization with satisfactory exclusion from the arterial circulation.

influences risk factors for IA formation such as hypertension.^{19–22}

As Bourcier et al.¹³ hypothesized, individuals with rare variants in the *ANGPTL6* gene might require additional factors in order to trigger the development of IA. Therefore, heterozygous *ANGPTL6* genetic variants might only exhibit a deleterious effect in combination with certain risk factors such as high blood pressure. Other genes have also been previously reported to play a role in familial IA, such as *ADAMST15* and *PCNT*. *ADAMST15* has been shown to be significantly aggregated in families with IA.²³ It might be associated with IA formation through abnormal transcription of

metalloproteinases.²⁴ Two rare variants of the *PCNT* gene have been found in familial IA cases.²⁵ Deletions and mutations of this gene cause a type of dwarfism associated with IA in up to 20%.²⁵ This indicates the genetic complexity of this disease.

Our study has strengths: we present a large cohort of individuals with family history of IA sequenced using Sanger sequencing. Sanger sequencing remains the gold standard for mutation confirmation. Concordance between Sanger sequencing and whole-exome sequencing has been reported to be 97.3%.²⁶ This allowed us to check the variants found in our cases in a control population sequenced by whole-exome

Figure 4 Multiple Alignment Showing Moderate Conservation Across Species and ANGPTL6 Homolog of the Residues Affected by the Variants Identified in our Cohort (Variants Highlighted in Yellow)

sequencing. None of the rare genetic variants found in our cases were present in our own control dataset; if the genetic variants were present in the gnomAD and ExAC control populations, they were significantly rarer, reducing the likelihood of winner's curse.

Our study has limitations: we did not sequence affected or unaffected family members (except for 2 individuals who were related) as DNA for family members was not available and our aim was to externally validate the previous results of 4 rare coding variants within the *ANGPTL6* gene; information and DNA for relatives were generally not collected as part of the GOSH study. This would be especially of interest as 1 variant was present in both mother and daughter, indicating a potential familial clustering. We included patients with only 1 affected second-degree relative not fulfilling the classical criteria of familial IA. We do not believe that this limits our results as none of the patients harboring a rare variant only had 1 affected second-degree relative. In addition, we were not able to obtain the family

trees in 2 individuals; 1 was deceased (c.245T>A; p.Gln82Leu) and the other was not contactable (c.332G>A; p.Arg111His). We therefore cannot make any statements about penetrance in our population. Although we cannot make any statements about penetrance and we refrained from burden analysis due to cases and controls not having been sequenced with the same method, we demonstrated that the rare genetic variants found in our cases are far less frequent or absent in the considered control cohorts. Therefore, a genetic contribution to this disease, including by rare variant in the *ANGPTL6* gene, is highly likely.

We replicated some of the previously reported genetic variants in the *ANGPTL6* gene in a large cohort of individuals with family history of IA. Novel found genetic variants support that *ANGPTL6* loss of function might be a genetic risk factor influencing IA development. These findings remain to be confirmed in an independent cohort. The function of the *ANGPTL6* gene needs to be evaluated. It might offer a target

Figure 5 Homology Model of Human ANGPTL6 Protein and Respective Location of the Rare Variants Identified in This Study

	R81, L82	R111	E131
ANGPTL6 HOMO SAPIENS (HUMAN)	LRMRVGRHEELLRELO ^R LAADGAVAGEVRLRKESRGLSARLGQL ^R LAQLQHEAGPGAGPGADLG----		-----A ^E PAAALALSLEPESDH
ANGPTL6 PAN TROGLODYTES (CHIMPANZEE)	LRMRVGRHEELLRELO ^R LAADGAVAGEVRLRKESRGLSARLGQL ^R LAQLQHEAGPGAGPGADLGAEPAA		-----A ^E PAAALALSLEPESDH
ANGPT16 MUS MUSCULUS (MOUSE)	LRMRLGRHEELLRAL ^R RAAEGGALADEVRLREHSLTLNTRLGQL ^R LAQLQQEAPAEF----		-----DLG----A ^E PAAALGLSLEPESDH
ANGPT16 RATTUS NORVEGICUS (RAT)	LRMRVGRHEELLRAL ^R RAAEGGALADEVRLREHSLTLNTRLGQL ^R LAQLQQEAGAPF----		-----DLG----A ^E PAAALGLSLEPESDH
ANGPTL6 BOS TAURAS (COW)	LRRVSRHEELLREL ^R LAADGAVAGEVRLRKESRGLSARLGQL ^R LAQLQHEAGPGAAPS----		-----PG----A ^E PASALALSLEPESDH
ANGPTL6 CANIS LUPUS FAMILIARIS (DOG)	LRMRVGRHEELLRELO ^R LAADGAVAGEVRLRKESRGLSARLGQL ^R LAQLQHEAGPGAGPG----		-----PG----A ^E PAALLALSLEPESDY
ANGPT16 DANIO RERIO (ZEBRAFISH)	LRALLERQQDQLDM ^{RGR} LEEEGTLASEMRILRRSVNMNARIQL ^Y LAQLLSEIIQKKEQV----		-----A ^E EEQRRRVEPESDW
	S284		R358
ANGPTL6 HOMO SAPIENS (HUMAN)	GRHVV ^S WCEQQLEGGGWTVIQRQDGSVNFFTTWQHYKAGFGRPDGEYWLGLPEVYQLTSRGDHELLVLLDWDGGRGAR		RAHYDGF
ANGPTL6 PAN TROGLODYTES (CHIMPANZEE)	GRHVV ^S WCEQQLEGGGWTVIQRQDGSVNFFTTWQHYKAGFGRPDGEYWLGLPEVYQLTSRGDHELLVLLDWDGGRGAR		RAHYDGF
ANGPT16 MUS MUSCULUS (MOUSE)	GRRVV ^Y WCEQQLEGGGWTVIQRQDGSVNFFTTWQHYKAGFGRPEGEYWLGLPEVHQVTSRGDHELLLILLEDWGGRAR		RAHYDSF
ANGPT16 RATTUS NORVEGICUS (RAT)	GRRVV ^Y WCEQQLEGGGWTVIQRQDGSVNFFTTWQHYKVGFRPDGEYWLGLPEVHQVTSRGDHELLLILLEDWGGRAR		RAHYDSF
ANGPTL6 BOS TAURAS (COW)	-----VGFQPDGEYWLGLPEVHQVTSRGDHELLVLLDWDGGRGAR		RAHYDGF
ANGPTL6 CANIS LUPUS FAMILIARIS (DOG)	GRHVV ^S WCEQQLEGGGWTVIQRQDGSVNFFTTWQHYKVGFGQPDGEYWLGLPEVHQLTSRGDHELLVLLDWDGGRGAR		RAHYDGF
ANGPT16 DANIO RERIO (ZEBRAFISH)	TNRLLO ^Q AWCDQSRAGGWTVIQRQDGSVNFFRTWDQYKQGFNLDEGYWLGLLEHLWYLSQATYKLRVAMEDWDQGRQV		YAEYDSF

ANGPTL6 homology model created using Phyre2²⁷ and human fibrinogen beta chain (PDB ID: 3GHG) as the structural template.²⁸ Three recognizable domains are present: signal peptide (blue), coiled-coil domain (green), and fibrinogen-like domain (yellow).

for screening familial IA cases for *ANGPTL6* as well as other genes involved in IA, but also highlights the *ANGPTL6* pathway as a potential therapeutic target.

Acknowledgment

The authors thank the NIHR Clinical Research Network Practitioners, the investigators at participating centers, and the patients.

Study Funding

The GOSH study is funded by the Stroke Association. This work was partly undertaken at UCLH/UCL, which received a proportion of funding from the Department of Health's NIHR Biomedical Research Centres funding scheme. These funding bodies had no direct involvement in this study.

Disclosure

Isabel C. Hostettler and Henry Houlden received funding from the Alzheimer Research UK and Dunhill Medical Trust Foundation. Duncan Wilson and David J. Werring received funding from the Stroke Foundation/British Heart Foundation. Benjamin O'Callaghan, Enrico Bugiardi, Emer O'Connor, Jana Vandrovcova, Indran Davagnanam, Varinder Alg, Stephen Bonner, Daniel Walsh, Diederik Bulters, Neil Kitchen, Martin M. Brown, and Joan Grieve report no disclosures. Go to Neurology.org/N for full disclosures.

Publication History

Received by *Neurology* January 25, 2020. Accepted in final form October 8, 2020.

Appendix 1 Authors

Name	Location	Contribution
Isabel C. Hostettler, MD	Stroke Research Centre, University College London, Institute of Neurology, UK	Designed and conceptualized study, acquisition of data, analyzed the data, drafted the manuscript, revised the manuscript
Benjamin O'Callaghan, MSc	Neurogenetics Laboratory, The National Hospital of Neurology and Neurosurgery, London, UK	Contributed to the laboratory work, analyzed the data, drafted the manuscript, revised the manuscript
Enrico Bugiardi, MD	Neurogenetics Laboratory, The National Hospital of Neurology and Neurosurgery, London, UK	Contributed to the laboratory work, analyzed the data, drafted the manuscript, revised the manuscript
Emer O'Connor, MD	Neurogenetics Laboratory, The National Hospital of Neurology and Neurosurgery, London, UK	Contributed to the laboratory work, analyzed the data, drafted the manuscript, revised the manuscript
Jana Vandrovcova, PhD	Department of Neuromuscular Disorders, UCL Institute of Neurology, Queen Square, London, UK	Analyzed the data, interpreted the data, revised the manuscript for intellectual content

Appendix 1 (continued)

Name	Location	Contribution
Indran Davagnanam	Department of Neuroradiology, The National Hospital for Neurology and Neurosurgery, University College London Hospitals NHS Foundation Trust, Queen Square, UK	Analyzed the data, revised the manuscript for intellectual content
Varinder Alg, MD	Stroke Research Centre, University College London, Institute of Neurology, UK	Acquisition of data, revised the manuscript for intellectual content
Stephen Bonner, PhD	Department of Anaesthesia, The James Cook University Hospital, Middlesbrough, UK	Acquisition of data, revised the manuscript for intellectual content
Daniel Walsh, PhD	Department of Neurosurgery, King's College Hospital NHS Foundation Trust, London, UK	Acquisition of data, revised the manuscript for intellectual content
Diederik Bulters, FRCS	Department of Neurosurgery, University Hospital Southampton NHS Foundation Trust, UK	Acquisition of data, revised the manuscript for intellectual content
Neil Kitchen, PhD	Department of Neurosurgery, The National Hospital of Neurology and Neurosurgery, London, UK	Acquisition of data, revised the manuscript for intellectual content
Martin M. Brown, FRCP	Stroke Research Centre, University College London, Institute of Neurology, UK	Acquisition of data, revised the manuscript for intellectual content
Joan Grieve, MD	Department of Neurosurgery, The National Hospital of Neurology and Neurosurgery, London, UK	Acquisition of data, revised the manuscript for intellectual content
David J. Werring, PhD, FRCP	Stroke Research Centre, University College London, Institute of Neurology, UK	Designed and conceptualized study, interpreted the data, revised the manuscript for intellectual content
Henry Houlden, PhD, FRCP	Neurogenetics Laboratory, The National Hospital of Neurology and Neurosurgery, London, UK	Designed and conceptualized study, interpreted the data, revised the manuscript for intellectual content

Appendix 2 Coinvestigators

Name	Location	Contribution
Gareth Roberts	Royal Preston Hospital	Recruited patients
Timothy Jones	St George's Hospital, London	Recruited patients

Appendix 2 (continued)

Name	Location	Contribution
Giles Critchley	Hurstwood Park Neurologic Centre	Recruited patients
Pankaj Sharma	Imperial Healthcare, Charing Cross Hospital, London	Recruited patients
Richard Nelson	Frenchay Hospital, Bristol	Recruited patients
Peter Whitfield	Derriford Hospital, Plymouth	Recruited patients
Stuart Ross	Leeds General Infirmary	Recruited patients
Hiren Patel	Salford Royal Hospital	Recruited patients
Paul Eldridge	The Walton Centre, Liverpool	Recruited patients
Kari Saastamoinen	The Royal London Hospital	Recruited patients
Umang Patel	Sheffield Teaching Hospitals NHS Trust	Recruited patients
Enas Lawrance	Mayday Hospital, Croydon	Recruited patients
Subha Vandabona	Mount Gould Hospital, Plymouth	Recruited patients
David Mendelow	Newcastle University Hospitals NHS Trust	Recruited patients
Rachel Teal	Oxford Radcliffe Infirmary	Recruited patients

References

1. Vlak MH, Algra A, Brandenburg R, Rinkel GJ. Prevalence of unruptured intracranial aneurysms, with emphasis on sex, age, comorbidity, country, and time period: a systematic review and meta-analysis. *Lancet Neurol* 2011;10:626–636.
2. Johnston SC, Selvin S, Gress DR. The burden, trends, and demographics of mortality from subarachnoid hemorrhage. *Neurology* 1998;50:1413–1418.
3. Suarez JJ, Tarr RW, Selman WR. Aneurysmal subarachnoid hemorrhage. *New Engl J Med* 2006;354:387–396.
4. Kim ST, Brinjikji W, Lanzino G, Kallmes DF. Neurovascular manifestations of connective-tissue diseases: a review. *Interv Neuroradiol* 2016;22:624–637.
5. Flahault A, Trystram D, Nataf F, et al. Screening for intracranial aneurysms in autosomal dominant polycystic kidney disease is cost-effective. *Kidney Int* 2018;93:716–726.
6. Alg VS, Sofat R, Houlden H, Werring DJ. Genetic risk factors for intracranial aneurysms: a meta-analysis in more than 116,000 individuals. *Neurology* 2013;80:2154–2165.
7. Bourcier R, Redon R, Desal H. Genetic investigations on intracranial aneurysm: update and perspectives. *J Neuroradiol* 2015;42:67–71.
8. Huttunen T, von und zu Fraunberg M, Frosen J, et al. Saccular intracranial aneurysm disease: distribution of site, size, and age suggests different etiologies for aneurysm formation and rupture in 316 familial and 1454 sporadic eastern Finnish patients. *Neurosurgery* 2010;66:631–638; discussion 638.
9. Oike Y, Ito Y, Maekawa H, et al. Angiopoietin-related growth factor (AGF) promotes angiogenesis. *Blood* 2004;103:3760–3765.
10. Camenisch G, Pisabarro MT, Sherman D, et al. ANGPTL3 stimulates endothelial cell adhesion and migration via integrin alpha v beta 3 and induces blood vessel formation in vivo. *J Biol Chem* 2002;277:17281–17290.
11. Le Jan S, Amy C, Cazes A, et al. Angiopoietin-like 4 is a proangiogenic factor produced during ischemia and in conventional renal cell carcinoma. *Am J Pathol* 2003;162:1521–1528.
12. Santulli G. Angiopoietin-like proteins: a comprehensive look. *Front Endocrinol* 2014;5:4.
13. Bourcier R, Le Scouarnec S, Bonnaud S, et al. Rare coding variants in ANGPTL6 are associated with familial forms of intracranial aneurysm. *Am J Hum Genet* 2018;102:133–141.
14. Hostettler IC, Alg VS, Shahi N, et al. Characteristics of unruptured compared to ruptured intracranial aneurysms: a multicenter case-control study. *Neurosurgery* 2017.
15. Lek M, Karczewski KJ, Minikel EV, et al. Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 2016;536:285–291.
16. Hato T, Tabata M, Oike Y. The role of angiopoietin-like proteins in angiogenesis and metabolism. *Trends Cardiovasc Med* 2008;18:6–14.
17. Hendricks AE, Bochukova EG, Marenne G, et al. Rare variant analysis of human and rodent obesity genes in individuals with severe childhood obesity. *Sci Rep* 2017;7:4394.
18. Burkhard P, Stetefeld J, Strelkov SV. Coiled coils: a highly versatile protein folding motif. *Trends Cell Biol* 2001;11:82–88.
19. Ono M, Shimizugawa T, Shimamura M, et al. Protein region important for regulation of lipid metabolism in angiopoietin-like 3 (ANGPTL3): ANGPTL3 is cleaved and activated in vivo. *J Biol Chem* 2003;278:41804–41809.
20. Shimamura M, Matsuda M, Yasuno H, et al. Angiopoietin-like protein3 regulates plasma HDL cholesterol through suppression of endothelial lipase. *Arteriosclerosis Thromb Vasc Biol* 2007;27:366–372.
21. Wang Y, Lam KS, Lam JB, et al. Overexpression of angiopoietin-like protein 4 alters mitochondria activities and modulates methionine metabolic cycle in the liver tissues of db/db diabetic mice. *Mol Endocrinol* 2007;21:972–986.
22. Oike Y, Akao M, Yasunaga K, et al. Angiopoietin-related growth factor antagonizes obesity and insulin resistance. *Nat Med* 2005;11:400–408.
23. Yan J, Hitomi T, Takenaka K, et al. Genetic study of intracranial aneurysms. *Stroke* 2015;46:620–626.
24. Zhang J, Clatterbuck RE. Molecular genetics of human intracranial aneurysms. *Int J Stroke* 2008;3:272–287.
25. PCNT point mutations and familial intracranial aneurysms. *Neurology* 2019;93:231.
26. Hamilton A, Tetreault M, Dymont DA, et al. Concordance between whole-exome sequencing and clinical Sanger sequencing: implications for patient care. *Mol Genet Genomic Med* 2016;4:504–512.
27. Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. The Phyre2 web portal for protein modeling, prediction and analysis. *Nat Protoc* 2015;10:845–858.
28. Kollman JM, Pandi L, Sawaya MR, Riley M, Doolittle RF. Crystal structure of human fibrinogen. *Biochemistry* 2009;48:3877–3886.

Neurology®

ANGPTL6 Genetic Variants Are an Underlying Cause of Familial Intracranial Aneurysms

Isabel C. Hostettler, Benjamin O'Callaghan, Enrico Bugiardini, et al.
Neurology 2021;96:e947-e955 Published Online before print October 26, 2020
DOI 10.1212/WNL.0000000000011125

This information is current as of October 26, 2020

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/96/6/e947.full
References	This article cites 27 articles, 8 of which you can access for free at: http://n.neurology.org/content/96/6/e947.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Genetics http://n.neurology.org/cgi/collection/all_genetics Stroke prevention http://n.neurology.org/cgi/collection/stroke_prevention Subarachnoid hemorrhage http://n.neurology.org/cgi/collection/subarachnoid_hemorrhage
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.