Teaching NeurolImage: Dynamic MRI in the Evaluation of Cervical Myelopathy

Miguel Quintas-Neves, MD, and Ângelo Carneiro, MD, MSc

Neurology® 2021;97:e1359-e1360. doi:10.1212/WNL.00000000000012181

Correspondence

Dr. Quintas-Neves
mlqneves@gmail.com

Figure 1 Dynamic MRI of the Cervical Spine, Sagittal View

Sagittal T2-weighted imaging at rest (A), flexion (B), and extension (C). Spinal cord hyperintensity at the C1 level is depicted, suggesting gliosis/edema. There is also increased atlanto-dental interval (white line in A) and periarthral fibrosis. During flexion (B), severe canal stenosis is unveiled, favoring atlanto-axial instability.

A 44-year-old woman, with a childhood history of motor vehicle accident, was referred to neurosurgery due to progressive tetraparesis, inability to walk, generalized hyperreflexia, and bilateral extensor plantar reflexes (American Spinal Injury Association Impairment Scale D/Japanese Orthopaedic Association score 7), without bowel/bladder dysfunction. Dynamic MRI (Figures 1 and 2) showed spinal cord edema/gliosis at the C1 level and atlanto-axial subluxation. During flexion, severe spinal canal stenosis was observed, confirming atlanto-axial instability. Posterior arthrodesis was performed and resulted in resolution of instability with no neurologic deterioration after 3 months. Dynamic MRI in the supine position can be an accurate technique to unveil occult cervical canal stenosis.^{1,2} Although most cases are revealed or aggravated during extension, flexion should also be tested.²

Study Funding

The authors report no targeted funding.

Disclosure

The authors report no disclosures relevant to the manuscript. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

MORE ONLINE

Teaching slides

links.lww.com/WNL/B414

From the Department of Neuroradiology, Hospital de Braga, Portugal.

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Copyright © 2021 American Academy of Neurology

Copyright © 2021 American Academy of Neurology. Unauthorized reproduction of this article is prohibited.

Figure 2 MRI of the Cervical Spine, Axial View

Axial T2-weighted imaging at rest at the level of C1 confirms the bilateral spinal cord hyperintense lesion, in keeping with gliosis/edema.

Appendix Authors

Name	Location	Contribution
Miguel Quintas-Neves, MD	Department of Neuroradiology, Hospital de Braga, Portugal	Acquisition of data, clinical and imaging data review, literature review, final manuscript writing
Ângelo Carneiro, MD, MSc	Department of Neuroradiology, Hospital de Braga, Portugal	Clinical and imaging data review, literature review, final manuscript writing

References

1. Zeitoun D, El Hajj F, Sariali E, Catonné Y, Pascal-Mousellard H. Evaluation of spinal cord compression and hyperintense intramedullary lesions on T2-weighted sequences in patients with cervical spondylotic myelopathy using flexion-extension MRI protocol. *Spine J*. 2015;15(4):668-674.
2. Nanfang X, Shaobo W, Huishu Y, Xiaoguang L, Zhongjun L. Does dynamic supine magnetic resonance imaging improve the diagnostic accuracy of cervical spondylotic myelopathy? A review of the current evidence. *World Neurosurg*. 2017;100:474-479.

Neurology®

Teaching NeuroImage: Dynamic MRI in the Evaluation of Cervical Myelopathy

Miguel Quintas-Neves and Ângelo Carneiro

Neurology 2021;97:e1359-e1360 Published Online before print May 24, 2021

DOI 10.1212/WNL.00000000000012181

This information is current as of May 24, 2021

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/97/13/e1359.full
References	This article cites 2 articles, 0 of which you can access for free at: http://n.neurology.org/content/97/13/e1359.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): MRI http://n.neurology.org/cgi/collection/mri Spinal cord trauma http://n.neurology.org/cgi/collection/spinal_cord_trauma
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2021 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.