Rare Missense Functional Variants at *COL4A1* and *COL4A2* in Sporadic Intracerebral Hemorrhage

Jaeyoon Chung, PhD, Graham Hamilton, PhD, Minsup Kim, PhD, Sandro Marini, MD, Bailey Montgomery, Jonathan Henry, Art E. Cho, PhD, Devin L. Brown, MD, Bradford B. Worrall, MD, James F. Meschia, MD, Scott L. Silliman, MD, Magdy Selim, MD, PhD, David L. Tirschwell, MD, MSc, Chelsea S. Kidwell, MD, Brett Kissela, MD, MS, Steven M. Greenberg, MD, PhD, Anand Viswanathan, MD, PhD, Joshua N. Goldstein, MD, PhD, Carl D. Langefeld, PhD, Kristiina Rannikmae, MD, Catherine L.M. Sudlow, BMBCh, MSc, DPhil, FRCPE, Neshika Samarasekera, PhD, MRCP, Mark Rodrigues, PhD, FRCR, Rustam Al-Shahi Salman, PhD, FRCP Edin, James G.D. Prendergast, PhD, Sarah E. Harris, PhD, Ian Deary, MBChB, PhD, FRCP, Daniel Woo, MD, MSc, Jonathan Rosand, MD, MSc, Tom Van Agtmael, PhD,* and Christopher D. Anderson, MD, MMSc*

Correspondence

Dr. Anderson
cdanderson@
mgh.harvard.edu
or Dr. Van Agtmael
Tom.vanagtmael@
glasgow.ac.uk

Neurology® 2021;97:e236-e247. doi:10.1212/WNL.0000000000001227

Abstract

Objective

To test the genetic contribution of rare missense variants in *COL4A1* and *COL4A2* in which common variants are genetically associated with sporadic intracerebral hemorrhage (ICH), we performed rare variant analysis in multiple sequencing data for the risk for sporadic ICH.

Methods

We performed sequencing across 559 Kbp at 13q34 including *COL4A1* and *COL4A2* among 2,133 individuals (1,055 ICH cases; 1,078 controls) in United States–based and 1,381 individuals (192 ICH cases; 1,189 controls) from Scotland-based cohorts, followed by sequence annotation, functional impact prediction, genetic association testing, and in silico thermodynamic modeling.

Results

We identified 107 rare nonsynonymous variants in sporadic ICH, of which 2 missense variants, rs138269346 (*COL4A1*^{T110T}) and rs201716258 (*COL4A2*^{L203L}), were predicted to be highly functional and occurred in multiple ICH cases but not in controls from the United States–based cohort. The minor allele of rs201716258 was also present in Scottish patients with ICH, and rs138269346 was observed in 2 ICH-free controls with a history of hypertension and myocardial infarction. Rs138269346 was nominally associated with nonlobar ICH risk ($p = 0.05$), but not with lobar ICH ($p = 0.08$), while associations between rs201716258 and ICH subtypes were nonsignificant ($p > 0.12$). Both variants were considered pathogenic based on minor allele frequency (<0.00035 in European populations), predicted functional impact (deleterious or probably damaging), and in silico modeling studies (substantially altered physical length and thermal stability of collagen).

*These authors contributed equally to this work.

From the Center for Genomic Medicine (J.C., S.M., B.M., J.H., J.R., C.D.A.), Department of Neurology (B.M., J.H., S.M.G., A.V., J.R., C.D.A.), McCance Center for Brain Health (J.H., J.R., C.D.A.), and Department of Emergency Medicine (J.N.G.), Massachusetts General Hospital, Boston; Program in Medical and Population Genetics (J.C., J.R., C.D.A.), Broad Institute, Boston, MA; Glasgow Polyomics, Wolfson Wohl Cancer Research Centre, Garscube Campus (G.H.), and Institute of Cardiovascular and Medical Sciences, College of Medical, Veterinary and Life Sciences (G.H., T.V.A.), University of Glasgow, Bearsden, UK; Department of Bioinformatics (M.K., A.E.C.), Korea University, Sejong, South Korea; Stroke Program, Department of Neurology (D.L.B.), University of Michigan, Ann Arbor; Department of Neurology and Public Health Sciences (B.B.W.), University of Virginia Health System, Charlottesville; Department of Neurology (J.F.M.), Mayo Clinic Jacksonville; Department of Neurology (S.L.S.), University of Florida College of Medicine, Jacksonville; Department of Neurology, Stroke Division (M.S.), Beth Israel Deaconess Medical Center, Boston, MA; Department of Neurology, Harborview Medical Center (D.L.T.), University of Washington, Seattle; Department of Neurology (C.S.K.), The University of Arizona, Tucson; Department of Neurology and Rehabilitation Medicine (B.K., D.W.), University of Cincinnati, OH; Center for Public Health Genomics and Department of Biostatistical Sciences (C.D.L.), Wake Forest School of Medicine, Winston-Salem, NC; Centre for Medical Informatics, Usher Institute (K.R., C.L.M.S.), Centre for Clinical Brain Sciences (N.S., M.R., R.A.-S.S.), The Roslin Institute (J.G.D.P.), and Lothian Birth Cohorts Group, Department of Psychology (S.E.H., I.D.), University of Edinburgh; and British Heart Foundation Data Science Centre (K.R.), London, UK. Dr. Anderson is currently at the Department of Neurology, Brigham and Women's Hospital, Boston, MA

Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures. Funding information and disclosures deemed relevant by the authors, if any, are provided at the end of the article.

The Article Processing Charge was funded by UKRI, University of Glasgow.

This is an open access article distributed under the terms of the Creative Commons Attribution License 4.0 (CC BY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Glossary

CADD = combined annotation-dependent deletion; **CSVD** = cerebral small vessel disease; **ERICH** = Ethnic/Racial Variation in Intracerebral Hemorrhage; **GATK** = Genome Analysis Toolkit; **GOCHA** = Gene Discovery for Warfarin-Related Intracerebral Hemorrhage; **GWAS** = genome-wide association studies; **ICD-10** = International Classification of Diseases–10; **ICH** = intracerebral hemorrhage; **LBC** = Lothian Birth Cohorts; **LD** = linkage disequilibrium; **LINCHPIN** = Lothian study of Intracerebral Haemorrhage Pathology, Imaging and Neurologic Outcome; **MAF** = minor allele frequency; **MD** = molecular dynamics; **ns** = nanoseconds; **PDB** = Protein Data Bank; **QC** = quality control; **SNP** = single nucleotide polymorphism; **SVS** = small vessel ischemic stroke; **UKB** = UK Biobank; **VEP** = variant effect predictor; **WES** = whole exome sequencing; **WGS** = whole genome sequencing.

Conclusions

We identified rare missense variants in *COL4A1/A2* in association with sporadic ICH. Our annotation and simulation studies suggest that these variants are highly functional and may represent targets for translational follow-up.

Intracerebral hemorrhage (ICH) accounts for 10%–15% of strokes but is the most fatal and least treatable stroke type.^{1–3} More than half of patients with ICH die within the first year after the disease, and most survivors have prolonged disability.³ The molecular pathways underlying ICH remain poorly understood, limiting therapeutic development.

Genome-wide association studies (GWAS) have identified genetic loci associated with ICH risk and outcome.^{4,5} We recently determined a genome-wide significant association for cerebral small vessel disease (CSVD) including nonlobar ICH and small vessel ischemic stroke (SVS) with *COL4A1* and *COL4A2* (collagen IV α chain 1 and 2) at 13q34,6 validating previous studies.⁷

Rare mutations that most frequently affect Gly residues in the Gly-Xaa-Yaa repeat in *COL4A1/A2* cause mendelian early-onset cerebrovascular disease, ocular dysgenesis, and myopathy.^{8–12} However, the full spectrum of effects of *COL4A1/A2* mutations remain incompletely characterized.^{7,13,14} Sequencing of a few families or small number of patients ($n < 100$) suggested rare variants in *COL4A1/A2* can contribute to sporadic ICH.^{9–15}

To identify rare coding variants in *COL4A1/A2* that may underlie the aforementioned GWAS association with ICH and SVS, we performed targeted sequencing of the 13q34 region in United States–based studies (1,055 ICH cases and 1,078 ICH-free controls) as well as exonic sequencing of this locus in 192 ICH cases and whole genome sequencing (WGS) of 1,189 ICH-free controls from Scotland. We present annotation results, prediction of deleteriousness, and case/control segregation of rare variants across 13q34 from these datasets.

Methods

Standard Protocol Approvals, Registrations, and Participant Consents

The study protocols were approved for enrollment of the United States–based cohorts (Gene Discovery for Warfarin-

Related Intracerebral Hemorrhage [GOCHA] and Ethnic/Racial Variation in Intracerebral Hemorrhage [ERICH]) and the Scotland-based cohorts (the Lothian study of Intracerebral Haemorrhage Pathology, Imaging and Neurologic Outcome [LINCHPIN] and Lothian Birth Cohorts [LBC]) by the institutional review boards of Massachusetts General Hospital and University of Edinburgh. Informed consent was obtained from participants or an appropriate legal surrogate according to all recruiting sites.

Study Participants

United States–Based Participants

We collected DNA and phenotyping data on 1,055 patients with ICH (534 lobar and 521 nonlobar ICH) and 1,078 ICH-free controls from GOCHA⁵ and ERICH.¹⁶ There is no overlap between the participants in the present study and those included in previous targeted sequencing studies of *COL4A1/A2*.^{13,17} A total of 145 participants (18 ICH and 176 controls) in this study appeared in the previous ICH GWAS.⁵

Scotland–Based Participants

We collected DNA and phenotyping data on 192 ICH cases (40 lobar and 152 nonlobar) from LINCHPIN among the sample registered in the Edinburgh Lothian Audit of the Treatment of Cerebral Hemorrhage study.¹⁸

As a control population for the LINCHPIN cohort, we selected clinical and sequencing data from 1,189 individuals from the LBC study as an ethnically and geographically matched stroke-free control population to cases. Self-reported ethnicity of the patients with ICH and controls was white British.¹⁹ The LBC study recruited people living in Edinburgh and the Lothians who were born either in 1921 or 1936. Individuals who self-reported stroke were removed for this study.

Demographic information for the 2,133 United States–based and 1,381 Scotland-based participants are presented in table 1.

Targeted Sequencing at 13q34 and Quality Control (GOCHA/ERICH and LINCHPIN)

We performed targeted sequencing across 559 Kbp at 13q34 (chr13:110,701,882–111,265,349) including *COL4A1/A2* for 2,133 individuals (1,055 ICH cases and 1,078 controls) in GOCHA/ERICH on the 96plex Nimblegen SeqCap platform and for 192 ICH cases in LINCHPIN on the NextSeq 550 platform. The reads with high quality are aligned to the human genome (GRCh37) using BWA (version 0.7.10).²⁰ We applied preprocessing steps including duplicate removal, local realignment, indel realignment using Genome Analysis Toolkit (GATK) IndelRealigner, and base quality recalibration using GATK Base Recalibration before the calling. Then GATK best practices pipeline was used for calling the variants from those 2 different platforms, separately. The variants were called and filtered using HaploTypeCaller.

Variants passing the GATK variant quality score recalibration metric were retained. We included only single nucleotide variants or indels with a depth of 10 or higher. We excluded variants with a call rate <0.98, case–control call rate difference >0.005, and differential missingness between cases and controls ($p < 0.05$). Samples were excluded if they had a low average call rate (<0.98), low mean sequence depth (<30), low mean genotype quality (<85), or differential missingness between cases and controls ($p < 0.05$) and Hardy-Weinberg equilibrium test (p value < 10^{-6}) on cases and controls separately.

Whole Genome Sequencing (LBC)

WGS of the LBC sample has been performed previously, with sequencing protocol and quality control (QC) metrics as reported.²¹ The variants selected from the GOCHA/ERICH and LINCHPIN studies were further evaluated in the LBC sequencing data as a control population.

Variant Selection and Annotation

We used the Ensembl variant effect predictor (VEP) software to annotate functional consequences of variants with the Ensembl annotation database on human genome assembly GRCh37.²² VEP provides various information of functional consequences of variants (splice acceptor variant, splice donor variant, start lost, stop lost, stop gained, frameshift variant, inframe insertion, inframe deletion, and missense variant) and functional impacts predicted by SIFT (deleterious or tolerate),²³ PolyPhen (probably damaging, possibly damaging, benign, or unknown),²⁴ combined annotation-dependent deletion (CADD),²⁵ and LoFtool (0–1).²⁶ We also used SnpEff, which categorizes the effects of variants by their genomic locations (e.g., introns, regulatory sites, splicing, missenses, nonsense) into high, moderate, low, or modifier.²⁷ We selected nonsynonymous variants with high/moderate by SnpEff, deleterious by SIFT, probably/possibly damaging by PolyPhen, CADD scores >20.0, and LoFtool scores <0.1 in the GOCHA/ERICH and LINCHPIN datasets.

Single-Variant and Variant-Set Association Tests in GOCHA/ERICH

We used the CATS online calculator²⁸ to estimate power to detect ICH association for single variant analysis. We performed single-variant association tests of those variants selected by annotation methods for the ICH risk by subtype (all-mixed, lobar, and nonlobar ICH) using logistic regression in R (r-project.org/). The association models were adjusted for age and sex. Significance thresholds for the single variant test was set as $\alpha = 0.05$ with Bonferroni correction for the numbers of single variants (number of variants: 39; $p < 1.28 \times 10^{-3}$).

Identification and Selection of Potential Causal Rare Variants in ICH (GOCHA/ERICH)

Because our sample size remained extremely underpowered for identifying rare functional variants by conventional association tests, we focused on rare nonsynonymous variants in *COL4A1/A2*, particularly those that occur in more than 1 ICH case but not in controls, or in more than 1 control but not in cases. Variants were first explored in GOCHA/ERICH as a discovery dataset and then examined in LINCHPIN and LBC as an external validation of the findings in an independent population.

We applied PLINK²⁹ software to assess linkage disequilibrium (LD) and haplotype inferences at this locus, testing whether the common *single nucleotide polymorphism* (SNP), rs9515201, previously identified in a GWAS of CSVD risk, lies within a haplotype with our identified rare variants.

Because the genetic architectures of lobar and nonlobar ICH are known to differ,⁵ we tested ICH subtype-specific effects for identified variants using the BinomiRare exact test,³⁰ which uses Poisson-binomial probabilities to calculate the association strength of variants based on the probabilities of diseased individuals carrying minor alleles of the variants under the null hypothesis that the variants are not associated with the disease. A classical logistic regression analysis was not applied for single-variant test because our selected variants occur only in 1 group (e.g., cases or controls), but not both.³¹ The BinomiRare regression models for ICH subtypes were adjusted for age and sex.

Our identified variants were evaluated using gnomAD³² for allele frequencies across populations and Geno2MP (geno2mp.gs.washington.edu/Geno2MP) for phenomic effect in carriers with variants or their family members.³³

To further assess our selected variants, we also explored the recent whole exome sequencing (WES) of 200,000 individuals in UK Biobank (UKB).^{34,35} We leveraged 137,766 unrelated British White individuals based on kinship relatedness (removing at least 1 of a related pair of individuals) and predicted ancestries from principal components for population structure provided by the UKB.³⁶ For ICH-related phenotypes, we used 2 traits in the electric health record including ICD-10 code for ICH (I61; UKB field number 41,202) and “vascular/heart problems diagnosed by doctor” (UKB field number 6,150).

Table 1 Sample Demographics

	GOCHA and ERICH studies			LINCHPIN		
	Lobar ICH	Nonlobar ICH	Controls	Lobar ICH	Nonlobar ICH	LBC, controls
Total (% female)	534 (50.9)	521 (38.35)	1,078 (44.9)	40 (52.5)	152 (40)	1189 (52)
Age, y, mean (SD)	73.0 (12.6)	69.3 (13.3)	70.2 (12.3)	80.2 (7.6)	70 (14.7)	77.9 (5.9)
ICH volume, mL, mean (SD)	36.0 (33.8)	15.1 (19.8)	—	—	—	—
ICH volume, mL, median (IQR)	24.2 (40.9)	8.3 (14.6)	—	—	—	—

Abbreviations: ERICH = Ethnic/Racial Variation in Intracerebral Hemorrhage; GOCHA = Gene Discovery for Warfarin-Related Intracerebral Hemorrhage; ICH = intracerebral hemorrhage; IQR = interquartile range; LBC = Lothian Birth Cohorts; LINCHPIN = Lothian study of Intracerebral Haemorrhage Pathology, Imaging and Neurologic Outcome.

The age at onset for these conditions was not available in the UKB database. The differences in the numbers of carriers of our selected variants and noncarriers between cases and controls in the UKB was examined by a χ^2 test.

Analysis of Previously Identified Variants on COL4A1/A2 in GOCHA/ERICH

We investigated coding variants in *COL4A1/A2* previously identified by other sequencing studies in ICH (rs200786329,¹³ rs117412802, rs62621875, and rs201105747¹⁷) or its related diseases including hereditary angiopathy, nephropathy, aneurysms, and cramps (rs113994104, rs113994105, and rs113994106),¹² small vessel disease (rs121912857 and rs113994107),⁹ cerebrovascular disease (rs672601346),³⁷ and porencephaly (rs113994112,⁸ rs113994114¹⁵). The information about their positions and amino acid changes are provided in table e-1 (data available from Dryad, doi.org/10.5061/dryad.z34tmpgcq). Previously reported variants that were also identified in our sequencing data were analyzed for association with ICH risk using BinomiRare.

External Validation of Selected Variants in Scotland-Based Datasets

Selected nonsynonymous rare variants appearing only in cases or controls in the United States–based dataset were explored for validation in LINCHPIN and LBC to determine whether they segregated with cases or controls in these independent datasets. Because LINCHPIN and LBC utilized different recruitment methods and were sequenced using different approaches, discovery of candidate variants in these datasets for validation in the United States–based GOCHA/ERICH cases was not feasible due to concerns for bias.

In Silico Modeling of Protein Structure and Thermal Stability of Identified Variants

We further assessed structural effects of selected variants on protein structures of *COL4A1/A2* using molecular dynamics (MD) simulation. The α chains in *COL4A1* and *COL4A2* (figure 1A) interact to form 1 triple-helical collagen type IV protomer, $\alpha1\alpha2(\text{IV})$, that consists of 3 protein domains: N-terminal 7S, central triple-helical collagen, and C-terminal NC1 (figure 1B).

Thus far, the triple-helical collagen structure for $\alpha1\alpha2(\text{IV})$ has not been experimentally determined in the Protein Data Bank (PDB; rcsb.org). Therefore, we predicted the tertiary heterotrimer structure of $\alpha1\alpha2(\text{IV})$ molecule using protein secondary/tertiary structure prediction tools including PSIPRED,³⁸ PFAM,³⁹ and BLASTp (blast.ncbi.nlm.nih.gov/Blast.cgi).⁴⁰ Briefly, we used the $\alpha1\alpha2(\text{IV})$ structure (PDB ID: 2CUO)⁴¹ as a template and predicted 2 structure models for our selected variant in *COL4A1* according to the number of copies of its mutations such as *COL4A1*^{WT/mt}/*COL4A2*^{WT} and *COL4A1*^{mt/mt}/*COL4A2*^{WT} and 1 model for the variant in *COL4A2* such as *COL4A1*^{WT/WT}/*COL4A2*^{mt} (figure 2A). To compare structural effects of our variants on $\alpha1\alpha2(\text{IV})$, we also generated additional structure models for 2 previously reported variants including rs200786329 (*COL4A1*)¹³ and rs117412802 (*COL4A2*).¹⁷ These 2 variants were chosen for the comparison because rs200786329^{*COL4A1*} and rs117412802^{*COL4A2*} were most recently identified in targeted sequencing of sporadic ICH cases, other than familial early-onset type IV collagen-related diseases. Moreover, our selected variants were located in the same Gly-Xaa-Yaa pattern or the same functional domain. Furthermore, rs200786329^{*COL4A1*} and rs117412802^{*COL4A2*} showed strong functional changes in terms of intracellular accumulation and extracellular deficiency of *COL4A1* or *COL4A2* protein levels compared to other variants.^{13,17}

MD simulations were performed with the predicted tertiary structures of $\alpha1\alpha2(\text{IV})$ molecules containing our identified variants and the 2 previously reported variants using DESMOND (deshawresearch.com/resources_desmond.html)⁴² as described⁴³ across 500 nanoseconds (ns). We included a modeling assumption that the atom positions of replaced amino acids stabilized after 100 ns from the start. Maestro (schrodinger.com/maestro; Schrödinger, LLC, 2016) was used to extract and analyze 800 sequential snapshots of simulated protein structures of *COL4A1* and *COL4A2* across 400 ns (100–500 ns).

In addition, the effect of the tripeptide composition (Gly-Xaa-Yaa) containing our identified variants on collagen triple helix stability was estimated using the collagen thermal stability calculator (compbio.cs.princeton.edu/csc),⁴⁴ which predicts the melting temperature (T_m).

A

Exon #48: rs77786415, rs200681591, rs200192119, rs12877501*, rs761217889, rs747313370

Exon #47: rs201158095, rs775127177, rs760207140, rs201300563, rs377451586

Exon #51: rs755291402

Exon #46: rs201790726

Exon #45: rs202207552

Exon #43: rs199956455

Exon #42: chr13:111155531

Exon #37: rs117412802*

Exon #31: rs1388143812

Exon #30: rs771353368

Exon #29: rs768662646

Exon #28: rs201058867, rs761169216

Exon #21: rs374820618

Exon #8: chr13:111082302

Exon #5: chr13:111077181

Exon #48: rs12877501*

Exon #46: rs760682735

Exon #48: rs12877501*

Exon #37: rs117412802*

Exon #25: rs145172612, chr13:110839649

Exon #22: rs570739274

Exon #17: rs370677625

Exon #6: rs138269346

Exon #3: rs34004222*

Exon #3: rs34004222*

GOCHA/ERICH

LINCHPIN

COL4A1

COL4A2

Recombination rate (cM/Mb)

110.8 Mb 110.9 Mb 111.0 Mb 111.1 Mb 111.2 Mb

46.18

39.99

28.74

rs9515201

Chromosome 13, 13q34

B

COL4A1

COL4A2

rs138269346

rs201716258

1110T

H203R

C-Terminal NC1

Triple-helix collagen

N-Terminal 7S

Structure of collagen type IV protomer

N-Terminal 7S

Triple-helix collagen

C-Terminal NC1

Col4a1

Col4a2

Col4a1

(A) Highly functional nonsynonymous variants appearing in ICH cases predicted by annotation methods including SnpEff (high or moderate), SIFT (deleterious), PolyPhen (probably/possibly damaging), combined annotation-dependent deletion (CADD) (>20.0), and LoFtool (>0.1). Green: N-terminal 7S domain; blue: triple helix collagen domain; brown: C-terminal NC domain. *Variants that exist in both datasets. (B) Selected COL4A1 and COL4A2 variants. ERICH = Ethnic/Racial Variation in Intracerebral Hemorrhage; GOCHA = Gene Discovery for Warfarin-Related Intracerebral Hemorrhage.

Data Availability

Sequencing data used in this study are available on dbGAP (ncbi.nlm.nih.gov/gap/; accession ID: phs000416.v2.p1). Additional data supporting these findings are available by the authors upon reasonable request.

Results

Identification and Genetic Association Tests of Rare Variants in *COL4A1/A2*

Following targeted sequencing and stringent QC of the 13q34 region in the United States–based GOCHA/ERICH studies, we retained 1,055 patients with ICH and 1,078 ICH-free controls with 11,815 variants. Among these variants, 0.91% were nonsynonymous (106 missense, 1 nonsense) and 0.70% were synonymous coding variants.

In the Scotland-based LINCHPIN study, 192 patients with ICH remained after QC with 1,380 variants at the 13q34 locus. From these, 1.01% were nonsynonymous (15 missense and 1 nonsense) and 1.88% were synonymous coding variants.

Selecting for nonsynonymous variants with high/moderate impact by SnpEff, deleterious predicted by SIFT, probably/possibly damaging predicted by PolyPhen, CADD scores >20.0, and LoFtool scores >0.1, we were left with 39 variants in GOCHA/ERICH and 4 variants in LINCHPIN, with 3 of these missense variants, including rs34004222, rs117412802, and rs12877501, common between studies. These nonsynonymous variants, predicted to be highly functional based on multiple annotation methods, are shown in figure 1A. None of these variants is in significant LD ($r^2 > 0.4$) with rs9515201, the leading common variant that we previously identified in a recent CSVD GWAS.⁶ Detailed information on these variants appears in table e-2 (data available from Dryad, doi.org/10.5061/dryad.z34tmpgcq). Annotation information for all detected variants in both sequencing datasets is described in detail in table e-3 (GOCHA/ERICH) and table e-4 (LINCHPIN) (data available from Dryad, doi.org/10.5061/dryad.z34tmpgcq).

According to our statistical power calculation (figure e-1, doi.org/10.5061/dryad.z34tmpgcq), we have extremely limited power (less than 20%) for detecting genetic association of such a rare variant (minor allele frequency [MAF] <0.001) with risk of sporadic ICH. We therefore limited single variant tests to those 39 variants but did not observe a variant with significant ($p < 1.2 \times 10^{-3}$) association with ICH subtypes (table e-5, data available from Dryad, doi.org/10.5061/dryad.z34tmpgcq).

Selection of Potential Causal Variants in ICH (GOCHA/ERICH)

Due to this expected low statistical power for single rare variant discovery in the United States–based GOCHA/ERICH dataset, we chose to focus on nonsynonymous mutations that appear in at least 2 participants exclusively in 1 group (e.g., ICH) but is absent in the other group (e.g., ICH-free controls). We

did not observe any nonsynonymous variants that occurred in at least 2 controls but not in cases, but we did identify 2 rare missense variants (rs138269346 and rs201716258) appearing in at least 2 ICH cases without occurrence in the controls (table 2). rs138269346 (*COL4A1* Ile110Thr) is in exon 5 of *COL4A1* corresponding to the N-terminal 7S domain and rs201716258 (*COL4A2* His203Leu) is in exon 10 of *COL4A2* corresponding to the triple-helix collagen domain where it affects an X residue of the Gly-Xaa-Yaa repeat (figure 1B). rs138269346^{COL4A1} was present in 4 ICH cases (2 lobar and 2 nonlobar ICH cases) and rs201716258^{COL4A2} was observed in another 2 ICH cases (1 lobar and 1 nonlobar ICH case). These ICH cases were each heterozygous for the minor alleles of these variants and no individual carried the minor alleles of both variants. These variants were not associated with age at disease onset, sex, or ICH volume (table e-6, data available from Dryad, doi.org/10.5061/dryad.z34tmpgcq).

According to our LD calculation and haplotype analysis, these 2 rare variants are not in LD ($r^2 < 0.01$) with the lead SNP of our previous GWAS, rs9515201,⁶ and we did not observe haplotypes carrying minor alleles of the identified rare variants and rs9515201 (table e-7, data available from Dryad, doi.org/10.5061/dryad.z34tmpgcq).

Functional Annotations of rs138269346 and rs201716258

Based on gnomAD, these 2 variants are rare in European populations (MAF <0.0005) and even rarer in other populations (table 2). Rs138269346^{COL4A1} was deemed pathogenic due to its annotation across multiple prediction tools: moderate by SnpEff, deleterious by SIFT, probably damaging by PolyPhen, and probably damaging by LoFtool. It showed a very high CADD score of 23.8, suggesting that this variant is among the top 0.42% ($10^{-2.38}$) of most predicted deleterious variants in the human genome. The other variant, rs201716258^{COL4A2}, was also considered to have a negative effect by SnpEff (moderate) and LoFtool (probably damaging), but not by SIFT (tolerated), PolyPhen (benign), and CADD (4.7).

Validation of rs138269346 and rs201716258 in Scotland-Based Participants

We set out to validate our identified variants using the independent LINCHPIN dataset. We detected 2 patients with ICH who carry the minor allele of rs201716258^{COL4A2} but no carriers for rs138269346^{COL4A1}. For rs201716258, both individuals developed nonlobar ICH in the right ventricle (ICH epicenter was lentiform) with a low small vessel disease score.⁴⁵ The ages of patients (1 male, 1 female) were 85 and 73 at the time of ICH, and there was no history of ischemic stroke.

Turning to the population-based LBC cohort, we identified 2 ICH-free controls who carry the minor allele of rs138269346^{COL4A1} but no carriers for rs201716258^{COL4A2}. One of these 2 ICH-free controls carrying the rs138269346^{COL4A1} in LBC self-reported a history of hypertension and myocardial infarction.

Figure 2 Protein Structure Prediction and Molecular Dynamics Simulation

(A) Procedure of protein structure prediction. (B) Comparison of physical lengths of protein prediction models affected by our identified variants including rs138269346 (COL4A1) and rs20176258 (COL4A2) in black as well as the previously reported variants including rs200786329 (COL4A1)¹⁷ and rs117412802 (COL4A2)²¹ in gray. SNP = single nucleotide polymorphism.

Further Evaluation of rs138269346 and rs201716258 in Geno2MP and UKB

Within individuals contributing to the Geno2MP database, carriers of these variants expressed phenotypic abnormalities previously

identified in patients with rare collagen IV mutation syndromes.¹⁴ In Geno2MP, we found 5 heterozygous rs138269346^{COL4A1} carriers and 6 heterozygous rs201716258^{COL4A2} carriers (table e-8, data available from Dryad, doi.org/10.5061/dryad.z34tmpgqc).

Table 2 Minor Allele Frequencies and Predicted Functional Effects of Identified Variants in the Discovery Dataset

Variant	Allele ref/alt	Gene			Protein	Minor allele frequencies in different populations						Predicted functional effect					
		N	Symbol	Exon	Type	Amino acids	Domain	European	African	East Asian	South Asian	Latino	SnPEff	SIFT	PolyPhen	CADD	LoFtool
rs138269346	A/G	4	COL4A1	5/52	Missense	I110T	N-75 domain	3.3 × 10 ⁻⁴	8.0 × 10 ⁻⁵	0	3.6 × 10 ⁻⁴	2.8 × 10 ⁻⁵	Moderate	Deleterious	Probably damaging	23.8	Probably damaging: -0.0123
rs201716258	A/G	2	COL4A2	10/48	Missense	H203R	Triple-helix	4.7 × 10 ⁻⁴	1.2 × 10 ⁻⁴	0	0	5.7 × 10 ⁻⁵	Moderate	Tolerated	Benign	4.7	Probably damaging: -0.0908

Abbreviations: CADD = combined annotation-dependent deletion; Ref/Alt: reference/alternative alleles.

One of the rs138269346^{COL4A1} carriers had multiple abnormalities in the eyes and the cardiovascular and nervous systems, and relatives of the other 4 carriers also had abnormalities in the eye, musculature, and cardiovascular and nervous systems. Two of rs201716258^{COL4A2} carriers had abnormalities in the ears and the cardiovascular system, respectively, and the relatives of the other 4 carriers also had abnormalities in the nervous system or musculature.

Among the 137,766 unrelated individuals in the 200K WES (table e-9), we found 1 rs138269346 carrier but none for rs201716258 out of 276 proxy patients with ICH (by ICD-10 code I61). Out of 41,032 individuals with vascular/heart problems, there are 33 carriers for rs138269346 and 63 carriers for rs201716258. According to our χ^2 analysis, rs138269346 was nominally associated with the proxy ICH ($\chi^2 = 3.4$; $p = 0.063$) although there is only 1 proxy ICH case carrying rs138269346. We did not observe significant association in the rest of the χ^2 tests (table e-10, data available from Dryad, doi.org/10.5061/dryad.z34tmpgcq).

Genetic Association Tests With ICH Subtypes

To determine which ICH subtype is relevant to these variants, we employed BinomiRare (table 3), which revealed that rs138269346^{COL4A1} is nominally associated with risk for nonlobar ICH ($p = 0.054$) and less associated with lobar ICH ($p = 0.077$) and all-mixed ICH ($p = 0.090$). However, rs201716258^{COL4A2} did not carry a significant association with any ICH subtype (all p values > 0.12).

Among the previously reported variants in COL4A1/A2, we could detect 3 nonsynonymous variants in COL4A2, including rs117412802, rs62621875, and rs201105747, which were previously reported in only patients with ICH and not in controls.¹⁷ However, in the GOCHA/ERICH dataset, we found carriers of these variants in both ICH cases (carriers of rs117412802: 20; and rs62621875: 1) and controls (rs117412802: 24; and rs62621875: 1). We identified rs201105747 only in 1 control. From the LINCHPIN dataset, we found only rs117412802 among 4 patients with ICH (3 lobar and 1 nonlobar ICH). We found no significant associations of these observed variants with ICH risk from BinomiRare (table e-11, data available from Dryad, doi.org/10.5061/dryad.z34tmpgcq).

In Silico Functional Analysis of rs138269346 and rs201716258

To gain insight into any structural potential consequences of these 2 missense variants, we performed MD simulations. Based on α chain composition of $\alpha 1\alpha 2$ (IV), each protomer can contain 1 or 2 variant $\alpha 1$ (IV) molecules and 1 $\alpha 2$ (IV). We therefore constructed 2 structural models for rs138269346^{COL4A1}: COL4A1s^{WT/I110T}/COL4A2^{WT} and COL4A1s^{I110T/I110T}/COL4A2^{WT}. Next, for rs201716258^{COL4A2}, 1 structure model was predicted: COL4A1s^{WT/WT}/COL4A2^{H203L} (figure 2A). From these simulations, we observed that 2 mutation structures of rs138269346^{COL4A1}, COL4A1s^{WT/I110T}/COL4A1^{WT} and COL4A1s^{I110T/I110T}/COL4A2^{WT}, result in average lengths

Table 3 Genetic Association Test Summary of the *COL4A1/A2* Variants Using BinomiRare

Variant ID (effect allele)	All ICH			Lobar ICH			Nonlobar ICH		
	N in cases	Expected N in cases	p Value	N in cases	Expected N in cases	p Value	N in cases	Expected N in cases	p Value
rs138269346 (G)	4	2.05	0.09	2	0.79	0.08	2	0.66	0.05
rs201716258 (T)	2	0.98	0.12	1	0.28	0.14	1	0.27	0.13

Abbreviation: ICH = intracerebral hemorrhage.

of 16.59 Å (equal to 10^{-10} m) (SD 5.58 Å) and 12.64 Å (SD 4.29 Å), respectively (figure 2B). These are 14.9% and 35.1% shorter than the average length of the wild type (average 19.49 Å; SD 6.41 Å). Likewise, we also found that the mutated structure for rs201716258^{COL4A2}, COL4A1^{WT/WT}/COL4A2^{H203L} (average 19.13 Å; SD 6.03 Å), became 15.7% shorter than the wild type (average 22.70 Å; SD 5.94 Å).

For comparison, we also generated structural models of the previously reported *COL4A1/A2* variants rs200786329^{COL4A1} and rs117412802^{COL4A2}. Interestingly, we found the same pattern in the simulations of the structure models with mutations of the previously reported variants. The average lengths of those mutant forms for rs200786329^{COL4A1} and rs117412802^{COL4A2} are 9.7%–25.8% shorter than their wild types (figure 2B).

It is well established that the amino acid sequence of the collagenous domain and mutations in collagen influence the thermal stability and melting temperature of the triple helix.⁴⁶ According to the collagen stability calculator,⁴⁴ COL4A1^{I100T} reduces predicted T_m by 2.2°C compared to COL4A1^{WT}, while COL4A2^{H203L} increases the T_m by 2.1–3.1°C compared to COL4A2^{WT} (figure e-2, data available from Dryad, doi.org/10.5061/dryad.z34tmgpcq), further lending support to their functionality.

Discussion

We conducted a targeted sequencing study of 1,055 ICH cases and 1,078 controls, and an additional dataset (192 ICH cases) at the 13q34 locus, which we and others have previously identified in genetic association studies for ICH and related manifestations of CSVD.^{6,7} 13q34 has been the target of prior sequencing studies in ICH, but the sample size in this current study ($n = 2,325$) is approximately 10 times larger than those prior efforts ($n < 200$).^{8,9,12,13,15,17,37} However, even with this boost in sample size, statistical power remains low for rare variants, so we focused on nonsynonymous variants appearing only in ICH cases but not in controls to restrict our search to the most likely pathogenic variants. Our study identified 2 novel rare missense variants in *COL4A1/A2*, appearing in cases only in our discovery dataset that (1) are predicted to have damaging effects, (2) are not in LD with common SNP identified in the previous GWAS for

CSVD, and (3) are predicted to substantially alter the physical length and the thermal stability of the type IV collagens. Furthermore, despite the limited statistical power of our sample size, we did observe a nominally significant association at rs138269346^{COL4A1} for nonlobar ICH risk.

In our attempt to externally validate these results in independent datasets, we detected 2 patients with ICH carrying rs201716258^{COL4A2} in LINCHPIN. No ICH case carriers for rs138269346^{COL4A1} were found, although 2 individuals with this variant were identified in the LBC control population. Given the size difference between the case populations in the 2 studies and the differences in recruitment between the case/control datasets and LBC, the significance of this replication failure at rs138269346^{COL4A1} is unclear.

The majority of mendelian disease-associated mutations in *COL4A1/A2* identified to date affect the Gly residue in the Gly-Xaa-Yaa repeat pattern, which is necessary for stable triple-helix formation, and more mutations have been described in *COL4A1* than *COL4A2*.¹⁴ However, the amino acids of our selected variants are located on the X residue in the pattern. Among the previously identified mutations for sporadic ICH, P352L^{rs200786329} (*COL4A1*)¹³ and E11223G^{rs117412802} (*COL4A2*)¹⁷ are also located on non-Gly residue in the tripeptide pattern, the Y and X residues, respectively. These previous variants (rs200786329 and rs117412802) significantly reduced the ratio of extracellular to intracellular *COL4A1* or *COL4A2* proteins compared to wild types in cultured cells, which is considered as a potential pathogenic mechanism underlying the type IV collagens related diseases.^{13,17} Interestingly, our MD simulation shows that these previous variants (rs200786329¹³ and rs117412802¹⁷) as well as our new variants (rs138269346 and rs20176258) substantially shorten the physical lengths of triple-helical structures, indicating that altered protein structures such as shortening of the helical domains could lead to changes in the protein functions including secretion of type IV collagen into extracellular matrix or secretion of mutant protein. In this regard, our thermal stability study predicted that COL4A1^{I100T} decreases (−2.2°C) and COL4A2^{H203L} increases (+2.1–3.1°C) T_m of collagen structures. These changes are not small in magnitude compared to the 2°C reduction affected by mendelian disease-associated Col2a1 mutations.⁴⁷ Therefore, it also may be possible that our identified variants exert their effects on type IV collagen

function through affecting the thermal stability of collagen, analogous to other collagen mutations.

Our study has several limitations. First, due to limited statistical power, we focused on rare variants appearing only in ICH cases but not controls, which could be considered conservative and increase false-negative associations for variants with low penetrance. Furthermore, we lack an analogous case/control ICH dataset with deep sequencing for direct validation of our observed variants. Targeted sequencing of 13q34 in LINCHPIN allowed us to verify the existence of rs201716258 in ICH cases, but the relatively small sample size and substantial difference in proportions of ICH subtypes compared to the discovery dataset makes the lack of observation of rs138269346 inconclusive. The population-based subjects from LBC are a welcome adjunct to the LINCHPIN ICH cases, but the varying burden of ICH-relevant disease histories and comorbidities and prospective nature of the LBC study, which is not continually updated, make interpretation of the appearance of rs138269346 in ICH-free individuals challenging. Due to the lack of genome-wide genotyping on the vast majority of subjects, population structure could not be assessed and adjusted for in our single variant association tests. While interesting and supportive of known phenotypic associations in rare *COL4A1/A2* syndromes, care must be taken in interpretation of phenotypes of variant carriers in Geno2MAP as this database is not a representative population sample and does not provide data to permit testing of statistical enrichment of variants. We observed differences in predicted effects of our identified variants across annotation methods. However, annotation methods could generate substantially different predictions of variant effects due to variation in the weighting of variant conservation and other features.^{48,49} Because of their rarity, imputation of these variants in existing ICH GWAS datasets using the HRC and TOPMed reference panels return very poor imputation quality scores, preventing additional forms of replication at this time.

Furthermore, there are challenges in integrating UKB data into our study. First, heterogeneous phenotyping of ICH cases in the UKB is a major concern that could substantially affect our genetic models.⁵⁰ For example, ICH status based on the ICD-10 code in electronic health records is less precise than manual phenotyping of ICH cases due to a lack of distinction between sporadic primary ICH and secondary ICH caused by trauma, brain tumor, hemorrhagic transformation of ischemic stroke, vascular malformation, and other factors, even among health care practitioners. In addition, the relatively young age, low number of ICH cases, and lack of ICH adjudication in the UKB make it ill-suited as a validation dataset for this application.

This work marks the continued progression of genetic research into collagen IV from gene mapping in mendelian disease through to identification of risk variants for sporadic disease in the general population. These sequencing results build on extant GWAS of ICH, demonstrating that *COL4A1* and *COL4A2* contribute to sporadic ICH not only through as-yet poorly understood mechanisms related to associations at common

variants but also potentially by rare variants that alter protein structure. Our observations from both sequenced datasets as shared through this article add substantially to our understanding of the burden and spectrum of *COL4A1/COL4A2* rare variation in sporadic ICH. Given the significance of 13q34 in both rare and common variant studies of ICH, studies building on our observations may link the pathogenic processes that underlie monogenic and polygenic ICH risk at this locus.

Study Funding

Supported by US-NINDS R01NS103924, US NHLBI RS&G 224, UK MRC MR/R005567-1, G0900428, G1002605, and 203699/Z/16/Z. Targeted sequencing services were provided by the Northwest Genomics Center at the University of Washington, Department of Genome Sciences, under US Federal Government contract number HHSN268201100037C from the National Heart, Lung, and Blood Institute (RS&G 224). T. Van Agtmael is supported by funding from the UK Medical Research Council (MRC; MR/R005567-1), Stroke Association (PPA 2016/02), and Heart Research UK (RG 2664/17/20). The LINCHPIN study was supported by an MRC/Stroke Association clinical research training fellowship (G0900428) awarded to Dr. Samarasekera, a Wellcome Trust clinical research training fellowship awarded to Dr. Rodrigues (203699/Z/16/Z), and an MRC senior clinical fellowship (G1002605) awarded to Prof. Al-Shahi Salman; the DNA sequencing was supported by a project grant from the MRC (MR/R005567/1). Phenotype collection in the Lothian Birth Cohort 1921 was supported by the United Kingdom's Biotechnology and Biological Sciences Research Council (BBSRC), The Royal Society, and The Chief Scientist Office of the Scottish Government. Phenotype collection in the Lothian Birth Cohort 1936 was supported by Age UK (The Disconnected Mind project). Whole-genome sequencing of the Lothian Birth Cohorts was funded by the BBSRC.

Disclosure

J. Chung, G. Hamilton, M. Kim, S. Marini, B. Montgomery, J. Henry, A.E. Cho, D.L. Brown, B.B. Worrall, J.F. Meschia, S.L. Silliman, M. Selim, D.L. Tirschwell, C.S. Kidwell, B. Kissela, S.M. Greenberg, A. Viswanathan, J.N. Goldstein, C.D. Langefeld, K. Rannikmae, C.L.M. Sudlow, N. Samarasekera, and M. Rodrigues report no disclosures. R. Al-Shahi Salman reports funding from the British Heart Foundation, The Stroke Association, and GE Healthcare Limited, paid to the University of Edinburgh, outside the submitted work. J.G.D. Prendergast, S.E. Harris, I. Deary, D. Woo, J. Rosand, and T. Van Agtmael report no disclosures. C.D. Anderson reports sponsored research support from the NIH, the American Heart Association, Massachusetts General Hospital, and Bayer AG, and has consulted for ApoPharma, Inc. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

Publication History

Received by *Neurology* November 17, 2020. Accepted in final form April 19, 2021.

Appendix Authors

Name	Location	Contribution
Jaeyoon Chung, PhD	Massachusetts General Hospital, Boston	Drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data
Graham Hamilton, PhD	University of Glasgow, Bearsden, UK	Drafting/revision of the manuscript for content, including medical writing for content; analysis or interpretation of data
Minsup Kim, PhD	Broad Institute, Boston, MA	Analysis or interpretation of data
Sandro Marini, MD	Korea University, Sejong, South Korea	Analysis or interpretation of data
Bailey Montgomery	Massachusetts General Hospital, Boston	Analysis or interpretation of data
Jonathan Henry	Massachusetts General Hospital, Boston	Analysis or interpretation of data
Art E. Cho, PhD	Korea University, Sejong, South Korea	Analysis or interpretation of data
Devin L. Brown, MD	University of Michigan, Ann Arbor	Drafting/revision of the manuscript for content, including medical writing for content
Bradford B. Worrall, MD	University of Virginia Health System, Charlottesville	Drafting/revision of the manuscript for content, including medical writing for content
James F. Meschia, MD	Mayo Clinic, Jacksonville, FL	Drafting/revision of the manuscript for content, including medical writing for content
Scott L. Silliman, MD	University of Florida College of Medicine, Jacksonville	Drafting/revision of the manuscript for content, including medical writing for content
Magdy Selim, MD, PhD	Beth Israel Deaconess Medical Center, Boston, MA	Drafting/revision of the manuscript for content, including medical writing for content
David L. Tirschwell, MD, MSc	University of Washington, Seattle	Drafting/revision of the manuscript for content, including medical writing for content
Chelsea S. Kidwell, MD	The University of Arizona, Tucson	Drafting/revision of the manuscript for content, including medical writing for content
Brett Kissela, MD, MS	University of Cincinnati, OH	Major role in the acquisition of data
Steven M. Greenberg, MD, PhD	Massachusetts General Hospital, Boston	Major role in the acquisition of data
Anand Viswanathan, MD, PhD	Massachusetts General Hospital, Boston	Major role in the acquisition of data

Appendix (continued)

Name	Location	Contribution
Joshua N. Goldstein, MD, PhD	Massachusetts General Hospital, Boston	Major role in the acquisition of data
Carl D. Langefeld, PhD	Wake Forest School of Medicine, Winston-Salem, NC	Major role in the acquisition of data
Kristiina Rannikmae, MD	Wake Forest School of Medicine, Winston-Salem, NC	Major role in the acquisition of data
Catherine L.M. Sudlow, BMBCh, MSc, DPhil, FRCPE	Wake Forest School of Medicine, Winston-Salem, NC	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data
Neshika Samarasekera, PhD, MRCP	University of Edinburgh, Scotland, UK	Major role in the acquisition of data
Mark Rodrigues, PhD, FRCR	University of Edinburgh, Scotland, UK	Major role in the acquisition of data
Rustam Al-Shahi Salman, PhD, FRCP Edin	University of Edinburgh, Scotland, UK	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data
James G.D. Prendergast, PhD	University of Edinburgh, Scotland, UK	Major role in the acquisition of data
Sarah E. Harris, PhD	University of Edinburgh, Scotland, UK	Major role in the acquisition of data
Ian Deary, MBChB, PhD, FRCP	University of Edinburgh, Scotland, UK	Major role in the acquisition of data
Daniel Woo, PhD	University of Cincinnati, OH	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data
Jonathan Rosand, MD, MSc	Massachusetts General Hospital, Boston	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data
Tom Van Agtmael, PhD	University of Cincinnati, OH	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; analysis or interpretation of data
Christopher D. Anderson, MD, MMSc	Massachusetts General Hospital, Boston	Drafting/revision of the manuscript for content, including medical writing for content; major role in the acquisition of data; study concept or design; analysis or interpretation of data

References

- Qureshi AI, Tuhim S, Broderick JP, Batjer HH, Hondo H, Hanley DF. Spontaneous intracerebral hemorrhage. *N Engl J Med*. 2001;344(19):1450-1460.
- Feigin VL, Norrving B, Mensah GA. Global burden of stroke. *Circ Res*. 2017;120(3):439-448.
- Woo D, Haverbusch M, Sekar P, et al. Effect of untreated hypertension on hemorrhagic stroke. *Stroke*. 2004;35(7):1703-1708.
- Marini S, Devan WJ, Radmanesh F, et al. 17p12 influences hematoma volume and outcome in spontaneous intracerebral hemorrhage. *Stroke*. 2018;49(7):1618-1625.
- Woo D, Falcone GJ, Devan WJ, et al. Meta-analysis of genome-wide association studies identifies 1q22 as a susceptibility locus for intracerebral hemorrhage. *Am J Hum Genet*. 2014;94(4):511-521.
- Chung J, Marini S, Pera J, et al. Genome-wide association study of cerebral small vessel disease reveals established and novel loci. *Brain*. 2019;142(10):3176-3189.
- Rannikmäe K, Davies G, Thomson PA, et al. Common variation in COL4A1/COL4A2 is associated with sporadic cerebral small vessel disease. *Neurology*. 2015;84(9):918-926.
- Gould DB, Phalan FC, Breedveld GJ, et al. Mutations in Col4a1 cause perinatal cerebral hemorrhage and porencephaly. *Science*. 2005;308(5725):1167-1171.
- Gould DB, Phalan FC, van Mil SE, et al. Role of COL4A1 in small-vessel disease and hemorrhagic stroke. *N Engl J Med*. 2006;354(14):1489-1496.
- Sibon I, Coupry I, Menegon P, et al. COL4A1 mutation in Axenfeld-Rieger anomaly with leukoencephalopathy and stroke. *Ann Neurol*. 2007;62(2):177-184.
- Van Agtmael T, Schlotzer-Schrehardt U, McKie L, et al. Dominant mutations of Col4a1 result in basement membrane defects which lead to anterior segment dysgenesis and glomerulopathy. *Hum Mol Genet*. 2005;14(21):3161-3168.
- Plaisier E, Gribouval O, Alamowitch S, et al. COL4A1 mutations and hereditary angiopathy, nephropathy, aneurysms, and muscle cramps. *N Engl J Med*. 2007;357(26):2687-2695.
- Weng YC, Sonni A, Labelle-Dumais C, et al. COL4A1 mutations in patients with sporadic late-onset intracerebral hemorrhage. *Ann Neurol*. 2012;71(4):470-477.
- Meuwissen ME, Halley DJ, Smit LS, et al. The expanding phenotype of COL4A1 and COL4A2 mutations: clinical data on 13 newly identified families and a review of the literature. *Genet Med*. 2015;17(11):843-853.
- de Vries LS, Koopman C, Groenendaal F, et al. COL4A1 mutation in two preterm siblings with antenatal onset of parenchymal hemorrhage. *Ann Neurol*. 2009;65(1):12-18.
- Woo D, Rosand J, Kidwell C, et al. The Ethnic/Racial Variations of Intracerebral Hemorrhage (ERICH) study protocol. *Stroke*. 2013;44(10):e120-e125.
- Jeanne M, Labelle-Dumais C, Jorgensen J, et al. COL4A2 mutations impair COL4A1 and COL4A2 secretion and cause hemorrhagic stroke. *Am J Hum Genet*. 2012;90(1):91-101.
- Samarasekera N, Fonville A, Lerpiniere C, et al. Influence of intracerebral hemorrhage location on incidence, characteristics, and outcome: population-based study. *Stroke*. 2015;46(2):361-368.
- Deary IJ, Gow AJ, Pattie A, Starr JM. Cohort profile: the Lothian Birth cohorts of 1921 and 1936. *Int J Epidemiol*. 2012;41(6):1576-1584.
- Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics*. 2009;25(14):1754-1760.
- Halachev M, Meynert A, Taylor MS, et al. Increased ultra-rare variant load in an isolated Scottish population impacts exonic and regulatory regions. *PLoS Genet*. 2019;15(11):e1008480.
- McLaren W, Gil L, Hunt SE, et al. The Ensembl variant effect predictor. *Genome Biol*. 2016;17(1):122.
- Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc*. 2009;4:1073-1081.
- Adzhubei IA, Schmidt S, Peshkin L, et al. A method and server for predicting damaging missense mutations. *Nat Methods*. 2010;7(4):248-249.
- Kircher M, Witten DM, Jain P, O'Roak BJ, Cooper GM, Shendure J. A general framework for estimating the relative pathogenicity of human genetic variants. *Nat Genet*. 2014;46(3):310-315.
- Fadista J, Oskolkov N, Hansson O, Groop L. LoFtool: a gene intolerance score based on loss-of-function variants in 60 706 individuals. *Bioinformatics*. 2017;33(4):471-474.
- Cingolani P, Patel VM, Coon M, et al. Using *Drosophila melanogaster* as a model for genotoxic chemical mutational studies with a new Program, SnpSift. *Front Genet*. 2012;3:35.
- Skol AD, Scott LJ, Abecasis GR, Boehnke M. Joint analysis is more efficient than replication-based analysis for two-stage genome-wide association studies. *Nat Genet*. 2006;38(2):209-213.
- Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet*. 2007;81(3):559-575.
- Sofer T, BinomiRare: a robust test of the association of a rare variant with a disease for pooled analysis and meta-analysis, with application to the HCHS/SOL. *Genet Epidemiol*. 2017;41(5):388-395.
- Sofer T, Lee J, Kurniansyah N, et al. BinomiRare: a carriers-only test for association of rare genetic variants with a binary outcome for mixed models and any case-control proportion. *medRxiv*. 2021. doi: 10.1101/2021.01.08.21249450.
- Karczewski KJ, Francioli LC, Tiao G, et al. The mutational constraint spectrum quantified from variation in 141,456 humans. *Nature*. 2020;581(7809):434-443.
- Wang J, Al-Ouran R, Hu Y, et al. MARRVEL: integration of human and model organism genetic resources to facilitate functional annotation of the human genome. *Am J Hum Genet*. 2017;100(6):843-853.
- Szuskowski JD, Balasubramanian S, Sasson A, et al. Advancing human genetics research and drug discovery through exome sequencing of the UK Biobank. *medRxiv*. 2020. doi: 10.1101/2020.11.02.20222232.
- Jurgens SJ, Choi SH, Morrill VN, et al. Rare genetic variation underlying human diseases and traits: results from 200,000 individuals in the UK Biobank. *bioRxiv*. 2020. doi: 10.1101/2020.11.29.402495.
- Wu Y, Byrne EM, Zheng Z, et al. Genome-wide association study of medication-use and associated disease in the UK Biobank. *Nat Commun*. 2019;10(1):1891.
- Shah S, Kumar Y, McLean B, et al. A dominantly inherited mutation in collagen IV A1 (COL4A1) causing childhood onset stroke without porencephaly. *Eur J Paediatr Neurol*. 2010;14(2):182-187.
- McGuffin LJ, Bryson K, Jones DT. The PSIPRED protein structure prediction server. *Bioinformatics*. 2000;16(4):404-405.
- El-Gebali S, Mistry J, Bateman A, et al. The Pfam protein families database in 2019. *Nucleic Acids Res*. 2019;47(D1):D427-D432.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol*. 1990;215(3):403-410.
- Hongo C, Noguchi K, Okuyama K, Tanaka Y, Nishino N. Repetitive interactions observed in the crystal structure of a collagen-model peptide, [(Pro-Pro-Gly)9]3. *J Biochem*. 2005;138(2):135-144.
- Bowers KJ, Chow DE, Xu H, et al. Scalable algorithms for molecular dynamics simulations on commodity clusters. Proceedings of the 2006 ACM/IEEE Conference on Supercomputing; 11-17 Nov, 2006; 43-43.
- Kim M, Kwon J, Goo JI, Choi Y, Cho AE. Elucidation of the inhibition mechanism of sulfiredoxin using molecular modeling and development of its inhibitors. *J Mol Graph Model*. 2019;92:208-215.
- Persikov AV, Ramshaw JA, Brodsky B. Prediction of collagen stability from amino acid sequence. *J Biol Chem*. 2005;280(19):19343-19349.
- Arba F, Leigh R, Inzitari D, Warach SJ, Luby M, Lees KR. Blood-brain barrier leakage increases with small vessel disease in acute ischemic stroke. *Neurology*. 2017;89(21):2143-2150.
- Bächinger HP, Davis JM. Sequence specific thermal stability of the collagen triple helix. *Int J Biol Macromol*. 1991;13(3):152-156.
- Chakkalakal SA, Heilig J, Baumann U, Paulsson M, Zaucke F. Impact of arginine to cysteine mutations in collagen II on protein secretion and cell survival. *Int J Mol Sci*. 2018;19(2):541.
- McCarthy DJ, Humburg P, Kanapin A, et al. Choice of transcripts and software has a large effect on variant annotation. *Genome Med*. 2014;6(3):26.
- Miller JE, Veturi Y, Ritchie MD. Innovative strategies for annotating the "relationSNP" between variants and molecular phenotypes. *BioData Min*. 2019;12:10.
- Rubbo B, Fitzpatrick NK, Denaxas S, et al. Use of electronic health records to ascertain, validate and phenotype acute myocardial infarction: a systematic review and recommendations. *Int J Cardiol*. 2015;187:705-711.

Neurology®

Rare Missense Functional Variants at *COL4A1* and *COL4A2* in Sporadic Intracerebral Hemorrhage

Jaeyoon Chung, Graham Hamilton, Minsup Kim, et al.

Neurology 2021;97:e236-e247 Published Online before print May 24, 2021

DOI 10.1212/WNL.00000000000012227

This information is current as of May 24, 2021

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/97/3/e236.full
References	This article cites 49 articles, 10 of which you can access for free at: http://n.neurology.org/content/97/3/e236.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Cerebrovascular disease/Stroke http://n.neurology.org/cgi/collection/all_cerebrovascular_disease_stroke Association studies in genetics http://n.neurology.org/cgi/collection/association_studies_in_genetics Intracerebral hemorrhage http://n.neurology.org/cgi/collection/intracerebral_hemorrhage
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2021 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.