Teaching Video NeuroImage: Dissociation of Abdominal Reflexes in Rheumatoid Atlantoaxial Subluxation

Igor Vilela Brum, MD, and Guilherme Diogo Silva, MD

Neurology® 2022;98:e211-e212. doi:10.1212/WNL.00000000000013007

Correspondence

Dr. Brum
igor.brum@hc.fm.usp.br

Figure Spinal Cord Compression by Atlantoaxial Subluxation and Retrodental Pannus

T2-weighted sagittal (A) spine MRI showing spinal cord compression and high signal intensity at the level of C1-C2 (dashed arrow) because of anterior subluxation of the atlas (solid arrow) and retrodental pannus (small arrow) in a patient with rheumatoid arthritis. T2-weighted axial (B) image confirms spinal cord compression by atlantoaxial subluxation (arrow, B).

A 59-year-old woman with rheumatoid arthritis presented with progressive gait difficulties and electric shock-like pain triggered by neck flexion (Lhermitte sign) for more than 6 months. On examination, she had hyperreflexia in the 4 limbs and brisk deep abdominal reflexes with absent superficial abdominal reflexes (reflex dissociation) (Video 1). MRI showed spinal cord compression resulting from anterior subluxation of the atlas and retrodental pannus (synovial tissue proliferation) (Figure). Superficial abdominal reflexes may be absent in 20% of normal individuals.¹ However, the dissociation of abdominal reflexes suggests an upper motor neuron lesion above the T6 spinal segment.²

Study Funding

The authors report no targeted funding.

Disclosure

I.V. Brum reports no disclosures relevant to the manuscript; G.D. Silva reports no disclosures relevant to the manuscript. Go to [Neurology.org/N](https://www.neurology.org/N) for full disclosures.

MORE ONLINE

Teaching slides

links.lww.com/WNL/B633

Video

Appendix Authors

Name	Location	Contribution
Igor Vilela Brum, MD	Department of Neurology, University of Sao Paulo, Brazil	Drafting/revision of the manuscript for content, including medical writing for content, major role in the acquisition of data, study concept or design, and analysis or interpretation of data
Guilherme Diogo Silva, MD	Department of Neurology, University of Sao Paulo, Brazil	Drafting/revision of the manuscript for content, including medical writing for content

References

1. Madonick MJ. Statistical control studies in neurology: 8. The cutaneous abdominal reflex. *Neurology*. 1957;7(7):459-465.
2. Teasdall RD, Van Den Ende H. A note on the deep abdominal reflex. *J Neurol Neurosurg Psychiatry*. 1982;45(4):382-383.

Neurology®

Teaching Video NeuroImage: Dissociation of Abdominal Reflexes in Rheumatoid Atlantoaxial Subluxation

Igor Vilela Brum and Guilherme Diogo Silva

Neurology 2022;98:e211-e212 Published Online before print October 21, 2021

DOI 10.1212/WNL.00000000000013007

This information is current as of October 21, 2021

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/98/2/e211.full
References	This article cites 2 articles, 2 of which you can access for free at: http://n.neurology.org/content/98/2/e211.full#ref-list-1
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): All Spinal Cord http://n.neurology.org/cgi/collection/all_spinal_cord Clinical neurology examination http://n.neurology.org/cgi/collection/clinical_neurology_examination MRI http://n.neurology.org/cgi/collection/mri
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2021 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.