Neurology®

The most widely read and highly cited peer-reviewed neurology journal
The Official Journal of the American Academy of Neurology

Neurology Publish Ahead of Print

DOI: 10.1212/WNL.0000000000200008

A 50-Year-Old Man With Progressive Limb Weakness and Slurred Speech

Author(s):

Nina Xie, MD^{1,2}; Guang Yang, MD PhD³; Weiru Zhang, MD³; Hongwei Xu, MD^{1,2}; Qiying Sun, MD PhD^{1,2}

Corresponding Author:

Qiying Sun

sunqiying2015@163.com

Affiliation Information for All Authors: 1 Department of Geriatric Neurology, Xiangya Hospital, Central South University, Changsha 410008, China; 2 National Clinical Research Center for Geriatric Disorders, Changsha, Hunan 410078, China; 3 Department of General Medicine, Xiangya Hospital, Central South University, Changsha 410008, China.

Neurology® Published Ahead of Print articles have been peer reviewed and accepted for publication. This manuscript will be published in its final form after copyediting, page composition, and review of proofs. Errors that could affect the content may be corrected during these processes.

Contributions:

Nina Xie: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data; Study concept or design; Analysis or interpretation of data

Guang Yang: Drafting/revision of the manuscript for content, including medical writing for content; Analysis or interpretation of data

Weiru Zhang: Drafting/revision of the manuscript for content, including medical writing for content; Analysis or interpretation of data

Hongwei Xu: Drafting/revision of the manuscript for content, including medical writing for content; Analysis or interpretation of data

Qiyi Sun: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data; Study concept or design; Analysis or interpretation of data

Number of characters in title: 10

Abstract Word count: 126

Word count of main text: 1484

References: 13

Figures: 2

Tables: 1

Supplemental: patient consent-to-disclose form CARE checklistTable 1

Search Terms: [178] Amyotrophic lateral sclerosis

Acknowledgment: We thank the patient and his family for granting permission to publish this information.

Study Funding: The authors report no targeted funding

Disclosures: The authors report no disclosures relevant to the manuscript.

Abstract

A 50-year-old man presented with a nine-month history of progressive left arm weakness and dysarthria. Family history showed that his parents are cousins and one of his siblings died of motor neuron disease. Brain MRI showed T2-weighted white matter hyperintensities along the course of pyramidal tracts. Neurological examination and EMG revealed upper and lower motor neuron signs involving the bulbar, cervical, thoracic, and lumbosacral segments, which meets the criteria of a definite amyotrophic lateral sclerosis (ALS), according to the revised El Escorial criteria. Whole-exome genetic sequencing found two novel *LYST* missense mutations, confirming the diagnosis of Chediak Higashi syndrome (CHS), a rare autosomal-recessive hematological disorder. Our case indicates that CHS can present as ALS phenotype and the *LYST* might be a novel causative gene for ALS.

Section 1

A 50-year-old man with progressive left arm weakness was referred for evaluation. He had been well until nine months prior when he first noticed the left hand was not as flexible as usual. Gradually he had difficulty lifting heavy things. Seven months ago, his speech also became slurred. No other symptoms were reported. Six months ago, he was referred to us. Past medical history was unremarkable. Toxin exposure and substance abuse were denied. His parents are cousins. Neither of them has similar symptoms. One of his siblings was clinically diagnosed with motor neuron disease and died at the age of 37.

Upon neurological examination, he was alert, oriented, and slightly dysarthric. Mini-Mental State Exam scored 27. Cranial nerve tests revealed central facial palsy, lingual palsy, and tongue fasciculations (Video 1). The left-arm strength was rated 4/5 both proximally and distally, with increased muscle tone and brisk deep tendon reflexes. The strength of other limbs was rated 5/5. Hoffman and Babinski's signs were present bilaterally. Sensation, coordination, and gait were normal. General examinations for skin, heart, lung, and abdomen were unremarkable.

- 1. Where would you localize the lesion?*
- 2. What would be your initial lab workup?*

Section 2

The central facial palsy, lingual palsy, left arm weakness with increased muscle tone and hyperreflexia, and presence of pathological reflexes suggest upper motor neuron (UMN) damage involving the bulbar, cervical, and lumbosacral segments. The tongue fasciculations indicate lower motor neuron (LMN) damage in the bulbar segment. Given the presence of both upper and lower motor neuron signs, motor neuron disease

and its mimics are suspected. The differential diagnosis is extensive, including amyotrophic lateral sclerosis (ALS), amyotrophic lateral sclerosis -frontotemporal dementia (ALS-FTD), infection, metabolic abnormality, paraneoplastic syndromes, and hereditary disorders. Therefore, initial lab workup should include routine screenings for these etiologies, as well as nerve conduction studies/ electromyography (NCS/EMG) and cerebrospinal MRI.

The following investigations were unremarkable or negative: complete blood count, urine and stool analysis, liver and renal function, cardiac enzymes, electrolytes, blood glucose, lipids, coagulation, thyroid function, homocysteine, vitamins, serum tumor markers, paraneoplastic antibodies (anti-Hu, Yo, Ri, Ma2/Ta, CV2/CRMP5 and amphiphysin), infectious panel (tuberculosis, hepatitis, HIV, Lyme, and syphilis), urine and serum protein electrophoresis, very-long-chain fatty acids (VLCFA), and electrocardiogram. Brain MRI showed symmetric T2-hyperintense signals along the course of pyramidal tracts (Figure 1). Cervical spinal MRI and chest-abdomen-pelvis CT were normal, as was the initial NCS/EMG result done three months after onset.

What differential diagnosis would you consider at this stage?

Section 3

According to the revised EI Escorial criteria, this patient has UMN and LMN signs in one region and UMN signs in three regions, suggesting a possible ALS¹. The MRI findings also support this diagnosis. However, it is uncommon that the NCS/EMG test did not show any signs of neurogenic impairment. Alternative diagnosis needs to be considered. The absence of cognitive decline excludes ALS-FTD. Infectious causes such as Lyme disease, HIV, and syphilitic amyotrophy and metabolic disorders such as vitamin 12 deficiency and hyperthyroidism were also ruled out by the negative blood tests and history. The normal serum tumor markers, negative paraneoplastic antibodies, absence of M protein, and normal chest-abdomen-pelvis CT argue against

a paraneoplastic syndrome. Given the positive family history, a hereditary neurodegenerative disorder is the most likely diagnosis at this stage. Several possibilities are discussed below.

Adrenomyeloneuropathy (AMN) is a variant of adrenoleukodystrophy. It is caused by mutations in the *ABCD1* gene and has an X-linked recessive mode of inheritance. Typical onset age is in the late twenties. AMN can present as mixed UMN and LMN signs by causing myelopathy and peripheral neuropathy, with or without adrenal insufficiency. Less frequently, brain MRI can show cerebral demyelination. In patients with milder phenotypes, VLCFA levels can be normal^{2,3}. However, the onset age and absence of myelopathy and peripheral neuropathy in this patient make AMN less likely. Moreover, lingual palsy is very uncommon for AMN.

Spinocerebellar ataxia 3 (SCA3) is caused by the CAG repeat expansion mutations in the *ATXN3* gene. It can cause spasticity and tongue fasciculations in patients with early-onset age. However, most SCA3 is autosomal dominantly inherited. Had SCA3 been the diagnosis, at least one of the parents should be symptomatic⁴. Besides, there is no ataxia in this patient.

Adult hexosaminidase deficiency is caused by mutations in the *HEXA* gene and inherited in an autosomal recessive pattern. It is easily mistaken for ALS in the early stages, as neurologic symptoms such as spasticity, progressive muscle weakness, and dysarthria are often the initial manifestations. However, Adult hexosaminidase deficiency is more commonly seen in Ashkenazi Jewish, Celtic, and French-Canadian populations. NCS/EMG can detect signs of chronic anterior horn cell neuropathy⁵. This patient is of Asian descent and has normal NCS/EMG results.

Another autosomal recessive disorder that can cause UMN signs, LMN signs, and diffuse T2-weighted white matter hyperintensities is adult polyglucosan body disease (APBD). It is caused by mutations in the *GBE1* gene. Mutant GBE, a glycogen-branching enzyme, leads to the accumulation of misfolded glycogens, i.e. the polyglucosan bodies, in various tissues including muscles, peripheral nervous system, and central nervous system. The observation of periodic acid-Schiff-positive polyglucosan bodies on skin or peripheral nerve biopsy is pathognomonic⁶. However, the most common initial sign of APBD is urinary incontinence, which is not reported in this patient.

What would you do next to clarify the diagnosis?

Section 4

Although the possibility of diseases discussed above was relatively low, they still cannot be excluded. Genetic testing is warranted. Whole-exome genetic sequencing revealed compound heterozygous missense mutations (Exon 3: p.Val24Ala; Exon 21: p.Arg1986Gln) in the patient's *LYST* gene, which have not been reported previously. His parents are mutation carriers (Figure 2). The diagnosis of Chediak Higashi syndrome (CHS) was established. We further examined his peripheral blood smear. No giant cytoplasmic granules were observed in leukocytes. During the six-month follow-up period, he developed bucking when drinking water. The left limb weakness and dysarthria were not improved. Palmomental reflexes appeared bilaterally. Repeat needle EMG done nine months after onset indicated acute and chronic neurogenic impairment involving the bulbar, cervical, thoracic, and lumbosacral segments (eTable 1). Nerve conduction studies were normal. The EMG results and physical signs together suggest a definite ALS¹.

Discussion

CHS is a rare autosomal-recessive disorder caused by mutations in the *LYST* gene. *LYST* is a lysosomal trafficking regulator. Mutant *LYST* causes lysosomal dysfunction, leading to aberrant organellar protein trafficking and incompetent immunocytes. Genotypes of CHS are highly diverse. Loss-of-function and missense mutations correlate with typical and atypical CHS, respectively⁷. The typical form is characterized by immunodeficiency, oculocutaneous albinism, coagulation defects, a high risk of developing hemophagocytic lymphohistiocytosis (HLH), and presence of giant cytoplasmic granules in leukocytes. Patients usually die in the first decade without allogeneic hematopoietic cell transplantation (HCT).

In contrast, patients with atypical CHS survive into adulthood and manifest as progressive neurodegeneration, including parkinsonism, ataxia, peripheral neuropathy, and cognitive decline. Electroencephalogram may detect epileptiform discharges. Brain MRI may show cerebral atrophy. Symptomatic treatments are the mainstay option. Although atypical CHS patients escape the life-threatening complications in childhood, they are still at the risk of developing HLH by suddenly entering an accelerated phase. The outcome will be fatal without timely HCT. However, early diagnosis of atypical CHS is challenging due to the absence of hematological presentations^{8,9}. Family history provides critical diagnostic clues.

Our case described an atypical CHS patient with an ALS phenotype. To our knowledge, this was first reported in CHS. To confirm the diagnosis, gene analysis found two novel *LYST* missense mutations. As aforementioned, *LYST* regulates lysosome functions. The autophagy-endosome-lysosomal system plays a critical role in ALS pathogenesis. Several genes involved in the endolysosomal pathway, including *CHMP2B*, *VCP*, *ALS2*, have proved to be associated with a risk of ALS¹⁰⁻¹². Moreover, using the cDNA array and RT-PCR method, Daniel Offen et al. reported that the *LYST* mRNA level is markedly increased in the postmortem spinal cord

specimens of four sporadic ALS patients compared to controls¹³. These findings together with our case indicate that the *LYST* gene may be associated with ALS.

Notably, the giant cytoplasmic granules in leukocytes, formed by dysregulated lysosomes and vesicles, are thought to be a pathognomonic feature for CHS. However, they are not present in our patient's blood smear, although Wendy J. Introne et al. observed these granules in leukocytes of atypical CHS patients⁹. Given the absence of giant cytoplasmic granules and the lack of strong experimental data on the pathophysiological role of *LYST* in human motor neurons, the co-occurrence of ALS phenotype and *LYST* in our patient may be coincidental. To clarify whether there is a true causative relationship between them, *LYST* sequencing should be considered in ALS patients and more functional studies are needed. These data will be beneficial not only for understanding the pathophysiology of congenital immunodeficiency but also for ALS.

In summary, we reported a rare CHS case presenting as ALS. Our findings expanded the clinical phenotypes of CHS and implicated that *LYST* might be a novel causative gene for ALS.

Ethics approval

Written consent form was obtained from the patient. This study was approved by the Medical Ethics Committee of Central South University.

Figure 1 brain MRI

(A-C) FLAIR images showed symmetric hyperintense signals along the course of pyramidal tracts.

Figure 2 pedigree and sequencing results

(A) His parents are mutation carriers. A sibling died of motor neuron disease. (B-C) Sequence chromatogram showing the two novel compound heterozygous mutations (Exon 3: p.Val24Ala; Exon 21: p.Arg1986Gln) in the patient's *LYST* gene

Video 1 tongue fasciculation

Appendix 1. Authors

Name	Location	Contribution
Nina Xie MD	Department of Geriatric neurology, Xiangya Hospital, Central South University, Changsha 410008, China	Data collection and analysis Manuscript preparation
Guang Yang MD PhD	Department of General medicine, Xiangya Hospital, Central South University, Changsha 410008, China	Data collection and analysis Manuscript preparation
Weiru Zhang MD	Department of General medicine, Xiangya Hospital, Central South University, Changsha 410008, China	Data collection and analysis Manuscript revision
Hongwei Xu MD	Department of Geriatric neurology, Xiangya Hospital, Central South University, Changsha 410008, China	Data collection and analysis Manuscript revision
Qiyang Sun MD PhD	Department of Geriatric neurology, Xiangya Hospital, Central South University, Changsha 410008, China	Data collection and analysis Manuscript preparation

WNL-2022-200303_sup1 --<http://links.lww.com/WNL/B812>

WNL-2022-200303_vid1 -- <http://links.lww.com/WNL/B813>

Reference

1. van Es MA, Hardiman O, Chio A, et al. Amyotrophic lateral sclerosis. *The Lancet*. 2017;390(10107):2084-2098.
2. Azar C, Shor N, Nadjar Y. Adrenomyeloneuropathy Masquerading as Chronic Myelitis. *JAMA Neurol*. 2020;77(4):522-523.
3. van Geel BM, Bezman L, Loes DJ, Moser HW, Raymond GV. Evolution of phenotypes in adult male patients with X-linked adrenoleukodystrophy. *Annals of Neurology*. 2001;49(2):186-194.
4. França MC Jr DAA, Nucci A, Lopes-Cendes I. Muscle excitability abnormalities in Machado-Joseph disease. *Arch Neurol*. 2008;65(4):525-529.
5. Jahnova H, Poupetova H, Jireckova J, et al. Amyotrophy, cerebellar impairment and psychiatric disease are the main symptoms in a cohort of 14 Czech patients with the late-onset form of Tay-Sachs disease. *J Neurol*. 2019;266(8):1953-1959.

6. Souza PVS, Badia BML, Farias IB, et al. GBE1-related disorders: Adult polyglucosan body disease and its neuromuscular phenotypes. *Journal of Inherited Metabolic Disease*. 2021;44(3):534-543.
7. Nagai K, Ochi F, Terui K, et al. Clinical characteristics and outcomes of chediak-Higashi syndrome: a nationwide survey of Japan. *Pediatr Blood Cancer*. 2013;60(10):1582-1586.
8. Mathis S, Cintas P, de Saint-Basile G, Magy L, Funalot B, Vallat JM. Motor neuronopathy in Chediak–Higashi syndrome. *Journal of the Neurological Sciences*. 2014;344(1):203-207.
9. Introne WJ, Westbroek W, Cullinane AR, et al. Neurologic involvement in patients with atypical Chediak-Higashi disease. *Neurology*. 2016;86(14):1320.
10. Xie Y, Zhou B, Lin MY, Wang S, Foust KD, Sheng ZH. Endolysosomal Deficits Augment Mitochondria Pathology in Spinal Motor Neurons of Asymptomatic fALS Mice. *Neuron*. 2015;87(2):355-370.
11. Otomo A, Pan L, Hadano S. Dysregulation of the autophagy-endolysosomal system in amyotrophic lateral sclerosis and related motor neuron diseases. *Neurol Res Int*. 2012;2012:498428.
12. Chia R, Chiò A, Traynor BJ. Novel genes associated with amyotrophic lateral sclerosis: diagnostic and clinical implications. *The Lancet Neurology*. 2018;17(1):94-102.

13. Offen D, Barhum Y, Melamed E, Embacher N, Schindler C, Ransmayr G. Spinal cord mRNA profile in patients with ALS: comparison with transgenic mice expressing the human SOD-1 mutant. *J Mol Neurosci*. 2009;38(2):85-93.

Neurology®

Clinical Reasoning: A 50-Year-Old Man With Progressive Limb Weakness and Slurred Speech

Nina Xie, Guang Yang, Weiru Zhang, et al.
Neurology published online February 10, 2022
DOI 10.1212/WNL.0000000000200008

This information is current as of February 10, 2022

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/early/2022/02/10/WNL.0000000000200008.citation.full
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Amyotrophic lateral sclerosis http://n.neurology.org/cgi/collection/amyotrophic_lateral_sclerosis_
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright © 2022 American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.