Neurology®

The most widely read and highly cited peer-reviewed neurology journal
The Official Journal of the American Academy of Neurology

Neurology Publish Ahead of Print
DOI: 10.1212/WNL.0000000000200014

Pearls & Oysters: Bilateral mononeuropathic neuralgic amyotrophy triggered by *Bartonella henselae* infection responsive to intravenous immunoglobulin

Author(s):

Anelia Dietmann, MD, PhD¹; Eugénie Colin-Benoit, MD²; Gerd Tinkhauser, MD, PhD¹; Thomas Raphael Meinel, MD¹; Olivier Scheidegger, MD¹

Corresponding Author:

Anelia Dietmann, anelia.dietmann@gmail.com

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

Neurology® Published Ahead of Print articles have been peer reviewed and accepted for publication. This manuscript will be published in its final form after copyediting, page composition, and review of proofs. Errors that could affect the content may be corrected during these processes.

Affiliation Information for All Authors: 1. Department of Neurology, Inselspital, Bern University Hospital, University of Bern, Switzerland; 2. Department of Infectious Diseases, Inselspital, Bern University Hospital, University of Bern, Switz, Switzerland.

Equal Author Contribution:

Contributions:

Anelia Dietmann: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data; Study concept or design

Eugénie Colin-Benoit: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data

Gerd Tinkhauser: Drafting/revision of the manuscript for content, including medical writing for content

Thomas Raphael Meinel: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data

Olivier Scheidegger: Drafting/revision of the manuscript for content, including medical writing for content; Major role in the acquisition of data

Figure Count:

2

Table Count:

0

Search Terms:

[285] EMG, [120] MRI, [139] Bacterial infections, [181] Peripheral neuropathy, Neuralgic amyotrophy

Acknowledgment:

Study Funding:

Anelia Dietmann has been personally supported by research grants from the Bangerter Rhyhner Stiftung and a CTU Research Grant of the University of Bern.

Disclosures:

Anelia Dietmann received funding from following institutions: Bangerter Rhyner Stiftung, University Bern, Medical Faculty; Eugenie.Colin-Benoit has nothing to declare; Gerd Tinkhauser has nothing to declare; Thomas Raphael Meinel has nothing to declare; Olivier Scheidegger has nothing to declare.

Abstract

We present a case of a cat-owner with a scar on his right thenar eminence, followed by lymphadenopathy in the right axilla, general malaise and fever, and subsequent onset of bilateral neuralgic amyotrophy within one week. After comprehensive workup, cat scratch disease caused by *Bartonella henselae* was confirmed serologically and adequately treated. Despite antibiotic treatment, he presented clinically with persistent bilateral, asymmetric neuropathy of the median nerve, predominantly the interosseous anterior nerve, which was confirmed by multifocal swelling and hyperintense signal of the nerves on T2-weighted MR neurography. Electrophysiological examination confirmed axonal median neuropathies bilaterally. After an unsuccessful steroid treatment trial, he showed an excellent and sustained response to intravenous immunoglobulin despite a delay from symptom-onset to treatment of 10 months.

Pearls

- *Bartonella henselae* infection is a rare cause of bilateral neuralgic amyotrophy.

Oy-sters

- Patients with neuralgic amyotrophy, especially when bilateral, need a comprehensive workup to exclude rare infectious causes such as Hepatitis E and, in case of suggestive history, *Bartonella henselae* infection.
- An intravenous immunoglobulin treatment trial may be considered, even in cases with poor clinical prognosis.

Case report

A 50-year-old cat-owner with an unremarkable medical history recognized a small scratch injury on his right thenar region, without noticing a cat scratch or an insect bite. After 3 days, his thenar eminence became swollen, tender and painful with expanding redness around the wound. He incised the wound with secretion of purulent liquid. After another 2-3 days, he developed febrile temperatures and noticed swollen but painless lymph nodes in the right axilla. Inflammatory markers were slightly elevated (CRP 18.5mg/L, normal <5mg/L; leucocytes 14.7G/L, normal range 3.0-10.5G/L; with normal differential count). An oral therapy with amoxicillin-clavulanic acid 3x625mg/d was prescribed empirically by his general practitioner. Two days later he noticed electric sensations radiating anteriorly from his right elbow to his lower arm and hand for several hours, followed by pain in his hand and paresthesia in his volar thumb and index finger. Within the next two days, he noted numbness in his distal thumb and index fingers bilaterally with evolving inability to flex the distal interphalangeal joints. Furthermore, he noticed patchy numbness over

his right ventral thigh. The general practitioner took blood cultures and a serology for *B. henselae*. Blood cultures were negative but IgG titers (by indirect fluorescence assay) were highly elevated ($> 1:1024$), IgM negative. Antibiotic therapy was changed to azithromycin 500mg/d on the first day followed by 250mg/d for four further days. After one week with persistent neurological deficits, *Bartonella* serology was repeated showing even higher IgG titers of $> 1:4096$, IgM again negative. Twelve days after starting antibiotic treatment (9 days after starting azithromycin), a lumbar puncture was performed with normal cell count (4 mononuclear cells/ μ L) and normal values for protein (316mg/L, normal 150-450mg/L), glucose and lactate. However, two CSF-specific oligoclonal bands were detected without signs of blood brain barrier disturbance. A CSF culture remained negative as well as *Bartonella* polymerase chain reaction (PCR). Lymph node biopsy revealed an expansion of the interfollicular zone with isolated B plasma cells without signs of lymphoma, granulomas, or *Bartonella* detected by PCR. Serology tests for *Toxoplasma gondii*, *Treponema pallidum*, *Borrelia burgdorferi*, HIV, Hepatitis B, C and E viruses were negative. No vasculitic antibodies or ganglioside antibodies were detected in the serum.

One month after onset, the patient was referred to our center for neuromuscular diseases. On examination, the patient was unable to feel light touch, temperature and pin-prick in the volar right distal thumb and index finger and the left distal thumb. He had a Medical Research Council's scale (MRC) muscle strength grade M0 for flexion of the distal interphalangeal joints of both thumbs (Figure 1). Flexion of the distal index finger was M0 on the right side and M4 on the left side. The muscle strength for supination of the right hand was M4. On the right thigh, he indicated decreased sensation for light touch in a patchy distribution, not corresponding to a specific nerve. Deep tendon reflexes were brisk and symmetric in all extremities.

Motor nerve conduction study (NCS) of the median nerve with recording from the abductor pollicis brevis muscle showed slightly prolonged distal motor latency (right 4.4 ms, left 5.2ms; normal < 4.0ms) with normal amplitudes and normal conduction velocity of proximal segments on both sides. Sensory NCS of the median nerve was unremarkable. Motor and sensory NCS of the ulnar nerve on both sides were normal. Needle myography of the right flexor pollicis longus muscle and pronator quadratus muscle showed a prolonged insertional activity and moderate fibrillation potentials and positive sharp waves without detection of volitional motor activity. Needle myography of the right abductor pollicis brevis muscle and interosseous dorsalis I muscle were unremarkable.

At 6 months after disease onset a MR neurography (3 T, Magnetom SkyraFit, Siemens, Erlangen, Germany; spine and body coils used) of the right brachial plexus and arm nerves were performed using high-resolution T2-weighted, fat-suppressed sequences. The exam revealed irregular hyperintense signal abnormalities and short segments of enlarged nerve fascicle bundles of the right median nerve in the middle and distal upper arm, corresponding somatotopically to nerve fascicle bundles of the anterior interosseous nerve and proximal muscular branches of the median nerve (Figure 2). Signs of muscle denervation (diffuse hyperintense signal abnormalities) were found in the pronator teres, flexor carpi radialis, and flexor digitorum profundus muscles. The nerve structure and signal intensity of the plexus brachialis as well as the distal segments of the median nerve below the elbow were normal, all other arm nerves were without pathological imaging findings.

A treatment trial of oral prednisolone (1mg/kg body weight per day for 1 month) one month after symptom onset was unsuccessful. Another treatment trial with intravenous immunoglobulin (IVIG; 0.4mg/kg body weight over 5 days every 4 weeks)

was started 10 months after onset due to difficulty in obtaining reimbursement of IVIG treatment cost for off-label use. The patient showed a significant clinical response within the first two weeks after treatment resulting in only mild residual weakness of finger flexion of distal interphalangeal joints of thumbs (right M4, left M5), and index fingers (right M5-, left M5), without any residual sensory disturbance. After another 5 cycles of IVIG the patient achieved an almost complete recovery, showing only a minimal deficit in the flexion of the right distal thumb. The electrophysiological follow-up assessment revealed a normalization of the motor NCS of the median nerve bilaterally. Therefore, we are now starting a tentative treatment pause.

Discussion

Infections are suspected to cause or trigger neuralgic amyotrophy (NA) in a significant proportion of patients¹. In the last 10 years it has been increasingly recognized that acute hepatitis E virus infection was associated with NA, especially bilateral NA, accounting for up to 10% of cases in recent series². Case reports of NA in patients with acute cat-scratch disease (CSD) have also been described in literature³.

CSD is caused by the gram-negative intracellular bacteria *Bartonella henselae* and usually affects children and teenagers, typically presenting as a localized cutaneous infection followed by adjacent lymphadenopathy and fever⁴. Neurological involvement has been described in 10% of CSD patients, presenting with encephalopathy, meningitis, myelitis, polyradiculoneuritis, neuroretinitis and isolated optic neuritis^{5,6}.

Because isolation of *Bartonella* species in culture is difficult (long incubation period, specific laboratory conditions), current diagnostic workup is usually based on serologic testing, sometimes combined with PCR⁷. The indirect fluorescence assay

(IFA) is the most frequently used serologic method. In the early stages of the disease, titers to IgG and IgM may be low, requiring a second serum sample at a later date for diagnosis. A positive IgM test strongly suggests acute disease but the production of IgM is usually brief and negative results do not exclude acute infection. IFA IgG titers >1:256 strongly suggest active or recent infection⁴. PCR for *Bartonella* is available and provides the advantages of high specificity and rapid identification, the sensitivity however ranges from approximately 65 to 76% in lymph node specimens, depending on PCR targets and selection of patients^{8,9}. In this case, the combination of typical clinical presentation and positive serology confirmed the diagnosis of *B. henselae* infection.

The mechanism by which infections trigger NA are largely uncertain and understudied. Similar to Guillain-Barré Syndrome, infections may trigger an autoimmune response that causes damage to the peripheral nervous system¹. In our case, as well as in previously described cases, the absence of improvement of NA symptoms with antibiotic treatment is consistent with an immune-mediated reaction to CSD preceding NA³.

Our patient developed an asymmetric, bilateral NA affecting mainly the median nerves, predominantly the fascicles of the anterior interosseous nerve bilaterally, within the first week of a typical infection with *B. henselae*. Interestingly, despite the electrophysiological finding of severe axonal damage of the interosseous anterior nerve, an unsuccessful steroid therapy trial, and a long delay between diagnosis and treatment, the prompt response to IVIG within the first two weeks after the first treatment cycle was excellent. The patient, who was previously severely limited in his work as an electrical engineer, was able to fully resume his work after IVIG treatment. A tentative IVIG cessation is the next step. However, if the patient deteriorates after

stopping IVIG treatment, the diagnosis would need to be changed to possible multifocal chronic inflammatory polyneuropathy triggered by the *Bartonella henselae* infection, with definite electrophysiological criteria not fulfilled¹⁰.

In conclusion, we present a case of bilateral mononeuropathic neuralgic amyotrophy triggered by *Bartonella henselae* infection responsive to IVIG treatment. It is essential to search for infectious causes or triggers of NA. In case of persisting symptoms after treatment of antecedent infection, IVIG treatment may be an option, even in cases with poor clinical prognosis, considering the probable immune-mediated cause of disease and apparent response to IVIG.

1. Van Eijk JJJ, Groothuis JT, Van Alfen N. Neuralgic amyotrophy: An update on diagnosis, pathophysiology, and treatment. *Muscle and Nerve*. 2016;53:337–350.
2. Ripellino P, Pasi E, Melli G, et al. Neurologic complications of acute hepatitis E virus infection. *Neurol Neuroimmunol neuroinflammation*. 2020;7.
3. Stek CJ, van Eijk JJJ, Jacobs BC, et al. Neuralgic amyotrophy associated with *Bartonella henselae* infection. *J Neurol Neurosurg Psychiatry* [online serial]. 2011;82:707–708. Accessed at: <https://jnnp.bmj.com/lookup/doi/10.1136/jnnp.2009.191940>.
4. Florin TA, Zaoutis TE, Zaoutis LB. Beyond Cat Scratch Disease: Widening Spectrum of *Bartonella henselae* Infection. *Pediatrics* [online serial]. 2008;121:e1413–e1425. Accessed at: <http://pediatrics.aappublications.org/cgi/doi/10.1542/peds.2007-1897>.
5. Canneti B, Cabo-López I, Puy-Núñez A, et al. Neurological presentations of *Bartonella henselae* infection. *Neurol Sci* [online serial]. 2019;40:261–268. Accessed at: <http://link.springer.com/10.1007/s10072-018-3618-5>.
6. Carithers HA. Cat-scratch disease. An overview based on a study of 1,200 patients. *Am J Dis Child* [online serial]. 1985;139:1124. Accessed at: <http://archpedi.jamanetwork.com/article.aspx?doi=10.1001/archpedi.1985.02140130062031>.
7. Allizond V, Costa C, Sidoti F, et al. Serological and molecular detection of *Bartonella henselae* in specimens from patients with suspected cat scratch disease in Italy: A comparative study. Munderloh UG, editor. *PLoS One* [online serial]. 2019;14:e0211945. Accessed at: <https://dx.plos.org/10.1371/journal.pone.0211945>.
8. Hansmann Y, DeMartino S, Piémont Y, et al. Diagnosis of Cat Scratch Disease with Detection of *Bartonella henselae* by PCR: a Study of Patients with Lymph Node Enlargement. *J Clin Microbiol* [online serial]. American Society for Microbiology (ASM); 2005;43:3800. Accessed at: </pmc/articles/PMC1233974/>. Accessed October 21, 2021.
9. Sander A, Posselt M, Böhm N, Ruess M, Altwegg M. Detection of *Bartonella henselae* DNA by two different PCR assays and determination of the genotypes of strains involved in histologically defined cat scratch disease. *J Clin Microbiol* [online serial]. American Society for Microbiology (ASM); 1999;37:993–997. Accessed at: </pmc/articles/PMC88638/>. Accessed October 21, 2021.
10. Van den Bergh PYK, van Doorn PA, Hadden RDM, et al. European Academy of Neurology/Peripheral Nerve Society guideline on diagnosis and treatment of chronic inflammatory demyelinating polyradiculoneuropathy: Report of a joint Task Force—Second revision. *J Peripher Nerv Syst*. 2021;26:242–268.

Figure 1 Clinical presentation. Patient was asked to make a fist (A) and to flex the proximal and distal interphalangeal joints of his thumb (B) He was unable to flex both distal interphalangeal joints of digit I and II (MRC grad 0),

Figure 2 MRI High-resolution T2-weighted, fat-suppressed sequences of the right brachial plexus and arm nerves. A) Irregular hyperintense signal abnormalities and short segments of enlarged nerve fascicle bundles of the right median nerve in the right proximal upper arm (red arrow) and B) normal appearing median nerve (red arrow) in the right lower arm with signs of muscle denervation (diffuse hyperintense signal abnormalities) in the right pronator teres muscle (a), flexor carpi radialis muscle (b), and flexor digitorum profundus muscle (c).

Neurology®

Pearls & Oy-sters: Bilateral mononeuropathic neuralgic amyotrophy triggered by Bartonella henselae infection responsive to intravenous immunoglobulin

Anelia Dietmann, EugÃ©nie Colin-Benoit, Gerd Tinkhauser, et al.

Neurology published online February 10, 2022

DOI 10.1212/WNL.0000000000200014

This information is current as of February 10, 2022

Updated Information & Services	including high resolution figures, can be found at: http://n.neurology.org/content/early/2022/02/10/WNL.0000000000200014.full
Subspecialty Collections	This article, along with others on similar topics, appears in the following collection(s): Bacterial infections http://n.neurology.org/cgi/collection/bacterial_infections EMG http://n.neurology.org/cgi/collection/emg MRI http://n.neurology.org/cgi/collection/mri Peripheral neuropathy http://n.neurology.org/cgi/collection/peripheral_neuropathy
Permissions & Licensing	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: http://www.neurology.org/about/about_the_journal#permissions
Reprints	Information about ordering reprints can be found online: http://n.neurology.org/subscribers/advertise

Neurology® is the official journal of the American Academy of Neurology. Published continuously since 1951, it is now a weekly with 48 issues per year. Copyright Copyright © 2022 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the American Academy of Neurology. All rights reserved. Print ISSN: 0028-3878. Online ISSN: 1526-632X.